



WHO recommendations for care of the preterm or low-birth-weight infant



World Health
Organization

WHO recommendations for care of the preterm or low-birth-weight infant

WHO recommendations for care of the preterm or low birth weight infant

ISBN 978-92-4-005826-2 (electronic version)

ISBN 978-92-4-005827-9 (print version)

© **World Health Organization 2022**

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

Under the terms of this licence, you may copy, redistribute and adapt the work for non-commercial purposes, provided the work is appropriately cited, as indicated below. In any use of this work, there should be no suggestion that WHO endorses any specific organization, products or services. The use of the WHO logo is not permitted. If you adapt the work, then you must license your work under the same or equivalent Creative Commons licence. If you create a translation of this work, you should add the following disclaimer along with the suggested citation: "This translation was not created by the World Health Organization (WHO). WHO is not responsible for the content or accuracy of this translation. The original English edition shall be the binding and authentic edition".

Any mediation relating to disputes arising under the licence shall be conducted in accordance with the mediation rules of the World Intellectual Property Organization (<http://www.wipo.int/amc/en/mediation/rules/>).

Suggested citation. WHO recommendations for care of the preterm or low birth weight infant. Geneva: World Health Organization; 2022. Licence: CC BY-NC-SA 3.0 IGO.

Cataloguing-in-Publication (CIP) data. CIP data are available at <http://apps.who.int/iris>.

Sales, rights and licensing. To purchase WHO publications, see <http://apps.who.int/bookorders>. To submit requests for commercial use and queries on rights and licensing, see <https://www.who.int/copyright>.

Third-party materials. If you wish to reuse material from this work that is attributed to a third party, such as tables, figures or images, it is your responsibility to determine whether permission is needed for that reuse and to obtain permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

General disclaimers. The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of WHO concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by WHO in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by WHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall WHO be liable for damages arising from its use.

Design and layout by Green Ink Publishing Services Ltd.

Cover photograph: © WHO / Blink Media - Hilina Abebe

Contents

Acknowledgements	v
Acronyms and abbreviations	vi
Glossary	vii
Executive summary	viii
1. Introduction	1
2. Methods	5
3. Evidence and recommendations	11
A. Preventive and promotive care	12
A.1 Kangaroo mother care	12
A.2 Mother's own milk	17
A.3 Donor human milk	20
A.4 Multicomponent fortification of human milk	23
A.5 Preterm formula	27
A.6 Early initiation of enteral feeding	30
A.7 Responsive and scheduled feeding	33
A.8 Fast and slow advancement of feeding	36
A.9 Duration of exclusive breastfeeding	39
A.10 Micronutrient supplementation	42
A.11 Probiotics	58
A.12 Emollients	62
B. Care for complications	66
B.1 Continuous positive airway pressure for respiratory distress syndrome	66
B.2 Continuous positive airway pressure immediately after birth	70
B.3 Continuous positive airway pressure source	73
B.4 Methylxanthines for treatment of apnoea	76
B.5 Methylxanthines for extubation	79
B.6 Methylxanthines for prevention of apnoea	81
C. Family involvement and support	83
C.1 Family involvement in routine care	83
C.2 Family support	86
C.3 Home visits	91
C.4 Parental leave and entitlements	94
4. Implementation	97
5. Applicability issues	99
6. Research implications	100
7. Dissemination	102
8. Monitoring and evaluating the impact of the recommendations	103
9. Updating of the guideline	104
References	105
Annexes	118
Annex 1: Current WHO recommendations for care of the preterm or low-birth-weight infant	118
Annex 2: Contributors to the guideline	120
Annex 3: Summary of declarations of interest from the Guideline Development Group (GDG) members and how they were managed	122

Web Annexes

Web Annex A. Priority questions and outcomes

Web Annex B. Detailed list of research priorities

Web Annex C. Changes from approved scope of guideline

The web annexes are available at:

<https://apps.who.int/iris/bitstream/handle/10665/363698/9789240060043-eng.pdf>

Web Supplement. Evidence base

The web supplement is available at:

<https://apps.who.int/iris/bitstream/handle/10665/363699/9789240060050-eng.pdf>

Acknowledgements

The World Health Organization (WHO) Department of Maternal, Newborn, Child and Adolescent Health and Ageing (MCA) gratefully acknowledges the contributions that many individuals and organizations have made to the development of these recommendations.

WHO extends sincere thanks to the chairs of the Guideline Development Group (GDG) Gary Darmstadt and Vinod Paul, and to all the other GDG members: Nafisa Hamoud Al Jaifi, Shabina Ariff, Mats Blennow, Liz Comrie-Thomson, Socorro De Leon-Mendoza, Qi Feng, Patricia Fernandez Riera, Zelee Hill, Ayede Adejumoke Idowu, Carole Kenner, Victoria Nakibuuka Kirabira, Reinhard Klinkott, Silke Mader, Karim Manji, Rhonda Marriot, Monica Morgues, Sushma Nangia, Mohammad Shahidullah, Hoang Thi Tran, Rizwan Ullah Hassan, Andrew Weeks, Bogale Worku and Khalid Yunis. WHO also thanks the members of the External Review Group: Alexander Manu, Ebunoluwa Adejuyigbe, Carolyn MacLennan and Abiy Seifu Estifanos.

Roger Chou was the external guideline methodologist. Natalie Strobel provided technical support for analysis, preparation of Grading of Recommendations Assessment, Development and Evaluation (GRADE) tables and guideline content. Special thanks go to the authors of the systematic reviews, including the Cochrane Neonatal Group.

WHO acknowledges the observers at the GDG meetings representing partner organizations, including the Bill & Melinda Gates Foundation (Hema Magge), the International Pediatric Association (William Keenan), the United States Agency for International Development (USAID) (Lily Kak) and Save the Children (Steve Wall). WHO also appreciates the contributions of Gagan Gupta and Tedbabe Degefe Hailegebriel of UNICEF.

The WHO Steering Group coordinated the guideline development process: Rajiv Bahl, Vanessa Cavallera, Doris Chou, Karen Edmond (responsible technical officer), Larry Grummer Strawn, Shuchita Gupta, Anayda Portela and Suman Rao. WHO regional advisers who contributed to the guideline include Jamela Al-Raiby, Rajesh Mehta, Assumpta Muriithi, Khalid Siddeeg, Howard Sobel, Martin Weber and Teshome Desta Woldehanna. Other WHO staff who provided inputs on the guideline content include Ayesha Da Costa and Mercedes Bonet Semenas.

USAID funded this work. The views of the funding body have not influenced the content of these recommendations.

Acronyms and abbreviations

BSID	Bayley Scales of Infant and Toddler Development (edition indicated as BSID-II or BSID-III)	MD	mean difference
CERQual	Confidence in the Evidence from Reviews of Qualitative Research	MDI	Mental Development Index (BSID-II)
CI	confidence interval	MSD	Department of Mental Health and Substance Use (at WHO)
CPAP	continuous positive airway pressure	M-NICU	maternal–neonatal intensive care unit
DECIDE	Developing and Evaluating Communication strategies to support Informed Decisions and practice based on Evidence	NFS	Department of Nutrition and Food Safety (at WHO)
DOI	declaration of interests	NICU	neonatal intensive care unit
EBF	exclusive breastfeeding	OR	odds ratio
ERG	External Review Group	PDI	Psychomotor Development Index (BSID-II)
EST	Evidence Synthesis Team	PICO	population, intervention, comparator, outcome
EtD	Evidence-to-Decision	PMA	postmenstrual age
GDG	Guideline Development Group	PNC	postnatal care
GRADE	Grading of Recommendations Assessment, Development and Evaluation	RCT	randomized controlled trial
HAZ	height-for-age z score	RR	relative risk
IFD	Infant Flow Driver	SD	standard deviation
IQR	interquartile range	SGA	small for gestational age
IPA	International Pediatric Association	SMD	standardized mean difference
IU	international unit	SRH	Department of Sexual and Reproductive Health and Research (at WHO)
KMC	kangaroo mother care	spp.	several species
LAZ	length-for-age z score	UNICEF	United Nations Children's Fund
LMIC	low- or middle-income country	US\$	United States dollar
LMP	last menstrual period	USAID	United States Agency for International Development
MCA	Department of Maternal, Newborn, Child and Adolescent Health and Ageing (at WHO)	WAZ	weight-for-age z score
		WHO	World Health Organization
		WHZ	weight-for-height z score

Glossary

Low birth weight (LBW)	birth weight below 2.5 kg
Very LBW (VLBW)	birth weight below 1.5 kg
Extremely LBW	birth weight below 1 kg
Term gestation	born at 37 0/7 – 41 6/7 weeks of gestation (i.e. since mother's last menstrual period [LMP])
Preterm	born before 37 0/7 weeks of gestation
Very preterm	born before 32 0/7 weeks of gestation
Extremely preterm	born before 28 0/7 weeks of gestation
Post-term	born at or after 42 0/7 weeks of gestation
Chronological (or postnatal) age	age since birth (e.g. an infant born 10 weeks ago at 32 weeks' gestation is 10 weeks of age/chronological age/postnatal age)
Corrected age	chronological age minus the number of weeks or months born preterm (with term defined as 40 weeks, for the purpose of calculation) (e.g. an infant born at 32 weeks' gestation, who is 10 weeks old in chronological age, is only 2 weeks old in corrected age)
Postmenstrual age (PMA)	the age of a baby or fetus when counted from the first day of the mother's LMP before pregnancy (e.g. a baby that was born at 32 weeks' gestation, who is 10 weeks old in chronological age, is 42 weeks in PMA)
Stunting	length-for-age z score less than 2 standard deviation scores below the WHO child growth standards median
Underweight	weight-for-age z score less than 2 standard deviation scores below the WHO child growth standards median
Wasting	weight-for-length z score less than 2 standard deviation scores below the WHO child growth standards median

Sources:

Preterm birth. Geneva: World Health Organization; 2018 (<https://www.who.int/news-room/fact-sheets/detail/preterm-birth>, accessed 14 September 2022).

Low birth weight. In: Nutrition Landscape Information System (NLIS) [website]. Geneva: World Health Organization; 2022 (<https://www.who.int/data/nutrition/nlis/info/low-birth-weight>, accessed 14 September 2022).

WHO Global Database on Child Growth and Malnutrition. Geneva: World Health Organization; 2022 (<https://platform.who.int/nutrition/malnutrition-database>, accessed 31 October 2022).

Child growth standards [website]. Geneva: World Health Organization; no date (<https://www.who.int/tools/child-growth-standards/standards>, accessed 5 October 2022).

Executive summary

Introduction

Preterm infants have a gestational age below 37 weeks at birth and low-birth-weight (LBW) infants have a birth weight below 2.5 kg. Approximately 45% of all children under the age of five who die are newborns, and 60–80% of those newborns who die are premature and/or small for gestational age. Preterm and LBW infants have a 2- to 10-fold higher risk of mortality than infants born at term and with normal birth weight. Despite substantial progress over the last 10 years, the survival, health, growth and neurodevelopment of preterm and LBW infants remains concerning in many countries. Reasons include the complexities of caring for these vulnerable infants and preventing complications.


The care of preterm and LBW infants is a global priority. The WHO Departments of Maternal, Newborn, Child and Adolescent Health and Ageing (MCA) and Sexual and Reproductive Health and Research (SRH) have developed three guidelines for the care of preterm or LBW infants:

- *Guidelines on optimal feeding of low-birth-weight infants in low- and middle-income countries*, 2011;
- *WHO recommendations on interventions to improve preterm birth outcomes*, 2015; and
- *Recommendations for management of common childhood conditions*, 2012.

However, new evidence has emerged in many areas since the development of those guidelines. A review of 203 studies from low-, middle- and high-income countries about “what matters” to families about the care of their preterm or LBW infant reported that families want a positive outcome for their baby, to be involved in delivering care and to take an active role in deciding what interventions are given to their baby (these family values are listed in Table 1.1 in Chapter 1).

In December 2020, an international group of experts defined the scope and priority questions for the development of updated guidance about the care of preterm or LBW infants (see Figure 1 below, Table 1.2 in Chapter 1, and Web Annex A).

Figure 1. Scope of WHO recommendations for care of the preterm or low-birth-weight infant



Target audience

The recommendations in this guideline are intended to inform the development of national and subnational health policies, clinical protocols and programmatic guides. Therefore, the target audience includes national and subnational public health policy-makers, implementers and managers of maternal, newborn and child health programmes, health-care facility managers, supervisors/instructors for in-service training, health workers (including midwives, auxiliary nurse-midwives, nurses, paediatricians, neonatologists, general medical practitioners and community health workers), nongovernmental organizations, professional societies involved in the planning and management of maternal, newborn and child health services, academic staff involved in research and in the pre-service education and training of health workers, and those involved in the education of parents.

Guideline development methods

The guideline was developed using standard operating procedures in accordance with the process described in the *WHO handbook for guideline development*. This involved the convening of an Evidence Synthesis Team (EST) and an international Guideline Development Group (GDG) of experts. The process included: (i) identifying priority questions and outcomes, (ii) retrieval of the evidence, (iii) assessment and synthesis of the evidence, (iv) formulation of recommendations and write-up of the guideline, and (v) planning for dissemination, implementation, impact evaluation and updating of the recommendations.

The Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach was used to appraise the quality and certainty of the quantitative evidence for each priority question, and for the qualitative evidence, the reviews were appraised using the GRADE-CERQual (Confidence in the Evidence from Reviews of Qualitative Research) tool. The DECIDE approach (Developing and Evaluating Communication strategies to support Informed Decisions and practice based on Evidence), an evidence-to-decision tool, was used to guide the evidence search, evidence synthesis and judgements on the different criteria by the EST, and the formulation of recommendations by the GDG. This included assessment of the effects (benefits and harms) of the interventions on infant outcomes, and consideration of the values of families and health workers, acceptability and feasibility of the interventions, the resources required, and equity.

Recommendations were developed using WHO Guidelines Review Committee criteria: “strong” recommendations are generally applicable to all preterm or LBW infants; “conditional” recommendations mean that the intervention is recommended under certain conditions; and a “good practice statement” was made for an intervention that was obviously beneficial and should be done in most circumstances, even though there was no, little or only very-low-certainty evidence. The GDG members examined and interpreted the evidence, formulated the wording of the final recommendations and provided related remarks and considerations at virtual meetings held between November 2021 and January 2022.

Recommendations

This guideline includes 25 recommendations and 1 good practice statement (see Table 1) for care of the preterm and LBW infant. Of the recommendations, 11 are new and 14 are updated, and the good practice statement is new. There are 11 strong recommendations for all preterm or LBW infants and 14 recommendations that are conditional on particular contexts or conditions.

Sixteen recommendations are for preventive and promotive care, six are for care for complications and three are for family involvement and support. A good practice statement was made for parental leave and entitlements because the GDG determined that these have obvious benefits, although there was little evidence available.

The GDG provided remarks related to all the recommendations and the good practice statement, where needed. Users of the guideline should refer to these remarks, which are presented prominently, along with the recommendations in Chapter 3 of the guideline.

Table 1. WHO recommendations for the care of the preterm (< 37 weeks' gestation) or low-birth-weight (< 2.5 kg) infant

Domain	Recommendation	Status	Strength/type
A. PREVENTIVE AND PROMOTIVE CARE			
A.1a Any KMC	Kangaroo mother care (KMC) is recommended as routine care for all preterm or low-birth-weight infants. KMC can be initiated in the health-care facility or at home and should be given for 8–24 hours per day (as many hours as possible). (<i>Strong recommendation, high-certainty evidence</i>)	Updated	Strong
A.1b Immediate KMC	Kangaroo mother care (KMC) for preterm or low-birth-weight infants should be started as soon as possible after birth. (<i>Strong recommendation, high-certainty evidence</i>)	New	Strong
A.2 Mother's own milk	Mother's own milk is recommended for feeding of preterm or low-birth-weight (LBW) infants, including very preterm (< 32 weeks' gestation) or very LBW (< 1.5 kg) infants. (<i>Strong recommendation, low-certainty evidence</i>)	Updated	Strong
A.3 Donor human milk	When mother's own milk is not available, donor human milk may be considered for feeding of preterm or low-birth-weight (LBW) infants, including very preterm (< 32 weeks' gestation) or very LBW (< 1.5 kg) infants. (<i>Conditional recommendation, moderate-certainty evidence</i>)	Updated	Conditional
A.4 Multicomponent fortification of human milk	Multicomponent fortification of human milk is not routinely recommended for all preterm or low-birth-weight (LBW) infants but may be considered for very preterm (< 32 weeks' gestation) or very LBW (< 1.5 kg) infants who are fed mother's own milk or donor human milk. (<i>Conditional recommendation, low-to-moderate-certainty evidence</i>)	Updated	Conditional
A.5 Preterm formula	When mother's own milk and donor human milk are not available, nutrient-enriched preterm formula may be considered for very preterm (< 32 weeks' gestation) or very low-birth-weight (< 1.5 kg) infants. (<i>Conditional recommendation, low-certainty evidence</i>)	Updated	Conditional
A.6 Early initiation of enteral feeding	Preterm and low-birth-weight (LBW) infants, including very preterm (< 32 weeks' gestation) and very LBW (< 1.5 kg) infants, should be fed as early as possible from the first day after birth. Infants who are able to breastfeed should be put to the breast as soon as possible after birth. Infants who are unable to breastfeed should be given expressed mother's own milk as soon as it becomes available. If mother's own milk is not available, donor human milk should be given wherever possible. (<i>Strong recommendation, moderate-certainty evidence</i>)	Updated	Strong
A.7 Responsive and scheduled feeding	In health-care facilities, scheduled feeding may be considered rather than responsive feeding for preterm infants born before 34 weeks' gestation, until the infant is discharged. (<i>Conditional recommendation, low-certainty evidence</i>)	Updated	Conditional
A.8 Fast and slow advancement of feeding	In preterm or low-birth-weight (LBW) infants, including very preterm (< 32 weeks' gestation) or very LBW (< 1.5 kg) infants, who need to be fed by an alternative feeding method to breastfeeding (e.g. gastric tube feeding or cup feeding), feed volumes can be increased by up to 30 ml/kg per day. (<i>Conditional recommendation, moderate-certainty evidence</i>)	Updated	Conditional

Domain	Recommendation	Status	Strength/type
A.9 Duration of exclusive breastfeeding	Preterm or low-birth-weight infants should be exclusively breastfed until 6 months of age. (<i>Strong recommendation, very-low-certainty evidence</i>)	Updated	Strong
A.10a Iron supplementation	Enteral iron supplementation is recommended for human milk-fed preterm or low-birth-weight infants who are not receiving iron from another source. (<i>Strong recommendation, moderate-certainty evidence</i>)	Updated	Strong
A.10b Zinc supplementation	Enteral zinc supplementation may be considered for human milk-fed preterm or low-birth-weight infants who are not receiving zinc from another source. (<i>Conditional recommendation, low-certainty evidence</i>)	Updated	Conditional
A.10c Vitamin D supplementation	Enteral vitamin D supplementation may be considered for human milk-fed preterm or low-birth-weight infants who are not receiving vitamin D from another source. (<i>Conditional recommendation, low-certainty evidence</i>)	Updated	Conditional
A.10d Vitamin A supplementation	Enteral vitamin A supplementation may be considered for human milk-fed very preterm (< 32 weeks' gestation) or very low-birth-weight (< 1.5 kg) infants who are not receiving vitamin A from another source. (<i>Conditional recommendation, low-certainty evidence</i>)	Updated	Conditional
A.11 Probiotics	Probiotics may be considered for human-milk-fed very preterm infants (< 32 weeks' gestation). (<i>Conditional recommendation, moderate-certainty evidence</i>)	New	Conditional
A.12 Emollients	Application of topical oil to the body of preterm or low-birth-weight infants may be considered. (<i>Conditional recommendation, low-certainty evidence</i>)	New	Conditional

B. CARE FOR COMPLICATIONS

B.1 CPAP for respiratory distress syndrome	Continuous positive airway pressure (CPAP) therapy is recommended in preterm infants with clinical signs of respiratory distress syndrome. (<i>Strong recommendation, moderate-certainty evidence</i>)	Updated	Strong
B.2 CPAP immediately after birth	Continuous positive airway pressure (CPAP) therapy may be considered immediately after birth for very preterm infants (< 32 weeks' gestation), with or without respiratory distress. (<i>Conditional recommendation, low-certainty evidence</i>)	New	Conditional
B.3 CPAP pressure source (bubble CPAP)	For preterm infants who need continuous positive airway pressure (CPAP) therapy, bubble CPAP may be considered rather than other pressure sources (e.g. ventilator CPAP). (<i>Conditional recommendation, low-certainty evidence</i>)	New	Conditional
B.4 Methylxanthines for treatment of apnoea	Caffeine is recommended for the treatment of apnoea in preterm infants. (<i>Strong recommendation, moderate-certainty evidence</i>)	New	Strong
B.5 Methylxanthines for extubation	Caffeine is recommended for the extubation of preterm infants born before 34 weeks' gestation. (<i>Strong recommendation, moderate-certainty evidence</i>)	New	Strong
B.6 Methylxanthines for prevention of apnoea	Caffeine may be considered for the prevention of apnoea in preterm infants born before 34 weeks' gestation. (<i>Conditional recommendation, low-certainty evidence</i>)	New	Conditional

Domain	Recommendation	Status	Strength/ type
C. FAMILY INVOLVEMENT AND SUPPORT			
C.1 Family involvement	Family involvement in the routine care of preterm or low-birth-weight infants in health-care facilities is recommended. (<i>Strong recommendation, low- to moderate-certainty evidence</i>)	New	Strong
C.2 Family support	Families of preterm or low-birth-weight infants should be given extra support to care for their infants, starting in health-care facilities from birth and continued during follow-up post-discharge. The support may include education, counselling and discharge preparation from health workers, and peer support. (<i>Conditional recommendation, very-low-certainty evidence</i>)	New	Conditional
C.3 Home visits	Home visits by trained health workers are recommended to support families to care for their preterm or low-birth-weight infant. (<i>Strong recommendation, moderate-certainty evidence</i>)	New	Strong
C.4 Parental leave and entitlements	Parental leave and entitlements should address the special needs of mothers, fathers and other primary caregivers of preterm or low-birth-weight infants. (<i>Good practice statement</i>)	New	Good practice statement

1. Introduction

1.1 Background

Preterm infants have a gestational age below 37 weeks at birth and low-birth-weight (LBW) infants have a birth weight below 2.5 kg (1-3). Global estimates of prematurity and LBW range from 15% to 20%. In 2015, an estimated 20.5 million live births were LBW, with 91% of those being from low- and middle-income countries (LMICs), mainly in southern Asia (48%) and sub-Saharan Africa (24%) (1-3).

Approximately 45% of all children under the age of five who die are newborns (2.7 out of 5.9 million in 2014), and 60-80% of those newborns who die are premature and/or small for gestational age (4). Preterm and LBW infants have a 2- to 10-fold higher risk of mortality than infants born at term (at least 37 weeks' gestation) and with normal birth weight (at least 2.5 kg), and are particularly vulnerable to impaired respiration, difficulty feeding, growth failure,

poor body temperature regulation, and infection (5-7). Preterm and LBW infants have a higher risk of developmental disabilities, including cerebral palsy and retinopathy of prematurity, and long-term adult-onset chronic conditions such as cardiovascular disease (8,9).

The survival, health, growth and neurodevelopment of preterm and LBW infants remains concerning in many countries and the pace of improvement has been slow (10-13). Reasons include the complexities of caring for these vulnerable infants and preventing complications. A review of 203 studies from low-, middle- and high-income settings, about "what matters" to families about the care of preterm or LBW infants reported that families want a positive outcome for their baby, to be involved in providing care for their baby, and to take an active role in deciding what interventions are given to their baby (see values and preferences listed in Table 1.1) (14).

Table 1.1 Family values and preferences about the care of their preterm or low-birth-weight infant

Domain	Descriptor
Positive outcome	Positive outcome for the child
Active involvement in care	Delivering care
	Fathers and partners involved
	Opportunities for parenting
	Shared decision-making and consent
Coping at home	Accessing support in a crisis
	Autonomy
	Extended family support and community resources
	Health professional expertise in the community
	Preparation for discharge
	Transition arrangements
Emotional support for family	Support for all parents
	Additional support for mothers and fathers, acknowledging that they may have different emotional support needs
	Support from the wider family
	Support from other parents in similar situations

Domain	Descriptor
Health-care environment	Access to babies
	Orientation and familiarity with the neonatal intensive care unit (NICU)
	Balance between privacy and monitoring
	Staffing and equipment levels
Information needs met	Information about the baby
	Frequent updates
	How information is given
	Matching needs with information
Logistical support	Accommodation (comfort and facilities)
	Broader family support and impact
	Costs of treatment
	Parental leave
Positive relationships with staff	Compassion and sensitivity
	Consistency in care and communication
	Health professional expertise and care
	Respect, collaboration and trust

Source: Hurt et al., 2022 (14).

The care of the preterm and LBW infant is a global priority and a component of the United Nations *Every Woman Every Child* (EWEC) *Global Strategy for Women's, Children's and Adolescents' Health* 2016–2030 (4), the United Nations Children's Fund (UNICEF) *Every Child Alive* campaign (15), the World Health Organization (WHO) 2025 global nutrition targets (16), and the joint WHO-UNICEF *Every Newborn Action Plan* (ENAP) to end preventable deaths (17,18).

The WHO Departments of Maternal, Newborn, Child and Adolescent Health and Ageing (MCA) and Sexual and Reproductive Health and Research (SRH) have previously developed three guidelines for the care of preterm or LBW infants:

- *Guidelines on optimal feeding of low-birth-weight infants in low- and middle-income countries*, 2011 (19);
- *WHO recommendations on interventions to improve preterm birth outcomes*, 2015 (20); and
- *Recommendations for management of common childhood conditions*, 2012 (21).

However, new evidence has emerged in many areas since the development of those guidelines.

1.2 Target audience

The recommendations in this guideline are intended to inform the development of national and subnational health policies, clinical protocols and programmatic guides. Therefore, the target audience includes national and subnational public health policy-makers, implementers and managers of maternal, newborn and child health programmes, health-care facility managers, supervisors/instructors for in-service training, health workers (including midwives, auxiliary nurse-midwives, nurses, paediatricians, neonatologists, general medical practitioners and community health workers), nongovernmental organizations, professional societies involved in the planning and management of maternal, newborn and child health services, academic staff involved in research and in the pre-service education and training of health workers, and those involved in the education of parents.

1.3 Scope of the guideline

The recommendations cover the care of the preterm or LBW infant in any health-care facility or community setting from birth to 24 months of age unless otherwise indicated (see Table 1.2 and Figure 1.1). There are 25 recommendations and 1 good practice statement. They are summarized in Table 1 in the executive summary and presented in detail in Chapter 3. Of the recommendations, 11 are new and 14 are updated. There are 11 strong recommendations for all preterm or LBW infants and 14 recommendations that are conditional on particular contexts or conditions. Sixteen recommendations are for preventive and promotive


care (section A of Chapter 3), six are for care for complications (section B) and three are for family involvement and support (section C). A good practice statement was made for parental leave and entitlements because the GDG determined that these have obvious benefits, although there was little evidence available.

Other recommendations for care of the preterm or LBW infant (i.e. the items in Figure 1.1 that are in italics, not bold) are covered elsewhere, or will be included in a future update (see Annex 1). The recommendations presented here are also complementary to existing WHO guidelines for antenatal, intrapartum and postnatal care (20,22-24).

Table 1.2. Framework for the WHO recommendations for care of the preterm or low-birth-weight infant

Target population (P)	Preterm (<37 weeks' gestation) or low-birth-weight (LBW) (<2.5 kg) infants
Interventions (I)	A. Preventive and promotive care B. Care for complications C. Family involvement and support
Intervention period	From birth to 24 months of age
Comparators (C)	Usual care or no intervention
Comparator period	From birth to 24 months of age
Outcomes (O)	<ul style="list-style-type: none">▪ Critical outcomes: infant all-cause mortality, morbidity, growth, neurodevelopment at latest follow-up▪ Other outcomes: other infant outcomes that are specific for the intervention at latest follow-up
Outcome period	Unrestricted
Setting	Health-care facility or home, in any country or setting
Subgroups	<ul style="list-style-type: none">▪ Very preterm (< 32 weeks' gestation) or very LBW (< 1.5 kg)▪ Other, specific for the intervention

Figure 1.1 Scope of WHO recommendations for care of the preterm or low-birth-weight infant



2. Methods

This document was developed using the standard operating procedures described in the *WHO handbook for guideline development, second edition* (25). The process included: (i) identifying priority questions and outcomes, (ii) retrieval of the evidence, (iii) assessment and synthesis of the evidence, (iv) formulation of recommendations and write-up of the guideline, and (v) planning for the dissemination, implementation, impact evaluation and updating of the recommendations.

2.1 Contributors to the guideline

The groups involved in the development of the guideline are described below. The members of these groups are listed in Annex 2.

2.1.1 WHO Steering Group

The guideline development process was supervised by the WHO Steering Group, comprising staff members from four WHO departments: Maternal, Newborn, Child and Adolescent Health and Ageing; Mental Health and Substance Abuse; Nutrition and Food Safety; and Sexual and Reproductive Health and Research. The Steering Group drafted the initial scope of the guideline; identified priority questions in the “PICO” format (encompassing population, intervention, comparators and outcomes); prepared the guideline planning proposal; identified and invited systematic review teams, the guideline methodologist and members of the Guideline Development Group (GDG); supervised evidence retrieval, assessment and synthesis; organized the GDG meetings; prepared draft recommendations for the consideration of the GDG; compiled the final guideline document; and managed the guideline publication and dissemination.

2.1.2 Guideline Development Group (GDG)

The WHO Steering Group identified 25 external experts and stakeholders from the six WHO regions to form the GDG. Criteria included geographic representation, gender balance and no conflicts of interest. The final GDG was a diverse group of individuals with expertise in research, clinical practice, policy and programmes, guideline development methods and service delivery

approaches, including patient and consumer representatives.

The GDG participated in a virtual scoping meeting with the Steering Group in December 2020, and provided input on the PICO questions and related details that had been drafted to guide the evidence reviews. The GDG members examined and interpreted the evidence, formulated the wording of the final recommendations and provided related remarks and considerations at virtual GDG meetings between November 2021 and January 2022. The GDG also reviewed and approved the final guideline document.

2.1.3 External Review Group (ERG)

The ERG included four technical experts and stakeholders with expertise and experience in the provision of care for the preterm or LBW infant. The group was geographically representative and gender balanced. The ERG peer-reviewed the draft guideline document after the GDG had approved it. They assessed and provided feedback on: factual errors; clarity of language; guideline decision-making processes; values and preferences of persons affected by the recommendations (including families, health workers, managers and policy-makers); and the implications for implementation. It was not within the remit of this group to change recommendations that had been formulated by the GDG.

2.1.4 Evidence Synthesis Team (EST)

The EST comprised the guideline methodologist, systematic review teams and members of the WHO Steering Group. Within the EST, there were two work streams, each addressing multiple domains (see section 2.4). The work streams initially prepared an overview of systematic reviews (26) and a review of what matters to families about the care of their preterm or LBW infant (see Table 1.1) (14). They then appraised the quality of existing systematic reviews and commissioned new systematic reviews and structured searches. The EST members then reviewed each systematic review, prepared the Grading of Recommendations Assessment, Development and Evaluation (GRADE) Evidence-to-Decision (EtD) frameworks for each priority question and attended the GDG meetings.

2.1.5 External partners and observers

Representatives of the United Nations Children's Fund (UNICEF), Save the Children, the Bill & Melinda Gates Foundation, the International Pediatric Association (IPA) and the United States Agency for International Development (USAID) were invited to the GDG meetings as observers. These organizations are potential partners in the implementation of the guideline, as they have a history of collaboration with WHO in guideline dissemination and implementation. Observers were allowed to make comments during technical discussions at selected times during the GDG meetings. Observers did not participate in discussions on the final recommendations.

2.2 Declarations of interests by external contributors

In accordance with WHO procedures for declarations of interests (DOIs) (27), all GDG, EST and ERG members and other external collaborators were asked to declare in writing any competing interests (whether academic, financial or other), using the standard WHO DOI form, before engaging in the guideline development process. All experts were instructed to notify the responsible technical officer of any change in relevant interests, in order to update and review potential conflicts of interest accordingly. In addition, the GDG members were requested to submit an electronic copy of their curriculum vitae.

The names and short curriculum vitae of the GDG members were published on the WHO website for public review and comment two weeks prior to the first GDG meeting.

The WHO Steering Group reviewed all DOI forms and curriculum vitae to determine whether any conflicts of interest existed. All findings from the DOI forms were managed in accordance with the WHO DOI guidelines on a case-by-case basis. To ensure consistency, the Steering Group applied the criteria for assessing the severity of a conflict of interest in the *WHO handbook for guideline development* (25).

For this guideline, none of the declared interests were considered serious enough to pose any risk to the guideline development process or to reduce its credibility. Thus, all experts were only required

to declare such interests at the first GDG meeting. At each subsequent GDG meeting, GDG and EST members and observers were required to share any new potential conflicts of interest with the group.

Some GDG members had performed primary research related to one or more of the guideline recommendations. In these cases, the experts were restricted from participating in discussions or formulating any recommendations related to that specific area of interest. There were no important conflicts of interest among the ERG members.

A summary of the GDG DOIs and how conflicts of interest were managed is provided in Annex 3.

2.3 Identifying priority questions and outcomes

At the scoping meeting, the GDG decided on the priority questions in the PICO format (population, intervention, comparators, outcomes), based on the following criteria:

- values and preferences of families as outlined in the systematic review, "What matters to families about the care of their preterm or low-birth-weight (LBW) infant" (see Table 1.1) (14);
- public health importance;
- availability of new evidence; and
- questions not addressed by existing WHO guidelines or those identified for update.

The final scope of the guideline is presented in Table 1.2 and Figure 1.1. The PICO questions can be found in Web Annex A.

2.4 Evidence search, retrieval and review

The DECIDE approach (Developing and Evaluating Communication strategies to support Informed Decisions and practice based on Evidence) (28) was used to guide the evidence search, evidence synthesis and judgements by the EST, and the formulation of recommendations by the GDG. The DECIDE framework has nine core domains: benefits, harms, balance of effects, certainty, values, acceptability, resources, feasibility and equity (Table 2.2).

Table 2.2 Evidence-to-Decision (EtD) framework workstreams and methods

Work stream	Domain	Questions to be answered	Methods	Range of ratings
1	Benefits	How effective is the intervention?	Quantitative systematic reviews of effectiveness studies	Large, moderate, small, trivial, none, varies, unknown
	Harms	Are there important adverse events reported by the study from the intervention?	Quantitative systematic reviews of effectiveness studies	Large, moderate, small, trivial, none, varies, unknown
	Balance of effects	Does the balance between benefits and harms favour the intervention?	DECIDE approach ^a	Favours intervention, probably favours intervention, probably favours no intervention, favours no intervention, varies, unknown
	Certainty	What is the certainty of the effectiveness evidence?	GRADE ^b or GRADE-CERQual ^c assessment of the certainty of the body of evidence	Bias, imprecision, inconsistency, indirectness High, moderate, low, very low certainty
2	Values and preferences	Is there important variability in the values or preferences a family might have about the outcomes that would impact judgements about the balance of effects?	Qualitative systematic reviews of experimental, quasi-experimental and observational studies	Yes, probably yes, probably no, no, varies, unknown
	Acceptability	Is the intervention acceptable?	Qualitative systematic reviews of experimental, quasi-experimental and observational studies	Yes, probably yes, probably no, no, varies, unknown
	Resources	What resources are required and what are their costs?	Structured searches in resource, cost, feasibility and equity databases ^d	Negligible costs, low-to-moderate costs, large costs, varies, unknown
	Feasibility	What is the feasibility of the intervention? Can it be easily or conveniently implemented? Is the intervention acceptable and are the resources required achievable?	Structured searches in resource, cost, feasibility and equity databases ^d	Yes, probably yes, probably no, no, varies, unknown
	Equity	Can the intervention be provided in low-resource settings? Will the populations that need the intervention most receive it quickly and at low cost?	Structured searches in resource, cost, feasibility and equity databases ^d	Yes, probably yes, probably no, no, varies, unknown

^a DECIDE = Developing and Evaluating Communication strategies to support Informed Decisions and practice based on Evidence (28).

^b GRADE = Grading of Recommendations Assessment, Development and Evaluation (29).

^c GRADE-CERQual = Confidence in the Evidence from Reviews of Qualitative Research (30).

^d Searches = Structured searches in UNICEF supply catalogue (31), International Medical Products Price Guide (32) and the *WHO compendium of innovative health technologies for low-resource settings* (33).

For effects (benefits and harms), evidence was derived from systematic reviews of randomized controlled trials (RCTs) where possible. If reviews of RCTs were not available, then systematic reviews of non-randomized studies of interventions were used. An overview of systematic reviews was compiled to identify all eligible systematic reviews that had been conducted in the last three years (26). If systematic reviews were not available, they were commissioned from expert systematic review groups. All commissioned systematic reviews followed standard methods, including: a standard protocol published in advance; a clear PICO question; criteria for identification of studies, including search strategies for different bibliographic databases; methods for assessing risk of bias; and a data analysis plan. The protocols were reviewed and approved by members of the Steering Group. The language used to describe the evidence on effects was consistent with the Cochrane Effective Practice and Organisation of Care approach (EPOC) (34). The GDG carefully considered the benefits and harms, the balance of effects, and the certainty of the evidence of effectiveness for each PICO question.

For values and acceptability, a systematic review on what matters to families about the care of their preterm or LBW infant was commissioned (14). This systematic review also followed standard methods for qualitative reviews, including: a standard protocol published in advance; a clear research question; criteria for identification of studies, including search strategies for different bibliographic databases; methods for assessing quality; and a data analysis plan. The protocol was also reviewed and approved by members of the Steering Group.

For resources, feasibility and equity, structured searches were done using search terms from effectiveness reviews and guidance published in the last five years. Databases included: Excerpta Medica database (Embase), MEDLINE, UNICEF supply catalogue, International Medical Products Price Guide, and the *WHO compendium of innovative health technologies for low-resource settings* (31-33,35,36).

This evidence was then compiled into a GRADE EtD framework for each priority question (see section 2.8).

2.5 Grading of the quality and certainty of the evidence

The GRADE approach was used to appraise the quality and certainty of the quantitative evidence for each priority question. GRADE is a standard systematic approach for developing and presenting summaries of evidence for clinical practice recommendations (29). It uses standard tools, which are published online, including GRADE protocols and risk-of-bias tools for assessing randomized and non-randomized studies. A GRADE EtD framework is prepared for each quantitative outcome and the certainty of evidence is rated as “high”, “moderate”, “low” or “very low”. The standard criteria for baseline GRADE ratings are that RCTs provide high-certainty evidence while non-randomized trials and observational studies provide low-certainty evidence. This baseline certainty rating is then downgraded based on characteristics of the study design: risk of bias, inconsistency, imprecision, indirectness and publication bias. Magnitude of effect and dose response allow upgrading of certainty for observational studies. Further details of the standard GRADE approach can be found online (29). For this guideline, both the systematic review teams and the external guideline methodologist (members of the EST) independently performed grading of the quantitative evidence for each priority question and outcome. Consensus was reached through discussion among the methodologist and all members of the EST.

For the qualitative evidence, the reviews were appraised using the GRADE-CERQual (Confidence in the Evidence from Reviews of Qualitative Research) tool (30). This tool uses an approach that is similar conceptually to other GRADE tools and provides a transparent method for assessing and assigning the level of confidence that can be applied to qualitative evidence. The three domains are values, acceptability and feasibility, and each of them has four components: methodological limitations of the individual studies; adequacy of data; coherence; and relevance to the review question.

2.6 Review of evidence, decision-making and recommendations

The WHO Steering Group provided the EtD frameworks to the GDG members as soon as the documents had been drafted, and in advance of

the virtual GDG meetings. The GDG was asked to review and provide comments on the documents electronically before the GDG meetings where possible. At the virtual meetings, under the leadership of the GDG chairs, GDG members collectively reviewed the EtD frameworks, the draft recommendations and any comments received through preliminary feedback.

The meetings included: presentation of the evidence and EtD frameworks by the EST; consideration of each EtD domain; presentation of draft recommendations by the WHO Steering Group; deliberations on each recommendation; and discussion about justification, caveats or difficulties, implementation considerations and research gaps.

The purpose of the GDG meetings was to reach consensus on each recommendation, including its direction, strength and conditions, based on explicit consideration of all the domains within the EtD frameworks.

Recommendations were developed using WHO Guidelines Review Committee (GRC) criteria (Box 2.1) (25):

Box 2.1 Approach for developing recommendations and good practice statements

The recommendation is:

A **"strong recommendation"** if the intervention is applicable to all preterm or low-birth-weight infants

- Strong recommendations should be phrased as "is recommended", "is not recommended", "should receive", "should not receive".

A **"conditional recommendation"** if the intervention is recommended under certain conditions, which could be shared decision-making, or in certain populations or settings

- Conditional recommendations should be phrased as "may be considered".

A **"good practice statement"** if the intervention is obviously beneficial and should be done in most circumstances, even though there is no, little or only very-low-certainty evidence

The recommendations should be accompanied by a description of the certainty of the body of evidence: "high", "moderate", "low", or "very low".

Source: WHO, 2014 (25).

The final adoption of each recommendation was made by consensus, defined as the agreement by three quarters or more of the GDG. Consensus was reached for all recommendations in this guideline and there were no strong disagreements.

The GDG also identified important research gaps and implications. Where the certainty of available evidence was rated as low or very low, the GDG considered whether further research should be prioritized, based on whether the research would: contribute to improvements in care of the preterm or LBW infant; fill a knowledge gap that would inform new recommendations or change an existing recommendation; be likely to promote equity; and be feasible to implement. The research implications are summarized in Chapter 6 and full details can be found in Web Annex B.

2.7 Document preparation and peer review

Following the final GDG meeting, the WHO responsible technical officer prepared a draft of the full guideline document to accurately reflect the deliberations and decisions of the GDG. Other members of the WHO Steering Group provided comments on the draft document before it was sent electronically to the GDG members for review and further comment. Subsequently, the revised document was also sent to the ERG members for peer review. The Steering Group carefully evaluated the input of the GDG members and the ERG peer reviewers for inclusion in the guideline document and made revisions to the draft document as needed. Further modifications to the guideline were limited to corrections of factual errors and improvements in language to address any lack of clarity and to conform to WHO style.

2.8 Presentation of the recommendations and evidence

The recommendations are presented in the summary table in the executive summary of this guideline (Table 1). In Chapter 3, the recommendations and associated GDG remarks are presented at the start of the sections about each intervention, followed by background information and definitions, and a summary of the evidence for each recommendation. The evidence summaries present the evidence

on effectiveness (benefits and harms) of the interventions (sources and characteristics of the evidence, critical outcomes, other outcomes and subgroup analysis) followed by a summary of other evidence (values and acceptability, resources, feasibility and equity). Finally for each intervention, a summary of judgements is presented in a table, including justifications for the recommendation made (if any) and the EtD summary.

The GRADE data tables for each priority question are presented in the Web Supplement.¹ The GRADE tables contain the grading of: bias, inconsistency,

indirectness, imprecision, number of participants, relative and absolute effect, risk difference and 95% confidence intervals. Further detail on methods can be found in the *WHO handbook for guideline development* and other documents (25,29).

This guideline is also accompanied by three web annexes:²

- Web Annex A: Priority questions and outcomes
- Web Annex B: Detailed list of research implications
- Web Annex C: Changes from approved scope of guideline.

1 Available at: <https://apps.who.int/iris/bitstream/handle/10665/363699/9789240060050-eng.pdf>

2 Available at: <https://apps.who.int/iris/bitstream/handle/10665/363698/9789240060043-eng.pdf>

3. Evidence and recommendations

This guideline includes 25 recommendations and 1 good practice statement (summarized in Table 1 in the executive summary and presented in detail in this chapter) for care of the preterm (born before 37 weeks' gestation) or low-birth-weight (LBW; < 2.5 kg) infant. Of the recommendations, 11 are new and 14 are updated, and the good practice statement is new. There are 11 strong recommendations for all preterm or LBW infants and 14 recommendations that are conditional on particular contexts or conditions.

Sixteen recommendations are for preventive and promotive care, six are for care for complications, and three are for family involvement and support. A good practice statement was made for parental leave and entitlements because the Guideline Development Group (GDG) determined that these have obvious benefits, although there was little evidence available.

The GDG provided remarks related to all the recommendations and the good practice statement, where needed. Users of the guideline should refer to these remarks, which are presented prominently along with the recommendations in this chapter.

The recommendations have been divided into the following categories, as presented in this chapter:

**A. PREVENTIVE AND PROMOTIVE CARE
(16 recommendations)**

**B. CARE FOR COMPLICATIONS
(6 recommendations)**

**C. FAMILY INVOLVEMENT AND SUPPORT
(3 recommendations, and
1 good practice statement)**

A. Preventive and promotive care

A.1 KANGAROO MOTHER CARE

Recommendation and remarks

RECOMMENDATION A.1a (UPDATED)

Any KMC:

Kangaroo mother care (KMC) is recommended as routine care for all preterm or low-birth-weight infants. KMC can be initiated in the health-care facility or at home and should be given for 8–24 hours per day (as many hours as possible). *(Strong recommendation, high-certainty evidence)*

RECOMMENDATION A.1b (NEW)

Immediate KMC:

Kangaroo mother care (KMC) for preterm or low-birth-weight infants should be started as soon as possible after birth. (Strong recommendation, high-certainty evidence) *(Strong recommendation, high-certainty evidence)*

Remarks

- Any KMC
 - KMC can be given at home or at the health-care facility.
 - Infants who receive KMC should be secured firmly to the mother's chest with a binder that ensures a patent airway.
 - Whenever possible, the mother should provide KMC. If the mother is not available, fathers, partners and other family members can also provide KMC.
 - Infants who need intensive care should be managed in special units, where mothers, fathers, partners and other family members can be with their preterm or LBW infants 24 hours a day.
- Immediate KMC
 - At home, immediate KMC should be given to infants who have no danger signs (22).
 - At health-care facilities, immediate KMC can be initiated before the infant is clinically stable unless the infant is unable to breathe spontaneously after resuscitation, is in shock or needs mechanical ventilation. The infant's clinical condition (including heart rate, breathing, colour, temperature and oxygen saturation, where possible) must be monitored.

Background and definitions

Kangaroo mother care (KMC) is defined by WHO as early, continuous and prolonged skin-to-skin contact between the mother (or other caregiver) and the baby, and exclusive breastfeeding (20). In 2015, WHO recommended that KMC be given to hospitalized babies under 2.0 kg as soon as the

babies were clinically stable (20). However, there has been wide variation among care providers (i.e. parents/primary caregivers and health workers) in the timing and duration of KMC (37,38). New studies have also been published that assess the effects of KMC provided before clinical stabilization and also KMC initiated in community settings (39,40).

Summary of the evidence

Overview	A.1a Any KMC	A.1b Immediate KMC
PICO	<p>Population – Preterm or LBW infants</p> <p>Intervention 1 – KMC</p> <p>Comparator 1 – Conventional newborn care</p> <p>Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up</p>	<p>Population – Preterm or LBW infants</p> <p>Intervention 2 – KMC initiated early (within 24 hours of birth, also called immediate KMC)</p> <p>Comparator 2 – Initiating KMC later (more than 24 hours after birth)</p> <p>Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up</p>
Setting, timing, subgroups	<p>Setting – Health-care facility or home in any country or setting</p> <p>Timing of intervention – From birth</p> <p>Subgroups</p> <ul style="list-style-type: none"> • Gestational age at birth (< 34 weeks, ≥ 34 weeks) • Birth weight (< 2.0 kg, ≥ 2.0 kg) • Daily duration of KMC achieved (< 8 hours, 8–16 hours, > 16 hours) 	

Effectiveness: Comparison 1 – KMC versus conventional newborn care

Sources and characteristics of the evidence

For the first comparison of KMC versus conventional newborn care, the effectiveness evidence was derived from a systematic review of 27 RCTs conducted between 1994 and 2021 that enrolled 11 956 infants (41). Six studies were from high-income countries (Australia, the United Kingdom of Great Britain and Northern Ireland and the United States of America [USA]), four studies were from upper-middle-income countries (China, Colombia, Ecuador and Malaysia), 15 were from lower-middle-income countries (Bangladesh, India, Indonesia, Kenya and Nepal) and two studies were from a low-income country (Ethiopia). Twenty-five studies were conducted in health-care facilities and two were community-based. In all but one of the studies, the infants were stabilized before enrolment. KMC was started within 24 hours after birth in two studies, between 1 and 7 days after birth in 10 studies, and more than 7 days after birth in 12 studies, but 3 studies did not report the timing of initiation of KMC. The duration of KMC was less than 8 hours in nine studies, between 8 and 16 hours in nine studies and more than 16 hours in four studies, while five studies did not report the duration of KMC.

Critical outcomes

Sixteen trials reported all-cause mortality, 11 reported severe morbidity (9 reported severe infection, 11 hypothermia), 11 reported growth outcomes (weight gain) and 1 reported neurodevelopment (1 reported Griffith quotients, 1 reported Bayley Scales of Infant and Toddler Development, third edition [BSID-III]). No serious adverse events were reported. (Full details

are provided in GRADE Table A.1a, in the Web Supplement.³⁾

- **Mortality:** For KMC compared with conventional newborn care, high-certainty evidence from 12 trials of 10 505 participants suggests a decrease in all-cause mortality at discharge, at 40 weeks postmenstrual age (PMA; i.e. the baby's age when counted from the first day of the mother's last menstrual period before pregnancy – see Glossary) or at 28 days of age (relative risk [RR] 0.68, 95% confidence interval [CI] 0.53 to 0.86). High-certainty evidence from four trials of 8031 participants suggests a decrease in all-cause mortality at 6 months of age (RR 0.75, 95% CI 0.62 to 0.92).
- **Morbidity:** Moderate-certainty evidence from nine trials of 9847 participants suggests a decrease in severe infection or sepsis at 40 weeks PMA or at 28 days after birth (RR 0.85, 95% CI 0.79 to 0.92). Moderate-certainty evidence from 11 trials of 1169 participants suggests a decrease in hypothermia at discharge, at 40 weeks PMA or at 28 days after birth (RR 0.32, 95% CI 0.19 to 0.53).
- **Growth:** Low-certainty evidence from 11 trials of 1198 participants suggests an increase in weight gain (in grams per day) at 28 days after birth (mean difference [MD] 4.08, 95% CI 2.30 to 5.86).
- **Neurodevelopment:** Very-low-certainty evidence from one trial of 579 participants suggests little or no effect on Griffith quotients for psychomotor development (all subscales) at 12 months corrected age (i.e. the chronological age [age since birth or “postnatal age”] minus the number of weeks or months born preterm –

3 Available online: <https://apps.who.int/iris/bitstream/handle/10665/363699/9789240060050-eng.pdf>

see Glossary) (MD 1.05, 95% CI -0.75 to 2.85). Very-low-certainty evidence from one trial of 516 participants suggests little or no effect on cognitive neurodevelopment at 12 months of age using the BSID-III (MD 0.21, 95% CI -1.84 to 2.27) and other neurodevelopment measures (language, motor).

Other outcomes

There was an increase in exclusive breastfeeding at discharge, at 40 weeks PMA or at 28 days of age (RR 1.48, 95% CI 1.44 to 1.52; 9 trials, 9983 participants) and at 1–3 months follow-up (RR 1.39, 95% CI 0.99 to 1.97; 7 trials, 8139 participants). There was an increase in any breastfeeding at discharge, at 40 weeks PMA or at 28 days of age (RR 1.15, 95% CI 1.07 to 1.23; 12 studies, 10 146 participants) and at three months follow-up (RR 1.03; 95% CI 1.02 to 1.04; $I^2 = 70\%$; 7 studies, 8463 participants). There was also a decrease in the length of hospital stay (MD -0.39 days, 95% CI -0.79 to 0.0; 12 studies, 1214 participants).

Subgroup analyses

Subgroup differences for morbidity, growth and neurodevelopmental outcomes could not be assessed as there were insufficient studies. For all-cause mortality, no subgroup differences were seen for setting (health-care facility, community), gestational age (mean gestational age < 34 weeks, ≥ 34 weeks), birth weight (birth or enrolment weight < 2.0 kg, ≥ 2.0 kg) or daily duration of KMC achieved (< 8 hours/day, 8–16 hours/day and > 16 hours/day), although the analysis for daily duration of less than 8 hours was limited by small sample size and imprecision.

Effectiveness: Comparison 2 – KMC initiated early versus later

Sources and characteristics of the evidence

For the second comparison of KMC initiated early (< 24 hours after birth) versus KMC initiated late (≥ 24 hours after birth), the effectiveness evidence was derived from a systematic review of four RCTs totalling 3603 infants (41). One study was from a high-income country (Sweden), two were from low-income countries (Gambia and Madagascar) and one was a multicountry study conducted in Ghana, India, Malawi, Nigeria and the United Republic of Tanzania. All studies were conducted in health-care facilities. Two studies enrolled babies irrespective of clinical stability, while one study enrolled only stable infants and one study enrolled only unstable babies. KMC was started as soon after birth as possible in all studies. The mean age at initiation of KMC was

1.3 hours, 13.6 hours and 19 hours after birth in three studies, while one study did not report the age of initiation of KMC. The duration of KMC was less than 8 hours in one study, more than 16 hours in two studies and not reported in one study.

Critical outcomes

For the comparison of KMC initiated early compared with KMC initiated late, three trials reported all-cause mortality, three reported morbidity (2 reported severe infection, 3 hypothermia), one reported growth (weight gain) and none reported neurodevelopment outcomes. (Full details are provided in GRADE Table A.1b, in the Web Supplement.)

- **Mortality:** High-certainty evidence from three trials of 3533 participants suggests a decrease in all-cause mortality by 28 days of age (RR 0.78, 95% CI 0.66 to 0.92).
- **Morbidity:** Low-certainty evidence from two trials of 3415 participants suggests a decrease in the risk of sepsis by 28 days (RR 0.85, 95% CI 0.76 to 0.96). High-certainty evidence from three trials of 3513 participants suggests a decrease in the risk of hypothermia by discharge or 28 days (RR 0.74, 95% CI 0.61 to 0.90).
- **Growth:** Low-certainty evidence from one trial of 204 participants suggests little or no effect on weight gain by 28 days follow-up (measured in grams per day) (MD 2.20, 95% CI -5.26 to 0.86).

Other outcomes

There was an increase in exclusive breastfeeding (EBF) by hospital discharge (RR 1.12, 95% CI 1.07 to 1.16; 3 trials, 3464 participants). There was little or no effect on EBF by 28 days of age (RR 1.01, 95% CI 0.98 to 1.04; 3 trials, 2841 participants). There was a decrease in length of hospital stay (in days) (MD -0.30, 95% CI -0.31 to -0.29; 3 studies, 3498 participants).

Subgroup analyses

Differences for morbidity, growth and neurodevelopment could not be assessed as there were insufficient studies. For all-cause mortality, no subgroup differences were seen for setting (facility, community), gestational age (mean gestational age < 34 weeks, ≥ 34 weeks), birth or enrolment weight (< 2.0 kg, ≥ 2.0 kg) or daily duration of KMC (< 8 hours/day, 8–16 hours/day and > 16 hours/day).

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see

Table 1.1) reported that families want to be involved in delivering care to infants and want to take an active role in deciding what interventions are given to infants, including skin-to-skin contact and feeding (14). A systematic review of caregivers' perspectives on KMC reported that social support, access to care and cultural norms were important drivers of family perceptions, practices, attitudes and values about KMC (38). Important elements included: services free of charge for users; support from health workers; parents allowed unlimited visiting hours at the health-care facility; a private, quiet space in the hospital to provide KMC; and involvement of fathers and partners. Another synthesis of qualitative studies suggested that providing KMC can be restorative as well as energy-draining for mothers, fathers and partners (37).

Resources required and implementation considerations

Organization of care

KMC can be implemented at home and at all levels of newborn care (primary, secondary and tertiary) (42). Health services should ensure family involvement in the care of their preterm or LBW infant, irrespective of the infant's clinical condition. This should include a policy of "zero separation" between families and their preterm or LBW infant. This needs close collaboration between families and newborn and maternity care providers. Health-care facilities should ensure that families have access to beds, food, bathing and toilet facilities throughout the infant's hospital stay.

KMC is ideally initiated immediately after birth, or after initial resuscitation if that is needed. When it is not possible for the mother to provide KMC, other family members should provide it. To prepare for this situation, family members should be identified before delivery, counselled and allowed access to maternity and newborn care areas. If the infant needs to be transferred to a special or intensive care unit, the infant should be transported safely in KMC with the mother or another family member.

Choice of the best location for further management should be guided by the clinical condition of the infant. Stable larger infants could receive KMC in postnatal wards, while smaller ones could receive KMC in special care units (e.g. "step down" units, special care nurseries), and infants with complications could receive KMC in intensive care units. Many babies who need special or intensive care (e.g. level

2 or 3 care) are often separated from their mothers, although KMC is essential for these babies. Units that care for preterm babies and mothers with zero separation are needed (e.g. maternal-neonatal intensive care units [M-NICU] [43] or "couplet care" units [44]).

Health-care facilities should provide support so that mothers and families can continue KMC at home after discharge. All preterm and LBW infants must be followed up after discharge, ideally through home visits.

Preterm or LBW infants born at home should receive immediate KMC if they do not have danger signs, and should be transferred to a health-care facility if needed.

Infrastructure, equipment and supplies

A binder may help to keep the infant in skin-to-skin contact with the mother's or caregiver's chest. The infant should also have a warm hat, socks and a diaper/nappy. The mother or caregiver should wear whatever is comfortable, provided the clothes accommodate the baby.

Other arrangements can also make the baby and mother more comfortable, e.g. reclining beds and chairs. Other equipment and supplies needed are the same as for other newborn and maternal care, including a thermometer suitable for measuring body temperature down to 35°C.

If M-NICUs or couplet care units are used, they should have all the infrastructure, equipment and supplies that NICUs have for small or sick babies and that maternity wards have for mothers. For babies, this includes continuous positive airway pressure (CPAP) machines, pulse oximeters and radiant warmers or incubators if the infant is not in KMC. For mothers, this includes adult beds and an examination area where she can receive the health care she needs.

Workforce, training, supervision and monitoring

Health workers at all levels can provide KMC support to mothers and families. Training includes helping mothers keep infants in skin-to-skin contact, helping them with breastfeeding, and providing other neonatal care. Health workers should record the duration of KMC provided per day in a clinical register (or in home-based records in the community) and should monitor this on a regular basis.

Scale-up

KMC should be scaled up as an integrated intervention within programmes, not as a stand-alone programme. Scaling up means ensuring all preterm and LBW babies receive KMC across the whole country and across all countries. It needs multiple high-intensity (i.e. high-frequency and quality) interventions in the different domains described above (i.e. organization of care, health workforce, and infrastructure, equipment and supplies), but it also needs leadership and governance, financing, and health information systems.

- Leadership and governance can include: high-level leadership from national and subnational policy-makers, programme managers and facility directors; policies to enable zero separation; licensing standards for health-care facilities; pre-service education of health workers; and engagement with professional organizations.
- Health financing can include: dedicated line items in national budgets for KMC and expanded health insurance that includes KMC.

- Health information systems can include: monitoring of coverage and quality of KMC in routine health systems in health-care facilities and at the district and national levels.
- More detailed guidance on scaling up based on the results of implementation research (43,45-50) is being developed and will be published separately.

Feasibility and equity

Facility-based studies have shown that KMC can be provided to small babies, for more than 8 hours per day, and that it can be initiated immediately after birth irrespective of clinical stability (39,43,45,46). These studies were conducted in poor, remote and urban communities in “real world pragmatic” settings (40,51). However, community-initiated KMC and KMC for unstable babies have not been implemented outside research settings and global coverage remains low (52,53).

Summary of judgements

	Comparison 1: KMC vs conventional newborn care (A.1a)	Comparison 2: Immediate KMC vs later KMC (A.1b)
Justification	<ul style="list-style-type: none"> ▪ Evidence of large benefits: decreased mortality (<i>high-certainty evidence</i>), decreased infection (<i>moderate-certainty evidence</i>), decreased hypothermia (<i>moderate-certainty evidence</i>), increased weight gain (<i>low-certainty evidence</i>) and increased breastfeeding (<i>very-low-certainty evidence</i>) ▪ No evidence of harms 	<ul style="list-style-type: none"> ▪ Evidence of large benefits: decreased mortality (<i>high-certainty evidence</i>), decreased hypothermia (<i>high-certainty evidence</i>), decreased infections and increased weight gain (<i>low-certainty evidence</i>) ▪ No evidence of harms
Evidence-to-Decision summary		
Benefits	Large	Large
Harms	Trivial or none	Trivial or none
Certainty	Moderate	Moderate
Balance	Favours KMC	Favours immediate KMC
Values	No uncertainty or variability about outcomes	No uncertainty or variability about outcomes
Acceptability	Varies	Varies
Resources	Low to moderate	Low to moderate
Feasibility	Probably feasible	Probably feasible
Equity	Probably equitable	Probably equitable

A.2 MOTHER'S OWN MILK

Recommendation and remarks

RECOMMENDATION A.2 (UPDATED)

Mother's own milk is recommended for feeding of preterm or low-birth-weight (LBW) infants, including very preterm (< 32 weeks' gestation) or very LBW (< 1.5 kg) infants. (Strong recommendation, low-certainty evidence)

Remarks

- The GDG made a strong recommendation despite low-certainty evidence because of the consistent harm from infant formula on two critical outcomes (necrotizing enterocolitis and infection) and lack of evidence of benefit from infant formula.
- The GDG also considered that providing mother's own milk is the standard of care across all countries and the core of many national policies and programmes.
- Mothers should also be encouraged and supported before and after birth to provide their own breast-milk (including colostrum) for their infants.

Background and definitions

Mother's own milk confers important immune and nutritional advantages for preterm and LBW infants (54-56). Artificial formulas can be manipulated to contain higher amounts of important nutrients (such as protein) than mother's own milk (55,57). However,

formula milks do not contain the antibodies and immune modulators and primers present in human milk that protect the immature gastro intestinal tract of preterm and LBW infants (19,58,59). In 2011, WHO recommended that mother's own milk should be given to all preterm and LBW infants (19).

Summary of the evidence

OVERVIEW	A.2 Mother's own milk
PICO	Population – Preterm or LBW infants Intervention – Infant formula (term or preterm) Comparator – Mother's own milk Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Setting, timing, subgroups	Timing of the intervention – From birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none">▪ Gestational age at birth (< 32 weeks, ≥ 32 weeks)▪ Birth weight (< 1.5 kg, ≥ 1.5 kg)▪ Type of milk in the control group (mother's own milk as the sole diet, mother's own milk not the sole diet)

Effectiveness: Comparison – Any formula milk versus mother's own milk

Sources and characteristics of the evidence

The effectiveness evidence was derived from an updated systematic review of 42 studies reporting on 89 638 preterm or LBW infants from 20 countries (Australia, Belgium, Chile, China, Germany, Ghana, Greece, India, Israel, Italy, Japan, Nepal, the Netherlands, New Zealand, Poland, Romania, Spain, Sweden, the United Kingdom and the USA) (60).

Studies were included if they compared infants who received formula as the sole or predominant

(> 50%) diet (intervention group) with infants who received mother's own milk as the sole or predominant (> 50%) diet (comparison group) in the first 28 days after birth. Of the 89 638 participants, approximately 87% of infants were very preterm (< 32 weeks' gestation) or VLBW (< 1.5 kg). Studies typically excluded infants with congenital anomalies or gastrointestinal or neurological problems.

All the included studies were observational; there were no RCTs. Thirty-six studies were from hospitals and six were from the "whole population" (all infants born in the study area regardless of whether they

were admitted to hospital). The largest study (72 997 participants) was an observational study of all infants under 32 weeks' gestation admitted to 777 neonatal intensive care units (NICUs) in the USA. The studies used a combination of milks in the intervention and comparison groups.

In the intervention group, all 42 studies used formula milk as the sole or predominant (> 50%) diet. Among these studies, 24 studies gave formula milk as the sole diet, 13 mixed formula with mother's own milk, 5 mixed formula with donor milk and mother's own milk, and 6 did not state whether they mixed formula milk with other milks. Twenty-one studies used preterm formula, 5 used term formula, 2 used a combination of preterm and term formula, and 14 did not state which type of formula was used.

In the comparison group, all 42 studies used mother's own milk as the sole or predominant (> 50%) diet. Among these studies, 9 studies gave mother's own milk as the sole diet, 17 mixed mother's own milk with donor human milk, and the remainder did not state if they mixed mother's own milk with other milks. Twenty studies used fortifier, 6 did not use fortifier and 16 did not state whether fortifier was provided.

Babies all received their feeds from birth until discharge or 28 days of age. Twenty-five used parenteral nutrition, 10 did not use parenteral nutrition and the remainder did not state if parenteral nutrition was used.

Critical outcomes

For the comparison of any formula milk with mother's own milk, 5 studies reported all-cause mortality, 15 studies reported morbidity (15 reported necrotizing enterocolitis, 15 severe infection), 7 studies reported growth (3 reported weight-for-age z score [WAZ], 3 WAZ change, 9 length, 3 length-for-age z score [LAZ], 9 head circumference) and 8 studies reported neurodevelopment (8 reported cognitive outcomes, 3 language outcomes). (Full details are provided in GRADE Table A.2, in the Web Supplement.)

- **Mortality:** Low-certainty evidence from five observational studies of 9673 participants suggests little or no effect on all-cause mortality at latest follow-up (mean 116 days) (OR 1.26, 95% CI 0.91 to 1.76).
- **Morbidity:** Low-certainty evidence from 15 observational studies totalling 3013 participants suggests an increase in necrotizing enterocolitis

at latest follow-up (mean 44 days) (OR 2.99, 95% CI 1.75 to 5.11). Very-low-certainty evidence from 15 observational studies totalling 2562 participants suggests an increase in severe infection at latest follow-up (mean 31 days) (OR 1.52, 95% CI 0.98 to 2.37).

- **Growth:** Very-low-certainty evidence from three observational studies totalling 271 participants suggests little or no effect on weight (weight-for-age z score [WAZ]) between 39 and 416 weeks (MD 0.02, 95% CI -0.28 to 0.31). Very-low-certainty evidence from four observational studies totalling 74 130 participants suggests little or no effect on weight (WAZ change) from birth to discharge (mean 52 days) (MD 0.14, 95% CI -0.76 to 1.05). Very-low-certainty evidence from nine observational studies totalling 1048 participants suggests little or no effect on length (in centimetres) at latest follow-up (mean 58 days) (MD 0.33, 95% CI -0.4 to 1.05). Very-low-certainty evidence from three observational studies totalling 271 participants suggests little or no effect on length (LAZ) at 39 to 416 weeks (MD 0.06, 95% CI -0.81 to 0.92). Very-low-certainty evidence from nine observational studies totalling 1550 participants suggests little or no effect on head circumference (in centimetres) at latest follow-up (mean 45 days) (MD 0.26, 95% CI -0.35 to 0.87).
- **Neurodevelopment:** Very-low-certainty evidence from eight observational studies totalling 1560 participants suggests little or no effect on cognitive development at follow-up (range: 91 to 416 weeks) (standardized mean difference [SMD] 1.3 standard deviation [SD] lower, 95% CI -3.53 to 0.93). Very-low-certainty evidence from three observational studies totalling 587 participants suggests little or no effect on language development at follow-up (range: 39-104 weeks) (SMD 0.02 SD lower, 95% CI -0.39 to 0.43).

Subgroup analyses

There was no evidence of a subgroup difference by gestational age, birth weight, or type of milk in the control group for any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding

what interventions are given to infants, including what and how they are fed (14). Two qualitative reviews reported that parents understood the importance of expressing breast-milk for the care of their baby but also found it challenging to express breast-milk unless supported by hospital staff and adequately informed about resources (61,62). Reviews also report that families value having formula available if their circumstances demand it – for example, work commitments, maternity leave, night-time feeding, father/partner support (14).

Resources required and implementation considerations

Organization of care

Mother's own milk should be provided through direct breastfeeding wherever possible. If direct breastfeeding is not possible, then breast-milk can be expressed and provided using cups and gastric tubes.

Infrastructure, equipment and supplies

Breastfeeding requires no specific infrastructure, equipment or supplies. If expressed breast-milk is needed, milk can be expressed by hand or using

a manual breast pump. Supplies are also needed for cup and gastric tube feeding. National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can provide breastfeeding support to mothers and families. Standardized packages are needed for training, supervision and monitoring.

Feasibility and equity

Difficulties related to breastfeeding and expressing breast-milk in hospitals can include lack of privacy, inadequate training from busy health workers, and feelings of stress and inadequacy from mothers and families (63). There are also studies that report difficulties in providing mother's own milk when the mother and baby return home from hospital, including difficulties balancing work commitments, maternity leave, night-time feeding and father and partner support (14). There are many studies that report problems in sourcing clean water to reconstitute infant formula and wash receptacles in resource-limited settings (64,65).

Summary of judgements

Comparison: Any formula milk vs mother's own milk (A.2)

Justification

- No evidence of benefits of infant formula
- Evidence of moderate harms from using infant formula instead of mother's own milk: increased necrotizing enterocolitis (*low-certainty evidence*) and increased infections (*very-low-certainty evidence*)
- Evidence of little or no effect of using infant formula on mortality (*low-certainty evidence*), weight gain (*very-low-certainty evidence*) and neurodevelopment (*very-low-certainty evidence*)
- No evidence on other critical outcomes

Evidence-to-Decision summary

Benefits	Benefits of infant formula are trivial or none
Harms	Harms of infant formula are moderate
Certainty	Low
Balance	Does not favour infant formula, favours mother's own milk
Values	Probably no important uncertainty or variability about outcomes
Acceptability	Acceptability of infant formula varies, acceptability of mother's own milk does not vary
Resources	Low to moderate (costs of infant formula), negligible (costs of mother's own milk)
Feasibility	Feasibility of infant formula varies, feasibility of mother's own milk does not vary, where it is available
Equity	Equity of infant formula varies, equity of mother's own milk does not vary

A.3 DONOR HUMAN MILK

Recommendation and remarks

RECOMMENDATION A.3 (UPDATED)

When mother's own milk is not available, donor human milk may be considered for feeding of preterm or low-birth-weight (LBW) infants, including very preterm (< 32 weeks' gestation) or very LBW (< 1.5 kg) infants. (Conditional recommendation, moderate-certainty evidence)

Remarks

- The recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- The potential harm of necrotizing enterocolitis from infant formula was considered by the GDG to be more clinically important than the benefit of increased growth from infant formula.
- Donor human milk was pasteurized in all but one trial, so the GDG was not able to make a recommendation on the use of unpasteurized milk.
- Safe and affordable milk-banking facilities are needed for the provision of donor human milk.
- Mothers should also be encouraged and supported before and after birth to provide their own breast-milk (including colostrum) for their infants.

Background and definitions

When mother's own milk is not available, preterm or LBW infants must be given other milks. Donor human milk is provided through human milk banks (i.e. places where human milk is collected, treated and/or distributed) (56,66). Donor milk has differences in immune composition to mother's own milk.

Human milk banks also usually pasteurize milk to remove infective organisms, which further alters milk components (56,66). WHO LBW feeding guidelines in 2011 recommended feeding donor human milk rather than infant formula to preterm or LBW babies who cannot be fed mother's own milk (19). However, new studies have been published since that time.

Summary of the evidence

OVERVIEW	A.3 Donor human milk
PICO	Population – Preterm or LBW infants Intervention – Infant formula Comparator – Donor human milk Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none"> • Gestational age at birth (< 32 weeks, ≥ 32 weeks) • Birth weight (< 1.5 kg, ≥ 1.5 kg) • Amount of donor milk in the control arm (donor milk provided as the sole diet, mixed with infant formula)

Effectiveness: Comparison – Infant formula versus donor human milk

Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review published in 2019 of 12 RCTs enrolling 1879 preterm or LBW infants from neonatal units in eight countries (Austria, Canada, Finland, Hungary, Italy, the Netherlands, the United

Kingdom and the USA) (67). An updated search conducted on 1 October 2021 located no new trials. Participants were clinically stable preterm or LBW infants. Most were below 32 weeks' gestational age or below 1.8 kg at birth. Many trials excluded infants who were small for gestational age at birth and infants with congenital anomalies or gastrointestinal or neurological problems. The

trials varied according to whether formula or donor milk was provided as the sole diet (5 trials) or as a supplement to mother's own milk (7 trials). A mix of term and preterm formula was used. The donor milk was a mix of preterm and term donor milk and a mix of fortified and unfortified milk. In all trials except one, the donor human milk was pasteurized. In general, feeds were allocated for several weeks, or until participating infants reached a specified body weight (generally > 2 kg). One trial used the allocated feed for less than 10 days after birth. Infants then received preterm formula if their mother's own milk was insufficient.

Critical outcomes

For infant formula compared with donor human milk, seven trials reported all-cause mortality, nine reported morbidity (9 reported necrotizing enterocolitis, 5 invasive infection), nine reported growth (9 reported weight gain, 8 length, 8 head growth) and two reported neurodevelopment (neurodevelopmental disability). (Full details are provided in GRADE Table A.3, in the Web Supplement.)

- **Mortality:** Moderate-certainty evidence from seven trials totalling 1527 participants suggests little or no effect on all-cause mortality by hospital discharge (RR 1.1, 95% CI 0.8 to 1.5).
- **Morbidity:** Moderate-certainty evidence from nine trials totalling 1675 participants suggests an increase in risk of necrotizing enterocolitis by hospital discharge (RR 1.87, 95% CI 1.23 to 2.85). Moderate-certainty evidence from five trials totalling 1025 participants suggests little or no effect on risk of invasive infection by hospital discharge (RR 0.94, 95% CI 0.79 to 1.12).
- **Growth:** Moderate-certainty evidence from nine trials totalling 1028 participants suggests an increase in weight gain (in grams per kilogram per day) by hospital discharge (MD 2.51, 95% CI 1.93 to 3.08). Moderate-certainty evidence from eight trials totalling 820 participants suggests an increase in linear growth (crown-heel length, measured in millimetres per week) by hospital discharge (MD 1.21, 95% CI 0.77 to 1.65). Moderate-certainty evidence from eight trials totalling 894 participants suggests an increase in head growth (in millimetres per week) by hospital discharge (MD 0.85, 95% CI 0.47 to 1.23).

- **Neurodevelopment:** Moderate-certainty evidence from two trials totalling 400 participants suggests little or no effect on neurodevelopmental disability by 18 months of age (RR 1.21, 95% CI 0.62 to 2.35).

Two studies in the review also reported on long-term growth outcomes. Neither individual study nor meta-analyses of data from both studies showed differences in weight, length or head circumference at follow-up at 9 months, 18 months or 7.5–8 years of age. For the latest follow-up at 7.5–8 years of age, there was no difference in growth parameters between infants fed formula milk or donor human milk (weight [kg], MD -0.56, 95% CI -1.42 to 0.29; length [cm], 0.05, 95% CI -1.12 to 1.23; and head circumference [cm], MD -0.19, 95% CI -0.54 to 0.16; 2 studies, 420 participants).

Other outcomes

There was higher risk of feeding intolerance in the formula-fed group compared with the donor milk group (moderate-certainty evidence) (RR 4.92, 95% CI 1.17 to 20.70; 2 trials, 148 participants).

Subgroup analyses

For the analyses by gestational age and birth weight and amount of donor milk in the control arm, differences for all critical outcomes could not be assessed as there were insufficient studies.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding what interventions are given to infants, including what and how they are fed (14). A number of studies report the facilitators and barriers to donating and receiving donor human milk (68–71). These include preferences for receiving human rather than artificial milk, concerns about the effect of pasteurization and transportation, and concerns that the mother's own breast-milk supply will reduce (68–71). A large cross-sectional survey among health workers in urban Zimbabwe reported that the concept of donor human milk banking was acceptable, and that the participants would accept donor human milk for their children, and many would encourage their clients to donate human milk (68).

Resources required and implementation considerations

Organization of care

The provision of donor human milk requires access to a human milk bank where milk can be tested, pasteurized and transported safely.

Infrastructure, equipment and supplies

Infrastructure, equipment and supplies are needed for donor assessment (screening, informed consent, serological testing), milk expression, handling, storage, transport, pre-pasteurization testing, pasteurization, and post-pasteurization testing. Supplies are also needed for safe cup and gastric tube feeding.

Workforce, training, supervision and monitoring

Specialized staff are needed for the operation of donor human milk banks. Standardized packages are needed for training, supervision and monitoring. More detailed guidance on the operation of donor human

milk banks is being developed and will be published separately. Health workers at all levels can provide feeding support.

Feasibility and equity

A census of milk banks from a systematic literature review reported 572 milk banks globally in 60 countries, with the majority in high-income countries (68). It is well known that safe and affordable milk-banking facilities are needed for the provision of donor human milk. However, the base resources for donor milk feeding (i.e. donor recruitment, donor assessment [screening, informed consent, serological testing], milk expression, handling, storage, transport, pre-pasteurization testing, pasteurization, post-pasteurization testing) are much less available in low- and middle-income countries (LMICs), especially in smaller towns and villages (66,72). The use of donor milk varies widely within and between countries and is influenced by cultural practices, access, costs, awareness, supportive policies and resources (66,72).

Summary of judgements

Comparison: Infant formula vs donor human milk (A.3)

Justification

- In trials where most participants are very preterm (< 32 weeks' gestation) or VLBW (< 1.5 kg):
- Evidence of small benefits from using infant formula instead of donor human milk: increased in-hospital weight gain, length and head circumference (*moderate-certainty evidence*)
 - Evidence of moderate harms from using infant formula instead of donor human milk: increased necrotizing enterocolitis and feed intolerance (*moderate-certainty evidence*)
 - Evidence of little or no effect of using infant formula on mortality and neurodevelopment (*moderate-certainty evidence*)
 - No evidence on other critical outcomes

Evidence-to-Decision summary

Benefits	Benefits of infant formula are small
Harms	Harms of infant formula are moderate
Certainty	Moderate
Balance	Probably does not favour infant formula, probably favours donor human milk
Values	Probably no important uncertainty or variability about outcomes
Acceptability	Acceptability of infant formula and donor human milk varies
Resources	Resources for infant formula and donor human milk vary
Feasibility	Feasibility of infant formula and donor human milk varies
Equity	Equity of infant formula and donor human milk varies

A.4 MULTICOMPONENT FORTIFICATION OF HUMAN MILK

Recommendation and remarks

RECOMMENDATION A.4 (UPDATED)

Multicomponent fortification of human milk is not routinely recommended for all preterm or low-birth-weight (LBW) infants but may be considered for very preterm (< 32 weeks' gestation) or very LBW (< 1.5 kg) infants who are fed mother's own milk or donor human milk. *(Conditional recommendation, low- to moderate-certainty evidence)*

Remarks

- The potential harm of mortality and necrotizing enterocolitis from fortification was considered by the GDG to be very uncertain due to the low quality of the included trials. The GDG also considered that the benefits of multicomponent fortifier were clinically important for the weight, length and head circumference of very preterm (< 32 weeks) or very-low-birth-weight (VLBW) (< 1.5 kg) infants. Thus, the GDG decided not to routinely recommend multicomponent fortifier for all preterm or LBW infants and suggested that fortification may be considered for very preterm or VLBW infants. This recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- The GDG noted that there were limited data on the type of fortifier used in the studies. Based on most trials included in the evidence review, the GDG suggests that commercially available multicomponent fortifiers specifically formulated for preterm infants may be considered.
- The GDG also noted that there were limited data on the timing of initiation and duration of fortification in the studies. The GDG suggests that the initiation and duration of multicomponent fortification should be based on clinical judgement.
- Mothers should also be encouraged and supported before and after birth to provide their own breast-milk (including colostrum) for their infants.

Background and definitions

Commercially available multicomponent fortifiers for infant human milk feeding can be human or animal (often cows' milk) protein based, and contain carbohydrate, fat, protein, multivitamins, iron, zinc, calcium and phosphorous in varying amounts (56,73). They are provided as liquid or powder and mixed with mother's own or donor human milk (74,75). Some health workers advise families to add multicomponent fortifier to human milk feeds for preterm and LBW infants with the intent

to increase nutrient accretion (76,77). However, there are concerns that multicomponent fortifiers are associated with adverse events such as feed intolerance and necrotizing enterocolitis (56). WHO guidelines in 2011 recommended against the use of multicomponent fortifiers for all preterm and LBW babies but to use them for very-low-birth-weight (VLBW) babies (< 1.5 kg) or very preterm babies (< 32 weeks' gestation) who fail to gain weight (19). There have been new trials since that time.

Summary of the evidence

OVERVIEW	A.4 Multicomponent fortification of human milk
PICO	Population – Preterm or LBW infants Intervention – Human milk with multicomponent fortifier (human derived or non-human derived) Comparator – Human milk without multicomponent fortifier Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none"> • Gestational age at birth (< 32 weeks, ≥ 32 weeks) • Birth weight (< 1.5 kg, ≥ 1.5 kg) • Type of fortifier (human milk protein based, non-human milk protein based)

Effectiveness: Comparison – Multicomponent fortification versus unfortified breast-milk

Sources and characteristics of the evidence

The effectiveness evidence was derived from a 2019 Cochrane review of 18 small trials totalling 1456 preterm infants (78). An updated search conducted on 1 October 2021 located no new trials. All trials were conducted in specialist paediatric hospitals, typically in NICUs. The trials were conducted in 11 countries (Brazil, Canada, Denmark, Egypt, India, Italy, Oman, South Africa, Sweden, the United Kingdom and the USA). Babies were mostly very preterm (< 32 weeks' gestation) or VLBW (< 1.5 kg).

Trials used a range of different “base” milks to feed the infants which were identical in the intervention and the control arms. Six trials used only mother's own milk, one trial used only donor human milk, seven trials used a mixture of mother's own milk and donor milk, and four trials used a mixture of mother's own milk, donor milk and preterm formula. Participants received the intervention once they were tolerating a specified quantity of milk feeding, typically at least 100 ml/kg per day, or when receiving “full” enteral feeds, typically 150 ml/kg per day.

In the intervention arm in all trials, multicomponent fortifier was mixed into the base milk and was provided according to the manufacturer's

specifications. Fourteen trials used a commercially available, bovine-milk-based, powdered preparation and four trials used preterm formula powder as the multicomponent fortifier. No trials used human-milk-derived fortifier. The fortifier was provided until a prespecified body weight was attained (most commonly, 1.8–2.0 kg), until a prespecified PMA (most commonly 34–36 weeks) or until discharge home from hospital.

In the control arm, eight trials gave infants multiple supplements (i.e. multivitamins, iron, zinc, calcium and phosphorus) in similar quantities to the nutrients in multicomponent fortifier, five trials gave infants only vitamin D, and five trials gave no supplements at all. No trials gave infants additional carbohydrate or protein in the control arm.

Critical outcomes

For multicomponent fortification compared with unfortified breast-milk, two trials reported all-cause mortality, 13 reported morbidity (13 reported necrotizing enterocolitis), 14 reported growth (14 reported weight gain, 10 length gain, 11 head growth) and 1 reported neurodevelopment (Mental Development Index [MDI, BSID-II] and Psychomotor Development Index [PDI, BSID-II]). (Full details are provided in GRADE Table A.4, in the Web Supplement.)

- **Mortality:** Very-low-certainty evidence from two trials totalling 375 participants suggests an increase in all-cause mortality by discharge (RR 2.33, 95% CI 0.16 to 34.76).
- **Morbidity:** Low-certainty evidence from 13 trials totalling 1110 participants suggests an increase in necrotizing enterocolitis by hospital discharge (RR 1.37, 95% CI 0.72 to 2.63).
- **Growth:** Low-certainty evidence from 14 trials totalling 951 participants suggests an increase in weight gain (in grams per kilogram per day) by hospital discharge (MD 1.76, 95% CI 1.30 to 2.22). Low-certainty evidence from 10 trials totalling 741 participants suggests an increase in length gain (in centimetres per week) by hospital discharge (MD 0.11, 95% CI 0.08 to 0.15). Moderate-certainty evidence from 11 trials totalling 821 participants suggests an increase in head growth (in centimetres per week) by hospital discharge (MD 0.06, 95% CI 0.03 to 0.08).
- **Neurodevelopment:** Moderate-certainty evidence from one trial with 245 participants suggests little or no effect on MDI (BSID-II) by 18 months of age (MD 2.20, 95% CI -3.35 to 7.75). Moderate-certainty evidence from one trial totalling 245 participants suggests little or no effect on PDI (BSID-II) by 18 months of age (MD 2.40, 95% CI -1.90 to 6.70).

Other outcomes

There was little or no effect on length of hospital stay in weeks (MD -0.07, 95% CI -0.35 to 0.21; 6 trials, 526 infants), or feed intolerance (RR 1.05, 95% CI 0.65 to 1.67; 7 trials, 453 infants).

Subgroup analyses

The effect of gestational age and birth weight and type of fortifier could not be assessed as there were insufficient studies.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding what interventions are given to infants, including what and how they are fed (14). No other specific evidence was located about whether families value fortified feeds rather than unfortified feeds for their preterm or LBW baby, or find fortified feeds more or less acceptable than unfortified feeds.

Resources required and implementation considerations

Organization of care

Health-care facilities can provide multicomponent fortifier for preterm or LBW infants.

Infrastructure, equipment and supplies

The main commodity required is the fortifier, which should be a standard, nationally approved, multicomponent fortifier specially formulated for preterm or LBW infants. Commonly used fortifiers have similar amounts of carbohydrate, protein and micronutrients. Facilities for expressing breast-milk are also needed, as are facilities for the safe mixing of fortifier into expressed breast-milk. Supplies are also needed for cup or gastric tube feeding.

Workforce, training, supervision and monitoring

Health workers at all levels can provide support to mothers and families. Standardized packages are needed for training, supervision and monitoring.

Feasibility and equity

There was no specific evidence on the feasibility and equity of providing multicomponent fortifier for preterm or LBW infants.

Summary of judgements

Comparison: Multicomponent fortification vs unfortified breast-milk (A.4)

Justification	<p>In trials where most participants are very preterm (< 32 weeks' gestation) or VLBW (< 1.5 kg):</p> <ul style="list-style-type: none"> ▪ Evidence of small benefits: increase in in-hospital weight, length and head circumference (<i>moderate- to low-certainty evidence</i>) ▪ Evidence on harms uncertain: mortality (<i>very-low-certainty evidence</i>), necrotizing enterocolitis (<i>low-certainty evidence</i>) ▪ Evidence of little or no effect on neurodevelopment (<i>moderate-certainty evidence</i>) ▪ No evidence on other critical outcomes
----------------------	--

Evidence-to-Decision summary

Benefits	Small
Harms	Unknown
Certainty	Low
Balance	Varies
Values	Uncertainty or variability about outcomes
Acceptability	Unknown
Resources	Low to moderate
Feasibility	Varies
Equity	Not equitable

A.5 PRETERM FORMULA

Recommendation and remarks

RECOMMENDATION A.5 (UPDATED)

When mother's own milk and donor human milk are not available, nutrient-enriched preterm formula may be considered for very preterm (< 32 weeks' gestation) or very low-birth-weight infants. *(Conditional recommendation, low-certainty evidence)*

Remarks

- The recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- The GDG was not able to recommend a particular type of preterm formula. Based on most trials included in the evidence review, the GDG suggests that commercially available nutrient-enriched formulas specifically formulated for preterm infants may be considered.
- There was insufficient evidence to make a recommendation for infants who were born at 32–36 weeks' gestation or with birth weight of 1.5–2.4 kg. For these infants, the GDG considered that standard term formula or nutrient-enriched preterm formula may be considered, depending on clinical judgement.
- The GDG also noted that there was limited information on the timing of initiation and duration of preterm formula in the studies. The GDG suggests initiation and duration should be based on clinical judgement.
- Mothers should also be encouraged and supported before and after birth to provide their own breast-milk (including colostrum) for their infants.

Background and definitions

If human milk is not available, then preterm and LBW infants need to be given infant formula in the first six months after birth (56). Some studies suggest that feeding preterm infants with nutrient-enriched formula (or preterm formula) rather than formula developed for term infants (also called term formula, or non-nutrient-enriched formula) might increase

nutrient accretion, growth and neurodevelopmental outcomes (76,79,80). Preterm formula often has energy content over 72 kcal/100 ml and protein content over 1.7 g/100 ml (56,81). Term formula milks have varying energy and protein content, usually below these values (56,81). In 2011, WHO did not recommend preterm formula for feeding preterm and LBW infants (19).

Summary of the evidence

OVERVIEW	A.5 Preterm formula
PICO	Population – Preterm or LBW infants Intervention – Nutrient-enriched formula (or preterm formula) Comparator – Non-nutrient-enriched formula (or term formula) Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none"> • Gestational age at birth (< 32 weeks, ≥ 32 weeks) • Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – Preterm formula versus term formula

Sources and characteristics of the evidence

The effectiveness evidence was derived from a 2019 Cochrane systematic review of seven trials including 590 infants (81). An updated search conducted on 1 September 2021 located no new trials. The trials

were undertaken during the 1970s and 1980s in neonatal units in South Africa, Thailand, Türkiye, the United Kingdom and the USA. All infants were clinically stable preterm infants. Most were very low birth weight (< 1.5 kg). Few participants were extremely preterm (< 28 weeks), extremely low birth weight (< 1.0 kg) or growth restricted. The

trials excluded infants with congenital anomalies, or respiratory, gastrointestinal or neurological problems.

Preterm formula was defined in the systematic review as a formula with both energy content over 72 kcal/100 ml and protein content over 1.7 g/100 ml and term formula was defined as a formula with both energy content below 72 kcal/100 ml and protein content below 1.7 g/100 ml. In six trials, the formula was the sole diet while in one trial the formula was used in addition to human milk. The milk feeds were started when infants were clinically stable and able to tolerate enteral feeds in all trials. Trial participants continued to receive the intervention or control formula for two weeks or until they reached 2.0 kg. The target volume of milk intake for both groups was 150–180 ml/kg per day.

Critical outcomes

For preterm formula compared with term formula, two trials reported all-cause mortality, three reported morbidity (3 reported necrotizing enterocolitis), five reported growth (6 reported weight gain, 5 length gain, 5 head circumference) and two reported neurodevelopment (both reported MDI and PDI). (Full details are provided in GRADE Table A.5, in the Web Supplement.)

- **Mortality:** Low-certainty evidence from two trials totalling 424 participants suggests little or no effect on all-cause mortality by hospital discharge (RR 1.12, 95% CI 0.65 to 1.93).
- **Morbidity:** Low-certainty evidence from three trials totalling 489 participants suggests a decreased risk of necrotizing enterocolitis by hospital discharge (RR 0.72, 95% CI 0.41 to 1.25).
- **Growth:** Low-certainty evidence from six trials totalling 440 participants suggests an increase in weight gain (in grams per kilogram per day) by hospital discharge (MD 2.43, 95% CI 1.60 to 3.26). Low-certainty evidence from five trials totalling 386 participants suggests little or no effect on length gain (in millimetres per week) by hospital discharge (MD 0.22, 95% CI -0.70 to 1.13). Low-certainty evidence from five trials totalling 399 participants suggests an increase in head circumference gain (in millimetres per week) by hospital discharge (MD 1.04, 95% CI 0.18 to 1.89).
- **Neurodevelopment:** Moderate-certainty evidence from two trials totalling 310 participants suggests

an increase in MDI (BSID-II) at 18 months (MD 2.81, 95% CI -1.44 to 7.06). Low-certainty evidence from two trials totalling 310 participants suggests an increase in PDI (BSID-II) at 18 months (MD 6.56, 95% CI 2.87 to 10.26).

Subgroup analyses

For the analysis by gestational age and birth weight, differences for all critical outcomes could not be assessed as there were insufficient studies.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding what interventions are given to infants, including what and how they are fed (14). No other specific evidence was located about whether families value preterm formula rather than term formula for their preterm or LBW baby, or find preterm formula more or less acceptable than term formula.

Resources required and implementation considerations

Organization of care

Health workers and staff at other care facilities can provide preterm (nutrient-enriched) formula for preterm or LBW infants.

Infrastructure, equipment and supplies

The main commodity required is the preterm formula, which should be a standard, nationally approved formula, specially formulated for preterm or LBW infants. Facilities are needed for safe reconstitution of preterm formula. Supplies are also needed for cup or gastric tube feeding.

Workforce, training, supervision and monitoring

Health workers at all levels can provide support to mothers and families. Standardized packages are needed for training, supervision and monitoring.

Feasibility and equity

There was no specific evidence on the feasibility and equity of providing preterm formula for preterm or LBW infants.

Summary of judgements

Comparison: Preterm formula vs term formula (A.5)

Justification	In trials where most participants are very preterm (< 32 weeks' gestation) or VLBW (< 1.5 kg): <ul style="list-style-type: none">• Evidence of small benefits: increased in-hospital weight, head circumference, neurodevelopment (<i>low-certainty evidence</i>)• No evidence of harms• Evidence of little or no effect on mortality and necrotizing enterocolitis (<i>low-certainty evidence</i>)
----------------------	---

Evidence-to-Decision summary

Desirable	Small
Undesirable	Trivial or none
Certainty	Low
Balance	Probably favours preterm formula
Values	No uncertainty or variability about outcomes
Acceptability	Varies
Resources	Moderate
Feasibility	Probably not feasible
Equity	Probably not equitable

A.6 EARLY INITIATION OF ENTERAL FEEDING

Recommendation and remarks

RECOMMENDATION A.6 (UPDATED)

Preterm and low-birth-weight (LBW) infants, including very preterm (< 32 weeks' gestation) and very LBW (< 1.5 kg) infants, should be fed as early as possible from the first day after birth. Infants who are able to breastfeed should be put to the breast as soon as possible after birth. Infants who are unable to breastfeed should be given expressed mother's own milk as soon as it becomes available. If mother's own milk is not available, donor human milk should be given wherever possible. (*Strong recommendation, moderate-certainty evidence*)

Remarks

- Enteral feeding includes direct breastfeeding and feeding by cups, naso- or orogastric tubes.
- The trials included in the systematic review mostly did not state the stability of the babies, so careful consideration is needed in applying these recommendations to unstable babies. The GDG considers that initiation of enteral feeding in unstable babies should be based on clinical judgement.
- Infants should be given mother's own milk wherever possible. The provision of colostrum is especially important. If mother's own milk is not available, then donor human milk should be given wherever possible. If human milk is not available, infants can be fed formula as this is preferable to delayed initiation of enteral feeding and the use of parenteral nutrition.
- There was no difference in effectiveness by volume of initial feed, so a recommendation was not made on restricting the volume of feed.
- In all but one of the trials, the control group received parenteral nutrition. The benefits of early initiation of enteral feeding may be even greater when the alternatives are intravenous fluids or dextrose water rather than parenteral nutrition.

Background and definitions

WHO and UNICEF recommend early initiation of breastfeeding within 1 hour of birth for all healthy term infants (63). Clinicians continue to debate the optimal timing of feeding initiation for preterm and LBW infants for fear of potential health complications, including necrotizing enterocolitis (13,82,83). Additionally, women in communities

around the world may delay feeding due to the cultural practices of discarding colostrum, pain and discomfort after delivery, and concern about the developmental maturity of the infant, including the infant's inability to digest milk feeds (84,85). In 2011, WHO recommended early initiation of enteral feeding for stable preterm or LBW infants (19). However, there have been new studies since that time.

Summary of the evidence

OVERVIEW	A.6 Early initiation of enteral feeding
PICO	<p>Population – Preterm or LBW infants</p> <p>Intervention – Early initiation of enteral feeding (< 72 hours)</p> <p>Comparator – Delayed initiation of enteral feeding (> 72 hours)</p> <p>Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up</p>
Timing, setting, subgroups	<p>Timing of the intervention – Birth to 1 month of age</p> <p>Setting – Health-care facility or home in any country or setting</p> <p>Subgroups</p> <ul style="list-style-type: none"> • Gestational age at birth (< 32 weeks, ≥ 32 weeks) • Birth weight (< 1.5 kg, ≥ 1.5 kg) • Timing of feed initiation (days 1, 2, 3) • Milk volume (< 15 ml/kg per day, ≥ 15 ml/kg per day) • Milk type (human milk, formula, and mixed human milk with formula)

Effectiveness: Comparison – Early versus delayed initiation of enteral feeding

Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review of 14 trials enrolling 1511 preterm or LBW infants, which compared early initiation of enteral feeding (< 72 hours) with delayed initiation of enteral feeding (\geq 72 hours). The trials were from nine countries (Canada, Chile, Colombia, India, the Netherlands, Spain, the United Arab Emirates, the United Kingdom and the USA) (86). All trials were based in hospital NICUs. Ten trials restricted enrolment to very preterm infants (< 32 weeks' gestation) or VLBW infants (< 1.5 kg) and five enrolled all preterm or LBW infants. Three studies enrolled only small-for-gestational-age (SGA) infants. Early initiation time ranged from 1 to 3 days after birth and delayed initiation time ranged from 4 to 15 days after birth. Two studies initiated feeding by day 1 (i.e. < 24 hours), eight studies initiated by day 2 (i.e. < 48 hours) and five studies initiated by day 3 (i.e. < 72 hours). Enteral feed volumes ranged from 5 to 25 ml/kg per day. Only two studies provided babies with feed volumes > 15 ml/kg per day. Only one trial provided direct breastfeeding while the remaining 13 gave feeds by naso- or orogastric tube. Three studies gave the babies formula milk, one gave mother's own milk and the remaining 10 gave a mixture of milks (i.e. mother's own, donor human milk and/or formula). All infants received supplemental parenteral nutrition in the delayed initiation group, except for one study which did not specify.

Critical outcomes

For early feeding compared with delayed feeding for preterm or LBW infants, 12 studies reported all-cause mortality outcomes, 14 reported morbidity (14 reported necrotizing enterocolitis, 6 sepsis, 1 intraventricular haemorrhage), 7 reported growth outcomes (7 reported time to regain birth weight, 1 weight, 1 length and 3 head circumference). None of the trials reported on neurodevelopment. (Full details are provided in GRADE Table A.6, in the Web Supplement.)

- **Mortality:** Moderate-certainty evidence from 12 trials totalling 1292 participants suggests a decrease in all-cause mortality by hospital discharge (RR 0.69, 95% CI 0.48 to 0.99).
- **Morbidity:** Low-certainty evidence from 13 trials totalling 1484 participants suggests little or no effect on necrotizing enterocolitis by hospital discharge (RR 1.05, 95% CI 0.75 to 1.46). Low-certainty evidence from five trials totalling 626

participants suggests little or no effect on sepsis by discharge (RR 0.90, 95% CI 0.54 to 1.52).

Very-low-certainty evidence from one trial with 84 participants suggests a decrease in intraventricular haemorrhage by hospital discharge (RR 0.48, 95% CI 0.18 to 1.25).

- **Growth:** Low-certainty evidence from seven trials totalling 569 participants suggests little or no effect on time to regain birth weight (in days) (MD 0.26, 95% CI -0.63 to 1.15). Low-certainty evidence from three trials totalling 142 participants suggests little or no effect on weight (in grams) at latest follow-up (at chronological age 6–12 weeks) (MD -49.02, 95% CI -149.62 to 51.61). Very-low-certainty evidence from one trial with 40 participants suggests an increase in weight gain (in grams) from enrolment to 30 days follow-up (MD 51, 95% CI 32.4 to 69.6). Low-certainty evidence from two trials totalling 82 participants suggests little or no effect on length gain (in centimetres) at latest follow-up (at chronological age 32 weeks) (MD -0.62, 95% CI -1.51 to 0.27). Very-low-certainty evidence from two trials totalling 82 participants suggests little or no effect on head circumference (in centimetres) at latest follow-up (at discharge or chronological age 32 weeks) (MD -0.56, 95% CI -1.18 to 0.06).

Other outcomes

There was little or no effect on feed intolerance at discharge (RR 1.03, 95% CI 0.66 to 1.60; 2 trials, 187 participants) or length of hospital stay (days to discharge) (MD -3.2, 95% CI -5.74 to -0.66; 10 trials, 1100 participants).

Subgroup analyses

For the analyses by gestational age and birth weight, subgroup differences could not be assessed as there were insufficient studies on any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding what interventions are given to infants, including what and how they are fed (14). There have been studies of the barriers, facilitators, preferences, values and acceptability of early and late initiation of enteral feeding for preterm or LBW infants (84,85). Reasons for delay in initiation of feeding include cultural practices of discarding colostrum, pain and

discomfort after delivery, and concern about the developmental maturity of the baby, including the baby's inability to digest milk feeds. Reasons for early initiation include the importance of providing nurturing care to the baby as soon as possible, and concerns about the use of intravenous lines, dextrose water, total parenteral nutrition and lack of other nutritional support (84,85).

Resources required and implementation considerations

Organization of care

Early initiation of enteral feeding from the first day of life (the day of birth) can be implemented at home and at all levels of newborn care (primary, secondary and tertiary).

Infrastructure, equipment and supplies

National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can provide early initiation support to mothers and families. Standardized packages are needed for training, supervision and monitoring.

Feasibility and equity

There was no specific evidence on the feasibility and equity of early initiation of feeding for preterm or LBW infants.

Summary of judgements

Comparison: Early vs delayed initiation of enteral feeding (A.6)

Justification

- Evidence of moderate benefits: decreased mortality (*moderate-certainty evidence*), decreased length of hospital stay (*moderate-certainty evidence*), decreased intraventricular haemorrhage (*very-low-certainty evidence*)
- No evidence of harms
- Evidence of little or no effect on: necrotizing enterocolitis (*low-certainty evidence*), sepsis (*low-certainty evidence*), growth, i.e. time to regain birth weight, weight in grams, weight gain in grams, length at discharge (*low- to very-low-certainty evidence*), feed intolerance (*low-certainty evidence*)
- No evidence on other critical outcomes

Evidence-to-Decision summary

Benefits	Moderate
Harms	Trivial or none
Certainty	Moderate
Balance	Favours early initiation
Values	No uncertainty or variability about outcomes
Acceptability	Acceptable
Resources	Negligible
Feasibility	Feasible
Equity	Equitable

A.7 RESPONSIVE AND SCHEDULED FEEDING

Recommendation and remarks

RECOMMENDATION A.7 (UPDATED)

In health-care facilities, scheduled feeding may be considered rather than responsive feeding for preterm infants born before 34 weeks' gestation, until the infant is discharged. (Conditional recommendation, low-certainty evidence)

Remarks

- The recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- In making this decision, the GDG considered that the harms from responsive feeding (i.e. poor weight gain) outweighed the benefits (i.e. decreased length of hospital stay).
- Most data were about infants born before 34 weeks' gestation, so recommendations could not be made for infants born at or after 34 weeks' gestation.
- The included trials used a range of different feeding schedules and it was not possible to recommend a particular schedule. The GDG suggests 2–3 hourly scheduled feeding may be used for infants born before 34 weeks' gestation as this is a commonly used and feasible schedule.
- All studies were in hospitalized infants, so the GDG could not make a recommendation on feeding outside the hospital.
- Nurturing care and responsive caregiving are critical to the well-being of every preterm and LBW infant and should be implemented regardless of the type of feeding regime.

Background and definitions

Responsive feeding is often defined as feeding in response to infant visual and auditory cues (or signals) of hunger and satiety (87–89). Infant cues include crying, hand-mouth motions, suckling and awakesness. Scheduled feeding is defined in many studies as enteral feeding at regularly timed intervals, irrespective of infant cues (87–89). A 2016 Cochrane review suggested that responsive feeding led to

slower weight gain, but decreased the transition time from enteral tube to oral feeding (90). However, another systematic review reported that responsive feeding decreased the length of hospitalization and increased weight gain in infants (91). In 2011, WHO recommended that LBW infants who are orally fed but not breastfed should be fed based on infants' hunger cues, except when the infant remains asleep beyond three hours since the last feed (19).

Summary of the evidence

OVERVIEW	A.7 Responsive and scheduled feeding
PICO	Population – Preterm or LBW infants who receive any enteral feeding Intervention – Responsive feeding based on infant cues Comparator – Scheduled feeding Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none">• Gestational age at birth (< 32 weeks, ≥ 32 weeks)• Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – Responsive feeding versus scheduled feeding

Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review of eight RCTs reporting on 455 preterm or LBW infants from four countries (Canada, the Islamic Republic of Iran, Israel and the USA) (92). The studies were all conducted in NICUs and the responsive feeding was provided by health staff and not by families – that is, the health workers directly implemented a protocol of scheduled or responsive feeding regardless of whether a family member was present. The scheduled feeding regimes were mostly 2- to 3-hourly and the feeding volumes ranged from 120 to 180 ml/kg per day. The studies implemented the intervention for variable durations, with the minimum being 3 days and the maximum lasting until hospital discharge. Only one study recruited very preterm infants (< 32 weeks' gestation) while the remainder recruited preterm infants.

Critical outcomes

For responsive feeding compared with scheduled feeding for preterm or LBW infants, seven studies assessed growth outcomes (7 reported weight gain, 3 weight). No studies assessed mortality, morbidity or neurodevelopment outcomes. (Full details are provided in GRADE Table A.7, in the Web Supplement.)

- **Growth:** low-certainty evidence from two trials totalling 213 participants suggests a decrease in weight (in grams per day) by hospital discharge (MD -2.8, 95% CI -3.39 to -2.22). Low-certainty evidence from three trials totalling 183 participants suggests little to no effect on weight (in grams) by hospital discharge (MD -22.21, 95% CI -130.63 to 86.21). Very-low-certainty evidence from five trials totalling 372 participants suggests little to no effect on weight gain (in grams per kg per day) by hospital discharge (MD -0.99, 95% CI -2.45 to 0.46).

Other outcomes

Very-low-certainty evidence from three trials totalling 342 participants suggests a decrease in duration of hospitalization (days to discharge) (MD -1.42, 95% CI -5.43 to 2.59).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient studies.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding what interventions are given to infants, including what and how they are fed (14). No other specific evidence was located about whether families value responsive feeding more than scheduled feeding for their preterm or LBW baby or whether they find it more or less acceptable.

Resources required and implementation considerations

Organization of care

In facilities, infants born before 34 weeks' gestation can be fed every 2–3 hours. Infants born at 34 weeks' gestation or more can be fed every 3–4 hours or by responsive feeding. At home, there is no recommended scheduling; families and health workers can decide together, depending on clinical judgement and their preferences.

Infrastructure, equipment and supplies

National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can support mothers and families. Standardized packages are needed for training, supervision and monitoring.

Feasibility and equity

Administration of scheduled feeds for preterm and LBW babies varies markedly but common scheduling is 2- to 3-hourly feeding with volumes of 80–200 ml/kg per day for babies born before 34 weeks' gestation. Responsive feeding requires sensitivity and careful observation of the baby's behaviour and is more commonly implemented in settings with well staffed special care nurseries and NICUs (87). There was no specific evidence on the feasibility and equity of responsive and scheduled feeding for preterm or LBW infants.

Summary of judgements

Comparison: Responsive feeding vs scheduled feeding (A.7)

Justification

In trials where most participants are hospitalized infants born < 34 weeks' gestation:

- Evidence of small benefits from responsive feeding: decreased length of hospital stay (*very-low-certainty evidence*) in trials of infants born < 34 weeks' gestation
- Evidence of small harms from responsive feeding: decreased weight gain velocity in grams per day, and grams per kilogram per day (*low- to very-low-certainty evidence*), decreased weight gain in grams at discharge (*very-low-certainty evidence*)
- No evidence on other critical outcomes

Evidence-to-Decision summary

Benefits	Benefits of responsive feeding are trivial to none
Harms	Harms of responsive feeding are small
Certainty	Very low to low
Balance	Probably does not favour responsive feeding, probably favours scheduled feeding
Values	Uncertainty or variability about outcomes
Acceptability	Acceptability of responsive feeding and scheduled feeding varies
Resources	Resources needed for responsive feeding and scheduled feeding vary
Feasibility	Feasibility of responsive feeding and scheduled feeding varies
Equity	Equity of responsive feeding and scheduled feeding varies

A.8 FAST AND SLOW ADVANCEMENT OF FEEDING

Recommendation and remarks

RECOMMENDATION A.8 (UPDATED)

In preterm or low-birth-weight (LBW) infants, including very preterm (< 32 weeks' gestation) or very LBW (< 1.5 kg) infants, who need to be fed by an alternative feeding method to breastfeeding (e.g. gastric tube feeding or cup feeding), feed volumes can be increased by up to 30 ml/kg per day. (Conditional recommendation, moderate-certainty evidence)

Remarks

- The recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- The GDG noted that the trials enrolled infants immediately after birth (i.e. day 1 – within 24 hours of birth) so results are generalizable to very early feeding of LBW infants from this time.
- All trials excluded babies with congenital anomalies and birth asphyxia, so careful consideration is needed in applying these recommendations to infants with these conditions. Feed advancement should be based on clinical judgement for these infants.
- All trials compared fast advancement (increments of 30–40 ml/kg per day) with slow advancement (increments of 15–25 ml/kg per day). So the GDG took the conservative value of 30 ml/kg per day as the threshold for fast feed advancement. This value is also consistent with many national guidelines.
- All studies were in hospitalized infants, so the GDG could not make a recommendation on feeding outside the hospital.
- The GDG did not make separate recommendations for babies fed formula milk versus human milk as there was insufficient evidence (only one trial gave formula as the sole diet while the remainder gave human milk only or a mix of human milk and formula).
- The GDG considered that advancement should continue until full maintenance feed volumes are reached. These volumes should be based on local guidelines.
- The GDG noted that further research is needed to understand the neurodevelopmental effects of fast feed advancement.

Background and definitions

There is substantial variation in the definitions of fast and slow advancement of enteral feeding volumes for preterm and LBW babies in the first weeks after birth. Advancement increments commonly vary between 10 and 40 ml/kg per day (93,94). Up to the 1990s, the standard of care was a conservative (“slow rate”) approach because of concerns about

feed intolerance (e.g. gagging, vomiting and apnoea post-feed) and necrotizing enterocolitis (56). In 2011, WHO recommended that feeds could be advanced by up to 30 ml/kg per day with careful monitoring for feed intolerance in infants weighing under 1.5 kg (19). However, there have been new studies published since that time (95).

Summary of the evidence

OVERVIEW	A.8 Fast and slow advancement of feeding
PICO	Population – Preterm or LBW infants Intervention – Fast advancement of enteral feeds (≥ 30 ml/kg per day) Comparator – Slow advancement of enteral feeds (< 30 ml/kg per day) Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none"> • Gestational age at birth (< 32 weeks, ≥ 32 weeks) • Birth weight (< 1.5 kg, ≥ 1.5 kg) • Type of milk (human milk, formula milk)

Effectiveness: Comparison – Fast versus slow advancement of enteral feeds

Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review of 12 RCTs enrolling 4084 preterm or LBW infants (96). The trials were conducted in Bangladesh, Colombia, India, the Islamic Republic of Iran, Ireland, South Africa, Türkiye, the United Kingdom and the USA. The United Kingdom Speed of Increasing Milk Feeds trial (SIFT) was the largest trial ($n=2973$) (97). Most studies included clinically stable infants and excluded those with perinatal asphyxia or haemodynamic instability. The infants were typically randomized on days 1–4 after birth. Intervention (fast advancement) increments ranged from 30 to 40 ml/kg per day. Comparator (slow advancement) increments ranged from 10 to 25 ml/kg per day. The target volume of full feeding ranged from 120 to 180 ml/kg per day. Seven studies enrolled very preterm infants born before 32 weeks' gestation. Three studies used human milk, one used infant formula, and seven used a combination of the two.

Critical outcomes

For fast compared with slow advancement of enteral feeding for preterm or LBW infants, 11 trials reported all-cause mortality, 12 reported morbidity (12 reported necrotizing enterocolitis, 9 sepsis, 2 apnoea), 6 reported growth outcomes (6 reported time to regain birth weight, 1 WAZ at discharge, 1 weight at discharge, 1 weight gain, 1 head circumference) and 1 reported neurodevelopmental outcomes (disability). (Full details are provided in GRADE Table A.8, in the Web Supplement.)

■ **Mortality:** Moderate-certainty evidence from 11 trials with a total of 4132 participants suggests little or no effect on all-cause mortality by hospital discharge (RR 0.93, 95% CI 0.73 to 1.18).

■ **Morbidity:** Low-certainty evidence from two trials totalling 153 participants suggests a decrease in apnoea by hospital discharge (RR 0.72, 95% CI 0.47 to 1.12). Moderate-certainty evidence from 12 trials totalling 4291 participants suggests little or no effect on necrotizing enterocolitis by hospital discharge (RR 0.89, 95% CI 0.68 to 1.15). Moderate-certainty evidence from nine trials totalling 3648 participants suggests little or no effect on sepsis by hospital discharge (RR 0.92, 95% CI 0.83 to 1.03).

■ **Growth:** High-certainty evidence from six trials totalling 993 participants suggests a decrease in time to regain birth weight by hospital discharge (MD -3.69, 95% CI -4.44 to -2.95). Low-certainty evidence from one trial with 2793 participants suggests little or no effect on WAZ by hospital discharge (MD 0.0, 95% CI -0.08 to 0.08). Low-certainty evidence from one trial with 131 participants suggests little or no effect on weight gain (in grams per kilogram per day) by hospital discharge (MD 0.5, 95% CI -1.19 to 2.19). Low-certainty evidence from one trial with 100 participants suggests little or no effect on weight in grams by hospital discharge (MD -29.0, 95% CI -74.89 to 16.89). Low-certainty evidence from one trial with 2793 participants suggests little or no effect on head circumference (head circumference z score) by hospital discharge (MD -0.1, 95% CI -0.22 to 0.02).

■ **Neurodevelopment:** Low-certainty evidence from one trial of 2325 participants suggests little or no effect on neurodevelopment (neurodevelopmental disability measured using a validated test) at 24 months corrected age (RR 1.12, 95% CI 0.98 to 1.27).

Other outcomes

There was a decrease in length of hospital stay (days to discharge) (MD -3.08, 95% CI -4.34 to -1.81; 7 trials, 3864 participants) and little or no effect on feed intolerance by hospital discharge (RR 0.92, 95% CI 0.77 to 1.10; 8 trials, 1114 participants).

Subgroup analyses

No subgroup differences were seen for gestational age and birth weight for any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding what interventions are given to infants, including what and how they are fed (14). No specific evidence was located about whether families value fast versus slow feed advancement for their preterm or LBW baby or whether they find the different rates more or less acceptable.

Resources required and implementation considerations

Organization of care

Feed advancement should be based on clinical judgement for all infants at home and in health-care facilities. In facilities, there can be fast advancement of feed volumes by up to 30 ml/kg per day.

Infrastructure, equipment and supplies

National or local guidance on infrastructure, equipment and supplies for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can support mothers and families. Standardized packages are needed for training, supervision and monitoring.

Feasibility and equity

No specific evidence was located about the feasibility and equity of providing slow or fast feed advancement to preterm or LBW babies.

Summary of judgements

Comparison: Fast vs slow advancement of enteral feeds (A.8)

Justification

- Evidence of moderate benefits: decrease in apnoea (*moderate-certainty evidence*), decrease in time to regain birth weight (*high-certainty evidence*), decreased length of hospital stay (*moderate-certainty evidence*)
- Evidence on harms uncertain: impaired neurodevelopment (*low-certainty evidence*)
- Evidence of little or no effect on: mortality, necrotizing enterocolitis, sepsis, weight gain, head circumference (*low-certainty evidence*), feed intolerance (*moderate-certainty evidence*)

Evidence-to-Decision summary

Desirable	Small
Undesirable	Unknown
Certainty	Moderate
Balance	Probably favours fast feed advancement
Values	Uncertainty or variability about outcomes
Acceptability	Probably acceptable
Resources	Negligible
Feasibility	Probably feasible
Equity	Equitable

A.9 DURATION OF EXCLUSIVE BREASTFEEDING

Recommendation and remarks

RECOMMENDATION A.9 (UPDATED)

Preterm or low-birth-weight infants should be exclusively breastfed until 6 months of age. (*Strong recommendation, very-low-certainty evidence*)

Remarks

- The GDG made strong recommendation in favour of exclusive breastfeeding (EBF) until 6 months of age despite the very-low-certainty evidence because they considered the potential harms of less than 6 months of EBF to outweigh the potential harms of having at least 6 months of EBF.
- In making the decision, the GDG also considered the results of a systematic review of 42 studies (89 638 infants) comparing mother's own milk with infant formula in babies aged 0–6 months (60). This review showed consistent harm from the use of infant formula on a critical outcome (morbidity: necrotizing enterocolitis) in the first 6 months after birth. It also reported no evidence of benefit from infant formula over the same period.
- The GDG also considered that EBF until 6 months of age is the standard of care for preterm and LBW infants across many high-, middle- and low-income countries and is the foundation of many national policies and programmes.
- The GDG also felt that mothers should be encouraged and supported before and after birth to provide their own breast-milk (including colostrum) for their infants.

Background and definitions

WHO defines exclusive breastfeeding (EBF) as feeding no other foods or fluids (not even water) except breast-milk, medicines, vitamins and minerals (22). EBF until 6 months of age is recommended for full-term, normal-birth-weight infants (22). However, preterm and LBW infants are more vulnerable

to nutritional deficiencies (13,56). The risks of contamination of complementary foods and early infant formula feeding are also well known (98). In 2011, WHO recommended EBF until 6 months of age for preterm and LBW babies (19), but new studies have been published since that time.

Summary of the evidence

OVERVIEW	A.9 Duration of exclusive breastfeeding (EBF)
PICO	Population – Preterm or LBW infants Intervention – EBF to < 6 months of age Comparator – EBF until 6 months of age Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none">• Gestational age at birth (< 32 weeks, ≥ 32 weeks)• Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – Exclusive breastfeeding for less than six months versus for six months

Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review of two RCTs reporting on a total of 307 preterm or LBW infants from two countries (Honduras and India) (99). The trial in Honduras

randomized 119 term SGA EBF infants (mean birth weight in the intervention group was 2364 g [SD 137], mean birth weight in the control group was 2327 g [SD 183]) to receive nutrient-rich complementary foods starting from 4 months chronological age. The other study in India randomized 403 infants born before 34 weeks' gestation (mean birth weight in the intervention group was 1479 g [SD 308],

mean birth weight in the control group was 1492 g [SD 344]) to receive nutrient-rich complementary foods starting from 4 months corrected age. Fifty per cent (202/403) of these infants were EBF (104 intervention and 98 control) and 93% (188/202) of those EBF infants had WAZ outcome data (95 intervention and 93 control).

Critical outcomes

For EBF less than six months compared with EBF for six months for preterm or LBW infants, one trial reported morbidity (percentage of days with diarrhoea and/or fever), two trials reported growth outcomes (1 reported weight gain, 1 WAZ, 1 length gain) and one trial reported neurodevelopment (time to achieve motor developmental milestones). No trials reported mortality. (Full details are provided in GRADE Table A.9, in the Web Supplement.)

- **Morbidity:** Very-low-certainty evidence from one trial with 119 participants suggests a decrease in the percentage of days with diarrhoea from 16 to 26 weeks of chronological age (MD -2.6, 95% CI -5.2 to 0.0). Very-low-certainty evidence from one trial with 119 participants suggests little or no effect on the percentage of days with fever from 16 to 26 weeks chronological age (MD -0.7, 95% CI -3.4 to 2.0).
- **Growth:** Very-low-certainty evidence from one trial with 119 participants suggests a decrease in weight gain (in grams) from 4 to 6 months of chronological age (MD -13, 95% CI -143 to 117). Low-certainty evidence from one trial with 188 participants suggests little or no effect on WAZ at 12 months corrected age (MD 0.1, 95% CI -0.2 to 0.4). Very-low-certainty evidence from one trial with 119 participants suggests a decrease in the rate of length gain (in centimetres) from 4 to 6 months of chronological age (MD -0.2, 95% CI -0.6 to 0.2).
- **Neurodevelopment:** Very-low-certainty evidence from one trial with 108 participants suggests little or no effect on motor development milestones at specified chronological ages (in months) (raise head, MD 0.0, 95% CI -0.3 to 0.3; raise chest, MD -0.1, 95% CI -0.7 to 0.5; roll over, MD 0.0, 95% CI -0.7 to 0.7; able to crawl, MD 0.6, 95% CI -0.1 to 1.3; able to sit from lying position, MD 0.6, 95% CI 0.0 to 1.2). Very-low-certainty evidence from one trial with 99 participants suggests an increase in the percentage of infants who can walk by the chronological age of 12 months (RR 1.47, 95% CI 0.69 to 3.13).

Other outcomes

There was a decrease in anaemia (haemoglobin level < 10.5 g/dl) (RR 0.10, 95% CI 0.01 to 0.77, 1 trial, 104 participants) but not in infants who received iron supplements (RR 1.07, 95% CI 0.22 to 5.28; 1 trial, 29 participants).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding what interventions are given to infants, including what and how they are fed (14).

There are studies that report the difficulties in providing mother's own milk when the mother and baby return home, including difficulties balancing work commitments, maternity leave, night-time feeding and father/partner support (14). There are also studies that report family concerns with infant formula, including concerns about nutrient composition, water supply, contamination and cost (64,65). Studies also report families valuing having formula available if their circumstances demand it, such as work commitments, maternity leave, night-time feeding, father and partner support (64,65). No specific evidence was located about whether families value EBF for up to 6 months of age for their preterm or LBW baby or whether they find the different durations of EBF more or less acceptable.

Resources required and implementation considerations

Organization of care

Promotion of exclusive breastfeeding for six months should be done at the community and facility level and be integrated within standard national programmes. This should occur throughout the antenatal and postnatal periods and up until the infant reaches 6 months of age.

Infrastructure, equipment and supplies

National or local guidance for infrastructure, equipment and supplies for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can promote exclusive breastfeeding for six months. Standardized packages are needed for training, supervision and monitoring.

Feasibility and equity

There was no specific evidence on the feasibility and equity of duration of EBF for preterm or LBW infants.

Summary of judgements

Comparison: Exclusive breastfeeding (EBF) for less than six months vs for six months (A.9)

Justification

- Evidence of small benefits: decrease in percentage of days with diarrhoea (*very-low-certainty evidence*), increase in neurodevelopment, i.e. percentage of infants who can walk by the age of 12 months (*very-low-certainty evidence*)
- Evidence of small harms: decrease in weight gain in grams at 26 weeks (*very-low certainty evidence*)
- Evidence of little or no effect on other morbidity (percentage of days with fever), other growth (weight-for-age z score [WAZ], length in centimetres), and other neurodevelopmental milestones (raise head, raise head and chest, roll over, crawl, sit from lying position) (*very-low-certainty evidence*)
- No evidence on other critical outcomes

Evidence-to-Decision summary

Benefits	Benefits of EBF to < 6 months are small
Harms	Harms of EBF to < 6 months are small
Certainty	Very low
Balance	Does not favour EBF to < 6 months, favours EBF to 6 months
Values	Uncertainty or variability about outcomes
Acceptability	Acceptability of EBF to < 6 months varies
Resources	Resources for EBF to < 6 months are low to moderate
Feasibility	Feasibility of EBF to < 6 months varies
Equity	Equity of EBF to < 6 months varies

A.10 MICRONUTRIENT SUPPLEMENTATION

A.10a Iron supplementation

Recommendation and remarks

RECOMMENDATION A.10a (UPDATED)

Enteral iron supplementation is recommended for human milk-fed preterm or low-birth-weight infants who are not receiving iron from another source. (*Strong recommendation, moderate-certainty evidence*)

Remarks

- The GDG noted that there were limited data on dose, timing of initiation and duration of iron supplementation.
- Based on most trials included in the evidence review, the GDG suggests a daily dose of 2–4 mg/kg per day of elemental iron may be initiated when enteral feeds are well established, and may be continued until the infant receives iron from another source.

Background and definitions

Iron deficiency is associated with poor growth and development outcomes in term and preterm babies (100,101). Human milk may not meet the nutritional requirements of preterm or LBW infants because of their low iron stores, red blood cell expansion, catch-up growth and iatrogenic blood loss. The most recent systematic reviews of RCTs and non-randomized studies reported that enteral iron supplementation may improve haematological

outcomes in preterm and LBW babies but that there was insufficient evidence to assess effects on growth and neurodevelopmental outcomes (100,101). The optimal dose, optimal timing of initiation and the level and types of morbidity associated with iron supplementation were also unclear. In 2011, WHO recommended that VLBW infants fed mother's own milk or donor human milk should be given iron supplementation of 2–4 mg/kg per day starting at 2 weeks and continuing until 6 months of age (19).

Summary of the evidence

OVERVIEW	A.10a Iron supplementation
PICO	Population – Preterm or LBW infants who are fed mother's own milk or donor human milk Intervention – Iron supplementation Comparator – No iron supplementation Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none"> • Gestational age at birth (< 32 weeks, ≥ 32 weeks) • Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – Iron supplementation versus no iron supplementation

Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review of eight trials (11 publications) reporting on a total of 1093 infants from seven countries (Canada, Germany, India, the Netherlands, Sweden, the United Kingdom and the USA) (102). Most trials enrolled babies with birth weight below 1.5 kg or born before 32 weeks' gestation. The trials used iron supplementation

doses ranging from 1 to 7 mg/kg per day, median 2.2 (IQR 1.97–2.55) mg/kg per day. Supplementation commenced between 14 and 56 days chronological age. The mean duration of supplementation was 81 (SD 57) days and the median duration was 53 (IQR 40–98) days. One trial gave iron with multivitamin supplements and compared this with infants who received multivitamins alone. The remaining seven trials gave iron supplementation alone and compared this with placebo or no iron supplementation.

Critical outcomes

For enteral iron supplementation compared with no iron supplementation, four trials reported morbidity (4 reported sepsis, 2 necrotizing enterocolitis), five reported growth outcomes (5 reported weight, 3 length, 3 head circumference) and one reported on neurodevelopment (cognitive outcomes). No studies reported all-cause mortality. (Full details are provided in GRADE Table A.10, in the Web Supplement.)

■ **Morbidity:** Very-low-certainty evidence from four trials totalling 270 participants suggests little or no effect on sepsis prevalence at latest follow-up (median 8 [IQR 8–9] weeks) (RR 1.08, 95% CI 0.56 to 2.07). Very-low-certainty evidence from two trials totalling 194 participants suggests little or no effect on necrotizing enterocolitis prevalence at latest follow-up (median 9 [IQR 8.5–9.5] weeks) (RR 1.54, 95% CI 0.69 to 3.46).

■ **Growth:** Low-certainty evidence from five trials totalling 574 participants suggests an increase in weight in grams at latest follow-up (median 26 [IQR 8–36] weeks) (MD 35.31, 95% CI -64.53 to 135.15). Moderate-certainty evidence from three trials totalling 384 participants suggests an increase in length in centimetres at latest follow-up (median 26 [IQR 8–183] weeks) (MD 0.69, 95% CI 0.01 to 1.37). Low-certainty evidence from three trials totalling 385 participants suggests little or no effect on head circumference at latest follow-up (median 26 [IQR 8–183] weeks) (MD 0.09, 95% CI -0.4 to 0.21).

■ **Neurodevelopment:** Very-low-certainty evidence from one trial with 199 participants suggests little or no effect on cognitive development (measured using the Wechsler Intelligence Scale for Children, fourth edition [WISC-IV]) at latest follow-up (mean 365 weeks) (RR 0.31, 95% CI 0.09 to 1.02).

Other outcomes

Moderate-certainty evidence from two trials totalling 381 participants suggests a decrease in anaemia prevalence at latest follow-up (RR 0.25, 95% CI 0.10 to 0.62). Moderate-certainty evidence from five trials totalling 506 participants suggests an increase in haemoglobin prevalence at latest follow-up (mean 26 weeks) (MD 4.79, 95% CI 2.9 to 6.69). Very-low-certainty evidence from six trials totalling 607 participants suggests little or no effect on ferritin levels at latest follow-up (median 14 [IQR 8–26] weeks) (MD 8.76, 95% CI -0.85 to 18.37). Very-low-certainty evidence from two trials totalling 238 participants suggests little or no effect on feed

intolerance at latest follow-up (mean 8 weeks) (RR 1.05, 95% CI 0.49 to 2.27).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding what interventions are given to infants, including what and how they are fed (14). There was no specific evidence available about whether families value iron supplements for their preterm or LBW baby or whether they find them acceptable.

Resources required and implementation considerations

Organization of care

The supplements can be provided in the health-care facility or at home. The family needs accurate information on the dose and how to administer the supplement. National or local guidance for health-care facilities should be used.

Infrastructure, equipment and supplies

Iron supplements are commonly provided to LBW and preterm infants as oral liquid solution. Infants are commonly prescribed 2–4 mg/kg of elemental iron per day for the prophylaxis of iron deficiency anaemia. Concentrations of 5 mg of elemental iron per millilitre of liquid are often used (e.g. 1 ml/day to a 2 kg baby will provide 2.5 mg of elemental iron per day). Droppers or syringes can be used to administer the supplement to the infant. Doses are different for the treatment of iron deficiency anaemia. National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can support mothers and families. Standardized packages are needed for training, supervision and monitoring. Dispensing needs to be documented in clinical records.

Feasibility and equity

There was no specific evidence available about the feasibility of providing iron supplements to preterm or LBW babies.

Summary of judgements

Comparison: Iron supplementation vs no iron supplementation (A.10a)

Justification	<ul style="list-style-type: none"> • Evidence of small-to-moderate benefit: decreased anaemia, increased weight and length (<i>low-certainty evidence</i>) • No evidence of harms • Evidence of little or no effect on sepsis and necrotizing enterocolitis (<i>very-low-certainty evidence</i>), and on weight, head circumference and neurodevelopment (<i>low-certainty evidence</i>) • No evidence on other critical outcomes
----------------------	---

Evidence-to-Decision summary

Benefits	Small to moderate
Harms	Trivial or none
Certainty	Moderate
Balance	Favours iron supplementation
Values	No uncertainty or variability about outcomes
Acceptability	Probably acceptable
Resources	Low to moderate
Feasibility	Probably feasible
Equity	Probably equitable

A.10b Zinc supplementation

Recommendation and remarks

RECOMMENDATION A.10b (UPDATED)

Enteral zinc supplementation may be considered for human milk-fed preterm or low-birth-weight infants who are not receiving zinc from another source. (*Conditional recommendation, low-certainty evidence*)

Remarks

- The GDG noted that the evidence on harms (decreased neurodevelopment) was uncertain due to very-low-certainty evidence and imprecision.
- The recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- The GDG noted that there were limited data on the dose, timing of initiation and duration of supplementation. Based on most trials included in the evidence review, the GDG suggests a daily dose of 1–3 mg/kg per day of elemental zinc. The GDG also suggests that zinc may be initiated when enteral feeds are well established, and may be continued until the infant receives zinc from another source.

Background and definitions

Zinc is a trace element essential for physiological functions of the human body (103). Zinc deficiency is associated with dysfunction in epidermal, gastrointestinal, central nervous, immune, skeletal and reproductive systems (104,105). Human milk may not be able to meet the nutritional requirements of preterm or LBW infants because of their low zinc stores and catch-up growth (104–106). A recent (2021) Cochrane review of enteral zinc supplementation in hospitalized preterm infants

fed any type of milk (i.e. infant formula or human milk) reported that zinc supplementation reduced all-cause mortality and was associated with a probable improvement in short-term weight gain and linear growth, but had little or no effect on common morbidities of prematurity (107). However, there have been no recent systematic reviews of zinc supplementation in babies born at home or in the hospital or on babies fed human milk only. The optimal dose and timing of initiation are also unclear.

Summary of the evidence

OVERVIEW	A.10b Zinc supplementation
PICO	Population – Preterm or LBW infants who are fed mother’s own milk or donor human milk Intervention – Zinc supplementation Comparator – No zinc supplementation Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none">• Gestational age at birth (< 32 weeks, ≥ 32 weeks)• Birth weight (< 1.5 kg, ≥ 1.5 kg)• Dose of elemental zinc (< 3 mg/day, 3–5 mg/day and > 5 mg/day)

Effectiveness: Comparison – Zinc supplementation versus no zinc supplementation Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review of 14 RCTs totalling 9940 preterm or LBW infants from 11 countries (Bangladesh, Brazil, Chile, Egypt, India, the Islamic Republic of Iran, Italy, the Republic of Korea, Nepal, Spain and the United

Republic of Tanzania) (108). Most infants had a birth weight of at least 1.5 kg or were born at 32 weeks’ gestation or later. Among these, two large RCTs assessed the effects of zinc supplementation in a total of 2748 term LBW infants in Brazil and India. Zinc supplementation dosages across all 14 RCTs ranged from 1 mg/day up to 10 mg/day and commenced between birth and 35 days of age. Most

studies used a zinc dose of 3–5 mg/day. The mean duration of supplementation was 182 (SD 142) days and the median duration was 141 (IQR 98–183) days.

Critical outcomes

For zinc supplementation compared with no zinc supplementation, six trials reported all-cause mortality, six reported morbidity (2 reported hospitalization, 6 diarrhoea, 2 acute respiratory infection, 2 sepsis), eight reported growth outcomes (8 reported weight gain, 6 length gain, 5 head circumference) and two reported neurodevelopment (MDI and PDI [BSID-II]). (Full details are provided in GRADE Table A.11, in the Web Supplement.)

- **Mortality:** Low-certainty evidence from six trials totalling 8801 participants suggests a decrease in all-cause mortality at latest follow-up (median 26 [IQR 14–152.1] weeks) (RR 0.73, 95% CI 0.46 to 1.16). There was a similar effect on all-cause mortality when the two trials with term LBW infants were excluded (RR 0.68, 95% CI 0.43 to 1.09; 4 trials, 7192 participants).
- **Morbidity:** Moderate-certainty evidence from six trials totalling 1947 participants suggests a decrease in diarrhoea (events) at latest follow-up (median 26 [IQR 20.1–52.1] weeks) (RR 0.81, 95% CI 0.68 to 0.97). Very-low-certainty evidence from two trials totalling 172 participants suggests a decrease in acute respiratory infection at latest follow-up (median 13 [IQR 6–20] weeks) (RR 0.32, 95% CI 0.09 to 1.17). Low-certainty evidence from two trials totalling 265 participants suggests little to no effect on sepsis at latest follow-up (median 17 [IQR 14 to 20] weeks) (RR 1.12, 95% CI 0.62 to 2.02).
- **Growth:** Moderate-certainty evidence from 8 trials totalling 798 participants suggests an increase in weight (in grams) at latest follow-up (median 22 [IQR 13.5–39] weeks) (MD 378.57, 95% CI 275.26 to 481.88). Low-certainty evidence from six trials totalling 529 participants suggests an increase in length (in centimetres) at latest follow-up (median 36.1 [IQR 20–52.1] weeks) (MD 2.92, 95% CI 1.53 to 4.31). Low-certainty evidence from five trials totalling 466 participants suggests an increase in head growth (in centimetres) at latest follow-up (median 20 [IQR 13–24] weeks) (MD 0.56, 95% CI 0.23 to 0.9).
- **Neurodevelopment:** Very-low-certainty evidence from two trials totalling 301 participants suggests a decrease in MDI (BSID-II) scores at latest follow-up (52 weeks) (MD -4.18, 95% CI -1.85 to -6.51). Very-low-certainty evidence from two trials

totalling 301 participants suggests an increase in PDI (BSID-II) scores at latest follow-up (52 weeks) (MD 5.75, 95% CI -4.83 to 16.33).

Other outcomes

There was a decrease in hospitalization (at least one hospitalization) at latest follow-up (RR 0.70, 95% CI 0.24 to 2.00; 2 trials, 277 participants).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome. For the dose of elemental zinc, no subgroup differences were seen for any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding what interventions are given to infants, including what and how they are fed (14). There was no specific evidence available about whether families value zinc supplements for their preterm or LBW baby or whether they find them acceptable.

Resources required and implementation considerations

Organization of care

The supplements can be provided in the health-care facility or at home. The family needs accurate information on the dose and how to administer the supplement. National or local guidance for health-care facilities should be used.

Infrastructure, equipment and supplies

Zinc supplements are often provided as either 5 mg zinc capsules that are then opened and mixed with 5 ml of water (1 mg elemental zinc per ml) or zinc-containing multivitamin syrups (5 mg elemental zinc in 120 mls) (i.e. 42 µg elemental zinc per ml). Babies are often prescribed 1–5 mls of these formulations daily. Droppers or syringes can be used to administer the supplement to the infant. National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can support mothers and families. Standardized packages are needed for training, supervision and monitoring. Dispensing needs to be documented in clinical records.

Feasibility and equity

There was no specific evidence available about the feasibility and equity of providing zinc supplements to preterm or LBW babies.

Summary of judgements

Comparison: Zinc supplementation vs no zinc supplementation (A.10b)

Justification

- Evidence of small-to-moderate benefit: decreased mortality (*low-certainty evidence*), decreased diarrhoea (*moderate-certainty evidence*), decreased respiratory infection (*very-low-certainty*), increased weight, length, head circumference (*moderate-certainty evidence*) and increased psychomotor development scores (*very-low-certainty evidence*)
- Evidence on harms uncertain: decreased mental development scores (*low-certainty evidence*)
- Evidence of little or no effect on sepsis (*low-certainty evidence*)
- No evidence on other critical outcomes

Evidence-to-Decision summary

Benefits	Small to moderate
Harms	Unknown
Certainty	Low
Balance	Probably favours zinc supplementation
Values	Uncertainty or variability about outcomes
Acceptability	Probably acceptable
Resources	Low to moderate
Feasibility	Probably feasible
Equity	Probably equitable

A.10c Vitamin D supplementation

Recommendation and remarks

RECOMMENDATION A.10c (UPDATED)

Enteral vitamin D supplementation may be considered for human milk-fed preterm or low-birth-weight infants who are not receiving vitamin D from another source. (*Conditional recommendation, low-certainty evidence*)

Remarks

- The GDG noted that the evidence on harms (increased mortality) was uncertain due to low-certainty evidence and imprecision.
- The recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- The GDG also noted improvements in vitamin D deficiency and alkaline phosphatase levels but there were no data on other markers of bone health such as osteopenia or rickets.
- The GDG noted that there were limited data on the dose, timing of initiation and duration of supplementation. Based on most trials included in the evidence review, the GDG suggests a daily dose of 400–800 IU may be initiated when enteral feeds are well established, and may be continued until the infant receives vitamin D from another source.

Background and definitions

Vitamin D increases intestinal absorption of calcium and phosphorus, and enhances bone mineralization (42). Low vitamin D levels are associated with seizures, irritability, rickets (swollen, deformed, painful joints and bones), bone fractures, osteopenia (radiological evidence of thin bones) and metabolic bone disease (radiological evidence of widened or deformed bones) (109–111). Vitamin D deficiency has also been associated with increased risk of respiratory and diarrhoeal disease. Human milk may not be able to meet the nutritional requirements of preterm or LBW infants because of their low vitamin

D stores and catch-up growth (56). Babies born to darker-skinned mothers are at higher risk of vitamin D deficiency, especially those born in higher latitudes and in the winter months (112). In 2011, WHO recommended that VLBW infants with birth weight below 1.5 kg should be given vitamin D supplements (400–1000 IU per day) until 6 months of age (19). A systematic review published in 2020 reported improvements in vitamin D biomarkers (vitamin D levels, calcium levels, parathyroid hormone) after vitamin D supplementation was provided to all preterm infants (113).

Summary of the evidence

OVERVIEW	A.10c Vitamin D supplementation
PICO	<p>Population – Preterm or LBW infants who are fed mother's own milk or donor human milk</p> <p>Intervention – Vitamin D supplementation</p> <p>Comparator – No vitamin D supplementation</p> <p>Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up</p>
Timing, setting, subgroups	<p>Timing of the intervention – Birth to 6 months of age</p> <p>Setting – Health-care facility or home in any country or setting</p> <p>Subgroups</p> <ul style="list-style-type: none"> • Gestational age at birth (< 32 weeks, ≥ 32 weeks) • Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – Vitamin D supplementation versus no vitamin D supplementation

Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review of three RCTs totalling 2479 preterm or LBW infants from two countries (India and the USA) (114). One trial in India enrolled 2079 (84%) of these infants. Most had a birth weight of at least 1.5 kg or were born at 32 weeks' gestation or later. The trials used a dose of vitamin D supplementation ranging from 200 IU to 800 IU per day. Two trials compared vitamin D with placebo, while the third trial compared vitamin D plus multivitamins with multivitamins alone. Vitamin D supplementation began between birth and 7 days chronological age in all trials. The mean duration of supplementation was 19 (SD 19) days and the median duration was 26 (IQR 4 to 26) days.

Critical outcomes

For vitamin D supplementation compared with no vitamin D supplementation, two trials reported all-cause mortality, five reported morbidity (1 reported bronchopulmonary dysplasia, 1 reported "at least one serious morbidity"), two reported growth (2 reported WAZ, 2 LAZ/HAZ, 1 head circumference z scores) and two reported neurodevelopment (cognitive development and neurodevelopmental impairment). (Full details are provided in GRADE Table A.10c, in the Web Supplement.)

■ **Mortality:** Low-certainty evidence from two trials totalling 2179 participants suggests an increase in all-cause mortality at latest follow-up (RR 1.81, 95% CI 0.92 to 3.56).

■ **Morbidity:** Very-low-certainty evidence from one trial with 100 participants suggests a decrease in bronchopulmonary dysplasia at 8 weeks of age (RR 0.77, 95% CI 0.47 to 1.27). Very-low-certainty evidence from two trials totalling 2179 participants suggests little or no effect on any (at least one) serious morbidity at latest follow-up (median 17 [IQR 8–26] weeks) (RR 0.94, 95% CI 0.72 to 1.24). "At least one serious morbidity" is defined as any (at least one) serious morbidity assessed with: any severe morbidity (hospital admission, or outpatient visits with diagnoses selected based on clinical judgement that represented severe illness: pneumonia, persistent diarrhoea, dysentery, severe fever, severe protein energy malnutrition, ear infections, meningitis and septicaemia), RDS, early-onset sepsis (\leq 72 hours), late-onset sepsis

(> 72 hours) and culture-positive meningitis (115).

■ **Growth:** Moderate-certainty evidence from one trial with 1273 participants suggests an increase in WAZ at 6 months (MD 0.12, 95% CI 0.01 to 0.23). Low-certainty evidence from one trial with 912 participants suggests little or no effect on WAZ scores between 3 and 6 years of age (MD -0.07, 95% CI -0.18 to 0.04). Moderate-certainty evidence from one trial with 1258 participants suggests an increase in LAZ at 6 months (MD 0.12, 95% CI 0.03 to 0.21). Low-certainty evidence from one trial with 912 participants suggests little or no effect on height-for-age z scores (HAZ) between 3 and 6 years of age (MD 0.07, 95% CI -0.05 to 0.19). Low-certainty of evidence from one trial with 1259 participants suggests little or no effect on head circumference z scores at 6 months (MD -0.08, 95% CI -0.17 to 0.01).

■ **Neurodevelopment:** Very-low-certainty evidence from one trial with 70 participants suggests little or no effect on cognitive scores assessed at 104 weeks (RR 0.85, 95% CI 0.45 to 1.59). Very-low-certainty evidence from one trial with 71 participants suggests a decrease in neurodevelopmental impairment assessed at 104 weeks (RR 0.69, 95% CI 0.41 to 1.17).

Other outcomes

There was little or no effect on hospitalization (at least one hospitalization) at latest follow-up (6 months) (RR 0.84, 95% CI 0.42 to 1.66; 2 trials, 1468 participants). There was a decrease in serum alkaline phosphatase (ALP) (measured in IU per litre) (note: ALP should be \geq 500 IU/L) at 6 months follow-up (RR 0.37, 95% CI 0.10 to 1.36; 1 trial, 265 participants). There was a decrease in vitamin D deficiency (< 20 μ g/ml) at latest follow-up (6 months) (RR 0.58, 95% CI 0.49 to 0.68; 2 trials, 504 participants).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials reporting on any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding what interventions are given to infants, including

what and how they are fed (14). There was no specific evidence available about whether families value vitamin D supplements for their preterm or LBW baby or whether they find them acceptable.

Resources required and implementation considerations

Organization of care

The supplements can be provided in the health-care facility or at home. The family needs accurate information on the dose and how to administer the supplement. National or local guidance for health-care facilities should be used.

Infrastructure, equipment and supplies

Common methods of providing enteral vitamin D for preterm and LBW infants include infant multivitamin

formulations (e.g. vitamins D, A, C, B group).

Many formulations contain 400 IU vitamin D per 0.45–0.6 ml. Droppers or syringes can be used to administer the supplement to the infant. National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can support mothers and families. Standardized packages are needed for training, supervision and monitoring. Dispensing needs to be documented in clinical records.

Feasibility and equity

There was no specific evidence on the feasibility and equity of providing vitamin D supplements to preterm or LBW babies.

Summary of judgements

Comparison: Vitamin D supplementation vs no vitamin D supplementation (A.10c)

Justification

- Evidence of small benefit: decreased bronchopulmonary dysplasia (*very-low-certainty evidence*), increased weight and length (*moderate-certainty evidence*) and decreased neurodevelopmental impairment (*very-low-certainty evidence*)
- Evidence on harms uncertain: mortality (*low-certainty evidence*)
- Evidence of little or no effect on infections (*moderate-certainty evidence*), hospital admissions (*very-low-certainty evidence*), head circumference (*low-certainty evidence*), weight (*low-certainty evidence*) and length (*moderate-certainty evidence*)

Evidence-to-Decision framework

Benefits	Moderate
Harms	Unknown
Certainty	Moderate
Balance	Probably favours vitamin D supplementation
Values	No uncertainty or variability about outcomes
Acceptability	Varies
Resources	Low to moderate
Feasibility	Varies
Equity	Varies

A.10d Vitamin A supplementation

Recommendation and remarks

RECOMMENDATION A.10d (UPDATED)

Enteral vitamin A supplementation may be considered for human milk-fed very preterm (< 32 weeks' gestation) or very-low-birth-weight (< 1.5 kg) infants who are not receiving vitamin A from another source. (Conditional recommendation, low-certainty evidence)

Remarks

- The recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- There were no trials in infants born ≥ 32 weeks' gestation or ≥ 1.5 kg birth weight, so the GDG did not make a recommendation for those infants.
- The GDG noted that there were limited data on the dose, timing of initiation and duration of supplementation. Based on most trials included in the evidence review, the GDG suggests a daily dose of 1000–5000 IU may be initiated when enteral feeds are well established, and may be continued until the infant receives vitamin A from another source.

Background and definitions

Vitamin A regulates cellular growth and helps to maintain the integrity of the mucosa and epithelium of the respiratory and gastrointestinal tracts (116,117). Vitamin A may also boost immune function (118,119). Preterm infants are born with low cord blood and

liver storage of vitamin A (117). Supplementation with vitamin A has been reported to reduce bronchopulmonary dysplasia in studies of very preterm infants (born before 32 weeks' gestation) (116,120,121).

Summary of the evidence

OVERVIEW	A.10d Vitamin A supplementation
PICO	Population – Preterm or LBW infants who are fed mother's own milk or donor human milk Intervention – Vitamin A supplementation Comparator – No vitamin A supplementation Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none">• Gestational age at birth (< 32 weeks, ≥ 32 weeks)• Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – Vitamin A supplementation versus no vitamin A supplementation

Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review of RCTs of "low" daily dose (< 10 000 IU/day) enteral vitamin A supplementation for preterm and/or LBW infants, which included four trials and 800 participants from three countries (China, India and the United Kingdom) (122). All infants in the included trials had gestational age below 32 weeks or birth weight below 1.5 kg and most were born before 28 weeks' gestation (extremely

preterm) or with birth weight below 1.0 kg (extremely LBW). Doses ranged from 1500 to 10 000 IU/day and initiation of supplementation was between 1 and 4 days of age in the trials. Two trials provided supplementation until 28 days after birth while the other two trials continued until 34–36 weeks PMA.

Critical outcomes

For vitamin A supplementation compared with no vitamin A supplementation, four trials reported all-cause mortality, five reported morbidity (4 reported bronchopulmonary dysplasia, 1 pneumothorax, 1 pulmonary haemorrhage, 4 retinopathy

of prematurity, 2 patent ductus arteriosus, 1 periventricular leukomalacia, 3 sepsis, 1 seizures, 3 necrotizing enterocolitis, 2 intraventricular haemorrhage) and one reported growth (weight gain). No trials reported neurodevelopment. (Full details are provided in GRADE Table A.10d, in the Web Supplement.)

- **Mortality:** Moderate-certainty evidence from four trials totalling 800 participants suggests a decrease in all-cause mortality at latest follow-up (mean 10.3 weeks) (RR 0.74, 95% CI 0.53 to 1.02).
- **Morbidity:** Low-certainty evidence from four trials totalling 746 participants suggests a decrease in bronchopulmonary dysplasia at latest follow-up (mean 11.75 weeks) (RR 0.77, 95% CI 0.50 to 1.16). Low-certainty evidence from one trial with 154 participants suggests a decrease in pneumothorax at latest follow-up (10 weeks) (RR 0.75, 95% CI 0.46 to 1.21). Low-certainty evidence from one trial with 154 participants suggests a decrease in pulmonary haemorrhage at latest follow-up (10 weeks) (RR 0.60, 95% CI 0.30 to 1.21). Low-certainty evidence from four trials totalling 742 participants suggests a decrease in retinopathy of prematurity at latest follow-up (mean 11.75 weeks) (RR 0.69, 95% CI 0.37 to 1.30). Low-certainty evidence from two trials totalling 350 participants suggests a decrease in patent ductus arteriosus at latest follow-up (mean 7 weeks) (RR 0.66, 95% CI 0.21 to 2.06). Low-certainty evidence from one trial with 262 participants suggests a decrease in periventricular leukomalacia at latest follow-up (17 weeks) (RR 0.66, 95% CI 0.38 to 1.14). Low-certainty evidence from three trials totalling 646 participants suggests little to no effect on sepsis at latest follow-up (mean 12.3 weeks) (RR 0.87, 95% CI 0.64 to 1.19). Low-certainty evidence from one trial with 154 participants suggests little to no effect on seizures at latest follow-up (10 weeks) (RR 0.82, 95% CI 0.54 to 1.25). Very-low-certainty evidence from three trials totalling 604 participants suggests little to no effect on necrotizing enterocolitis at latest follow-up (mean 12.3 weeks) (RR 1.05, 95% CI 0.71 to 1.51). Very-low-certainty evidence from two trials totalling 450 participants suggests little to no effect on intraventricular haemorrhage at latest follow-up (mean 13.5 weeks) (RR 1.00, 95% CI 0.46 to 2.17).

- **Growth:** Low-certainty evidence from one trial with 188 participants suggests little to no effect on weight gain at latest follow-up (by hospital discharge or 16 weeks) (MD 0.02, 95% CI -0.2 to 0.24).

Other outcomes

There was a decrease in length of hospital stay (mean 6.3 weeks) (MD -8.76, 95% CI -32.1 to 14.58; 2 trials, 450 participants) and an increase in serum retinol concentration (measured in µg/ml) at latest follow-up (mean 8 weeks) (MD 4.7, 95% CI 1.2 to 8.2; 1 trial, 36 participants).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials reporting on any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding what interventions are given to infants, including what and how they are fed (14). There was no specific evidence available about whether families value vitamin A supplements for their preterm or LBW baby or whether they find them acceptable.

Resources required and implementation considerations

Organization of care

The supplements can be provided in the health-care facility or at home. The family needs accurate information on the dose and how to administer the supplement. National or local guidance for health-care facilities should be used.

Infrastructure, equipment and supplies

Common methods of providing enteral vitamin A for preterm and LBW infants include infant multivitamin formulations (e.g. vitamins D, A, C, B group) in 30–50 ml bottles. Many formulations contain 1000–5000 IU vitamin A per 0.45–0.6 ml. Droppers or syringes can be used to administer the supplement to the infant. National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can support mothers and families. Standardized packages are needed for training, supervision and monitoring. Dispensing needs to be documented in clinical records.

Feasibility and equity

There was no specific evidence available about the feasibility and equity of providing vitamin A supplements to preterm or LBW babies.

Summary of judgements

Comparison: Vitamin A supplementation vs no vitamin A supplementation (A.10d)

Justification

In trials where most participants are very preterm (< 32 weeks' gestation) or VLBW (< 1.5 kg):

- Evidence of small benefit: decreased mortality (*moderate-certainty evidence*), decreased bronchopulmonary dysplasia, pneumothorax, pulmonary haemorrhage, retinopathy of prematurity, patent ductus arteriosus and periventricular leukomalacia (*low-certainty evidence*)
- No evidence of harm
- Evidence of little or no effect on sepsis, seizures, weight (*low-certainty evidence*) and on necrotizing enterocolitis and intraventricular haemorrhage (*very-low-certainty evidence*)
- No evidence on other critical outcomes

Evidence-to-Decision summary

Benefits	Small or trivial to none
Harms	Trivial or none
Certainty	Low
Balance	Probably favours vitamin A supplementation
Values	Uncertainty or variability about outcomes
Acceptability	Probably acceptable
Resources	Low to moderate
Feasibility	Feasible
Equity	Probably equitable

A.10e Calcium and phosphorous supplementation

Recommendation and remarks

NO RECOMMENDATION

Remark

- The GDG decided not to make a recommendation on calcium or phosphorous supplementation as there was little evidence of benefits or harms on any critical outcome.

Background and definitions

Preterm and LBW infants have low skeletal stores of calcium and phosphorus (123). Previous systematic reviews have reported that calcium and phosphorous supplements given to human-milk-fed preterm or LBW infants had no effect on growth (weight, length,

head circumference) but improved bone biomarkers (serum alkaline phosphatase) (123,124). No effects have been reported on mortality, morbidity or neurodevelopment and no evidence was found on the optimal dose or timing of initiation.

Summary of the evidence

OVERVIEW	A.10e Calcium and phosphorous supplementation
PICO	Population – Preterm or LBW infants who are fed mother's own milk or donor human milk Intervention – Calcium and phosphorous supplementation Comparator – No calcium and phosphorous supplementation Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none"> Gestational age at birth (< 32 weeks, ≥ 32 weeks) Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – Calcium and phosphorous supplementation versus no calcium or phosphorous supplementation

Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review of three trials (2 RCTs and 1 non-randomized trial) reporting on a total of 162 preterm and/or LBW infants from two countries (the Islamic Republic of Iran and the United Kingdom) (125). Most babies in the trials had birth weight below 1.5 kg and/or had been born before 32 weeks' gestation.

Two trials assessed the effect of phosphorus supplementation only (dose of 15 mg/kg per day in 1 trial and 25 mg/kg per day in 1 trial) and the third trial assessed the effect of supplementation with calcium and phosphorous combined (calcium 45 mg/kg per day, phosphorus 25 mg/kg per day). All three trials gave supplements enterally, via naso- or orogastric tubes. Supplementation commenced between birth and 10 days chronological age in all three trials. The duration of supplementation was between 10 and 42 days in one trial and it could not be assessed in the other two.

Critical outcomes

For calcium and phosphorous supplementation compared with no calcium or phosphorous supplementation, three trials reported morbidity (2 reported rickets, 1 osteopenia) and one trial reported growth (length and head circumference). No trials reported all-cause mortality or neurodevelopment, and no trials reported on serious adverse events. (Full details are provided in GRADE Table A.10e, in the Web Supplement.)

Morbidity: Very-low-certainty evidence from three trials totalling 159 participants suggests a decrease in osteopenia or rickets at latest follow-up (mean 38.3 weeks) (RR 0.68, 95% CI 0.46 to 0.99).

Growth: Very-low-certainty evidence from one trial with 40 participants suggests little to no effect on weight (in grams) at 6 weeks of age (MD 138.5, 95% CI -82.16 to 359.16). Very-low-certainty evidence from one trial with 40 participants suggests little to no effects on length (in centimetres) at 6 weeks of age (MD 0.77, 95% CI -0.92 to 2.46). Very-low-

certainty evidence from one trial with 40 participants suggests little to no effect on head circumference (in centimetres) at 6 weeks of age (MD 0.33, 95% CI -0.3 to 0.96).

Other outcomes

There was little or no effect on serum alkaline phosphatase (IU/L) at 6 weeks of age (MD -126.11, 95% CI -298.5 to 46.27; 2 trials, 122 participants), serum calcium (mg/dl) at 6 weeks of age (MD 0.54, 95% CI -0.19 to 1.27; 1 trial, 40 participants), or serum phosphorus (IU/L) at 6 weeks of age (MD 0.07, 95% CI -0.22 to 0.36; 1 trial, 40 participants).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials reporting on any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding what interventions are given to infants, including what and how they are fed (14). There was no specific evidence available about whether families value calcium and phosphorous supplements for their preterm or LBW baby or whether they find them acceptable.

Summary of judgements

Comparison: Calcium and phosphorous supplementation vs no calcium or phosphorous supplementation (A.10e)

Justification

- Evidence of small benefit: decreased osteopenia, rickets (*very-low-certainty evidence*)
- Evidence of little or no effect on weight, length, head circumference (*very-low-certainty evidence*)
- No evidence on other critical outcomes

Evidence-to-Decision summary

Benefits	Unknown
Harms	Unknown
Certainty	Very low
Balance	Does not favour calcium and phosphorous supplementation
Values	Uncertainty or variability about outcomes
Acceptability	Probably acceptable
Resources	Low to moderate
Feasibility	Probably feasible
Equity	Probably equitable

Resources required and implementation considerations

Organization of care

The supplements can be provided in the health-care facility or at home. The family needs accurate information on the dose and how to administer the supplement. National or local guidance for health-care facilities should be used.

Infrastructure, equipment and supplies

Common methods of providing enteral calcium and phosphorous for preterm and LBW infants include a 5 ml suspension containing 125 mg of calcium, 55 mg of phosphorus and 200 IU of vitamin D, which is given three times a day at a dose of 2 ml/kg. Droppers or syringes can be used to administer the supplement to the infant. National or local guidance for health facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can support mothers and families. Standardized packages are needed for training, supervision and monitoring. Dispensing needs to be documented in clinical records.

Feasibility and equity

There was no specific evidence available about the feasibility and equity of providing calcium and phosphorous supplements to preterm or LBW babies.

A.10f Multiple micronutrient (MMN) supplementation

Recommendation and remarks

NO RECOMMENDATION

Remark

- The GDG decided not to make a recommendation on MMN supplementation as there was no evidence of benefits or harms on any critical outcome.

Background and definitions

Many health workers advise families to give MMN supplements to human-milk-fed preterm and LBW infants (76,123). The supplements commonly include A, D, E, B group vitamins, and some contain iron,

zinc, folate and magnesium (56). However, there has been no systematic review of the effect of MMN supplements on health and developmental outcomes in preterm and LBW infants.

Summary of the evidence

OVERVIEW	A.10f MMN supplementation
PICO	Population – Preterm or LBW infants who are fed mother's own milk or donor human milk Intervention – Enteral MMN supplementation Comparator – No MMN supplementation Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none"> Gestational age at birth (< 32 weeks, ≥ 32 weeks) Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – MMN supplementation versus no MMN supplementation

Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review of the effectiveness of MMNs defined as three or more micronutrients (vitamins A, D, E, B group, iron, zinc, folate or magnesium). Two RCTs were included, which enrolled a total of 414 preterm or LBW infants from two countries (Mexico and the United Republic of Tanzania) (126). The United Republic of Tanzania trial recruited 339 preterm or LBW infants. The Mexico trial recruited 75 preterm or LBW infants. The United Republic of Tanzania trial intervention was vitamin C, E, B group, folate and vitamin B12, which was compared with no MMN in the control group. The Mexico trial gave the same nutrients to the intervention group plus zinc, magnesium, vitamin D, vitamin A and iron, and compared this with vitamin A and iron in the control group. The United Republic of Tanzania trial initiated supplementation at 66 weeks of age and continued until 18 months of age, while the Mexico trial started

supplementation at 3 months, continuing until 24 months of age.

Critical outcomes

For enteral MMN supplementation compared with no MMN supplementation, two trials reported growth outcomes (weight-for-height z score [WHZ], HAZ, WAZ) and one trial reported neurodevelopmental outcomes (cognition, receptive language, expressive language, fine motor, gross motor). No trials reported mortality or morbidity outcomes, and no trials reported on serious adverse events. (Full details are provided in GRADE Table A.10f, in the Web Supplement.)

- **Growth:** Low-certainty evidence from two trials totalling 398 participants suggests little or no effect on wasting (WHZ < -2 SD) at latest follow-up (mean 91 weeks) (RR 0.86, 95% CI 0.50 to 1.48). Low-certainty evidence from two trials totalling 399 participants suggests little or no effect on stunting (HAZ < -2 SD) at latest follow-up (mean 91 weeks) (RR 1.17, 95% CI 0.83 to 1.66). Low-certainty evidence from two trials totalling

396 participants suggests little or no effect on underweight (WAZ < -2 SD) at latest follow-up (mean 91 weeks) (RR 1.22, 95% CI 0.85 to 1.22). There was also little or no effect on change in WHZ, HAZ or WAZ scores from baseline to endline in the studies.

- **Neurodevelopment:** At latest follow-up, very-low-certainty evidence from one trial with 27 participants suggests little or no effect on: cognition scores (78 weeks) (MD 2.64, 95% CI -0.48 to 5.67); receptive language scores (78 weeks) (MD 1.19, 95% CI -0.33 to 2.71); expressive language scores (78 weeks) (MD 0.94, 95% CI -1.13 to 3.01); fine motor scores (78 weeks) (MD 1.03, 95% CI -1.13 to 3.19); and gross motor scores (78 weeks) (MD 1.14, 95% CI -0.56 to 2.84). All of these neurodevelopment outcomes were measured using BSID-III.

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials reporting on any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting nutrition, and want to take an active role in deciding what interventions are given to infants, including what and how they are fed (14). There was no specific

evidence available about whether families value MMN supplements for their preterm or LBW baby or find them acceptable.

Resources required and implementation considerations

Organization of care

The supplements can be provided in the health-care facility or at home. The family needs accurate information on the dose and how to administer the supplement. National or local guidance for health-care facilities should be used.

Infrastructure, equipment and supplies

Common methods of providing enteral MMN supplements for preterm and LBW infants include infant multivitamin formulations (e.g. vitamins D, A, C, B group with added iron) in 30–50 ml bottles. Droppers or syringes can be used to administer the supplement to the infant. National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can support mothers and families. Standardized packages are needed for training, supervision and monitoring. Dispensing needs to be documented in clinical records.

Feasibility and equity

There was no specific evidence available about the feasibility and equity of providing MMN supplements to preterm or LBW babies.

Summary of judgements

Comparison: MMN supplementation vs no MMN supplementation (A.10f)

- | | |
|----------------------|---|
| Justification | <ul style="list-style-type: none"> • Evidence of little or no effect on weight, length and neurodevelopment (<i>low- to very-low-certainty evidence</i>) • No evidence on other critical outcomes |
|----------------------|---|

Evidence-to-Decision summary

Benefits	Unknown
Harms	Unknown
Certainty	Low to very low
Balance	Does not favour MMN supplementation
Values	Uncertainty or variability about outcomes
Acceptability	Probably acceptable
Resources	Low to moderate
Feasibility	Probably feasible
Equity	Probably equitable

A.11 PROBIOTICS

Recommendation and remarks

RECOMMENDATION A.11 (NEW)

Probiotics may be considered for human-milk-fed very preterm infants (< 32 weeks' gestation).

(Conditional recommendation, moderate-certainty evidence)

Remarks

- The recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- The GDG noted that there are many infant probiotic formulations available in the public domain that have variable quality control and formulation (127,128).
- The GDG considered that only probiotics especially formulated for preterm or LBW infants that meet regulatory standards should be used, and clear instructions for safe use should be given to health workers.
- The GDG did not make a recommendation for infants born after 32 weeks' gestation because the data were insufficient.
- Only five trials (254 participants) included infants fed formula as the sole diet, so the GDG did not make a recommendation for these infants.
- The GDG was not able to make a recommendation on type (i.e. genera, species or strain), formulation (e.g. powder or drops), dose, timing or duration of probiotic administration as there was insufficient evidence. The GDG considered that type, formulation, dose, timing and duration should be based on clinical judgement.

Background and definitions

Probiotics are formulations given by the enteral route that contain bacteria (e.g. *Bifidobacterium* spp. or *Lactobacillus* spp.) or fungi (e.g. *Saccharomyces* spp.) (129,130). A range of probiotic supplements are available commercially. Probiotics colonize the mucosal surface of the human gastrointestinal tract, modulate the intestinal microbiome and promote mucosal barrier functions (129,130). Probiotics

have been used to prevent and treat infectious or inflammatory gastrointestinal conditions primarily in adults, with only low-certainty evidence of any benefit for most conditions (131-133). There have also been many trials of probiotics in preterm and LBW infants in the last 10 years showing varying effects, including reductions in sepsis and necrotizing enterocolitis (134-137), but also increases in bacteraemia and fungaemia (134,138,139).

Summary of the evidence

OVERVIEW	A.11 Probiotics
PICO	Population – Preterm or LBW infants Intervention – Any probiotics Comparator – No probiotics Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none"> • Gestational age at birth (< 32 weeks, ≥ 32 weeks) • Birth weight (< 1.5 kg, ≥ 1.5 kg) • Probiotic species (<i>Bifidobacterium</i> spp., <i>Lactobacillus</i> spp., other spp.) • Type of enteral feed (human milk, formula, mixed)

Effectiveness: Comparison – Any probiotics versus no probiotics

Sources and characteristics of the evidence

The effectiveness evidence was derived from a Cochrane systematic review of 56 trials totalling 10 812 very preterm (< 32 weeks' gestation) or VLBW infants (< 1.5 kg) (127). An updated search conducted on 1 October 2021 located no new trials.

The average birth weight was 1.0–1.2 kg and average gestation at birth was 28–32 weeks. Four trials excluded infants who were born with birth weight below the 10th percentile for the reference population (i.e. small for gestational age, or SGA). Most trials were conducted during the past 20 years (4 trials were conducted pre-2000). The trials were from 21 countries (Australia, Bangladesh, Brazil, China, France, Germany, Greece, India, the Islamic Republic of Iran, Israel, Italy, Japan, Mexico, Pakistan, Poland, Slovenia, South Africa, Thailand, Türkiye, the United Kingdom and the USA). Fifty-five trials were individually randomized and one was cluster randomized. Twenty-one trials enrolled fewer than 100 participants, 20 enrolled 100–199, 12 enrolled 200–499 participants and 3 enrolled 500 participants or more. In most trials, participating infants were given human milk or formula feeding. Seven trials enrolled infants who received human milk only and five enrolled only formula-fed participants. The probiotic preparations varied, though were mostly lyophilized (freeze dried) or liquid commercially available products supplied by the manufacturer for use in the trial. Thirty-three trials used single-genus probiotics (most commonly, *Bifidobacterium* spp. or *Lactobacillus* spp.) and 23 used multi-genus combinations (most

commonly, *Bifidobacterium* spp. plus *Lactobacillus* spp.). Most trials initiated supplementation during the first week after birth, typically with the first enteral feed. In most trials, the intervention period was at least six weeks, typically lasting until discharge from hospital. Eleven of the trials administered the intervention for a shorter period (7–30 days). One trial continued the intervention until the infant reached 2.0 kg body weight.

Critical outcomes

For probiotics compared with no probiotics, 51 trials reported all-cause mortality, 54 reported morbidity (54 reported necrotizing enterocolitis, 47 culture-confirmed infection) and 6 reported neurodevelopment (severe neurodevelopmental impairment). No trials reported growth. (Full details are provided in GRADE Table A.11, in the Web Supplement.)

- **Mortality:** Moderate-certainty evidence from 51 trials totalling 10 170 participants suggests a decrease in all-cause mortality by hospital discharge (RR 0.76, 95% CI 0.65 to 0.89).
- **Morbidity:** Low-certainty evidence from 54 trials totalling 10 604 participants suggests a decrease in necrotizing enterocolitis by hospital discharge (RR 0.54, 95% CI 0.45 to 0.65). Moderate-certainty evidence from 47 trials totalling 9762 participants suggests a decrease in invasive infection by hospital discharge (RR 0.89, 95% CI 0.82 to 0.97).
- **Neurodevelopment:** Low-certainty evidence from five trials totalling 1518 participants suggests little or no effect on neurodevelopment (severe neurodevelopmental impairment assessed using a validated test) between 18 months and 3 years (RR 1.03, 95% CI 0.84 to 1.26).

Other outcomes

There was a decrease in length of hospital stay (in days) (MD -1.93, 95% CI -3.78 to -0.08; 22 trials, 5458 infants).

Subgroup analyses

Subgroup differences for growth and neurodevelopment could not be assessed as there were insufficient studies. No difference was found for mortality, necrotizing enterocolitis or sepsis for any of the subgroups: gestational age and birth weight, probiotic species or type of enteral feed.

Other studies

Eight studies (3080 participants) recruited infants with gestational age 32–36 weeks (mean 33 weeks (SD 4 weeks) and showed decreases in all-cause mortality (RR 0.50, 95% CI 0.43 to 1.17; 4 trials, 2334 participants, low-certainty evidence), necrotizing enterocolitis (RR 0.32, 95% CI 0.16 to 0.66; 6 trials, 1493 participants, low-certainty evidence), sepsis (RR 0.50, 95% CI 0.29 to 0.85; 6 trials, 2708 participants, low-certainty evidence) and neurodevelopmental impairment (RR 0.48, 95% CI 0.29 to 0.80; 1 trial, 249 participants, very-low-certainty evidence).

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, and want to take an active role in deciding what interventions are given to infants, including what and how they are fed (14). A study from the United Kingdom reported that families are willing to consider use of probiotics for

their preterm or LBW infants if there is evidence of benefit and safety (140). There was no other specific evidence available about whether families value probiotic supplements for their preterm or LBW baby or find them acceptable.

Resources required and implementation considerations

Organization of care

Probiotics can be provided in the health-care facility or at home. The family needs accurate information on the dose and how to administer the supplement. National or local guidance for health-care facilities should be used.

Infrastructure, equipment and supplies

Common probiotic preparations are either single-genus or multi-genus probiotic combinations (including *Bifidobacterium* spp. plus *Lactobacillus* spp). Dosing, amounts, frequency and duration vary. Probiotics can be provided as powder or liquids in bottles or mixed with infant milk or sterile water. National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can support mothers and families. Standardized packages are needed for training, supervision and monitoring. Dispensing needs to be documented in clinical records.

Feasibility and equity

There was no specific evidence available about the feasibility and equity of providing probiotics to preterm or LBW babies.

Summary of judgements

Comparison: Any probiotics vs no probiotics (A.11)

Justification	<p>In trials where most participants are very preterm (< 32 weeks' gestation) or VLBW (< 1.5 kg):</p> <ul style="list-style-type: none">• Evidence of moderate benefit: decreased mortality, necrotizing enterocolitis and invasive infection (<i>moderate-certainty evidence</i>)• No evidence of harms• Evidence of little or no effect on neurodevelopment (<i>low-certainty evidence</i>)• No evidence on other critical outcomes
---------------	--

Evidence-to-Decision summary

Benefits	Moderate
Harms	Trivial or none
Certainty	Moderate
Balance	Probably favours probiotics
Values	No uncertainty or variability about outcomes
Acceptability	Varies
Resources	Low to moderate
Feasibility	Varies
Equity	Varies

A.12 EMOLLIENTS

Recommendation and remarks

RECOMMENDATION A.12 (NEW)

Application of topical oils to the body of preterm or low-birth-weight infants may be considered.

(Conditional recommendation, moderate-certainty evidence)

Remarks

- The recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- The GDG noted that there were limited data on the type, dose, timing of initiation and duration of oil use. Based on most of the trials included in the evidence review, the GDG suggested that sunflower or coconut oils may be used and that initiation and duration of use may be based on clinical judgement. The GDG also felt that application of oils should be done gently to avoid disrupting skin integrity.
- The GDG decided not to make a recommendation on the use of ointments or creams due to little or no effect on mortality and morbidity (invasive infection, necrotizing enterocolitis, bronchopulmonary dysplasia, retinopathy of prematurity) and no evidence on other critical outcomes.

Background and definitions

Emollients are moisturizing treatments applied topically, i.e. directly to the skin. They include ointments (water-in-oil suspensions), creams (oil-in-water suspensions) and natural vegetable or plant topical oils (e.g. sunflower and coconut oils). The skin of preterm infants is developmentally immature (141,142) and can be easily abraded, which can allow entry of pathogenic organisms (143). Topical

emollients can improve skin integrity and barrier (protective) functions but they can also disrupt skin integrity, remove normal flora and microorganisms and increase colonization with other microorganisms (142). Emollients also contain fatty acids and other fluids that can be absorbed through the skin (141). However, there have been no recent systematic reviews of the effectiveness of topical ointments, creams or oils in preterm and LBW infants.

Summary of the evidence

OVERVIEW	A.12a Topical oil	A.12b Topical ointment or cream
PICO	Population – Preterm and LBW infants Intervention 1 – Topical oil Comparator 1 – No topical oil Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up	Population – Preterm and LBW infants Intervention 2 – Topical ointment or cream Comparator 2 – No topical ointment or cream Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none"> • Gestational age at birth (< 32 weeks, ≥ 32 weeks) • Birth weight (< 1.5 kg, ≥ 1.5 kg) 	

Effectiveness: Comparison 1 – Topical oil versus no topical oil

Sources and characteristics of the evidence

For this comparison, the effectiveness evidence was derived from a systematic review of 15 RCTs enrolling a total of 3718 infants (144) from nine countries (Bangladesh, Brazil, Egypt, France, Germany, India, the Islamic Republic of Iran and Pakistan). All trials used natural vegetable or plant oils: sunflower (8

trials), coconut (4 trials), and soybean, almond, vegetable and olive oil (1 trial each). The population was very preterm babies (< 32 weeks' gestation) in three trials. The intervention generally commenced within a few days of birth and continued until about 1–4 weeks chronological age or until hospital discharge. The oils were applied 2–6 times each day onto the whole skin surface (except the face and head) by the family or health worker.

Critical outcomes

For topical oil compared with no topical oil, 11 trials reported all-cause mortality, 9 reported morbidity (9 reported invasive infection, 1 necrotizing enterocolitis, 1 bronchopulmonary dysplasia, 1 retinopathy of prematurity), 7 reported growth (7 weight gain, 6 length, 6 head circumference) and 1 reported neurodevelopment (cognitive, language, motor and socioemotional outcomes [BSID-III]). No trials reported on serious adverse events. (Full details are provided in GRADE Table A.12a, in the Web Supplement.)

- **Mortality:** Low-certainty evidence from 11 trials totalling 1119 participants suggests little to no effect on all-cause mortality by hospital discharge (RR 0.94, 95% CI 0.82 to 1.08).
- **Morbidity:** Low-certainty evidence from nine trials totalling 3256 participants suggests a decrease in invasive infection by hospital discharge (RR 0.71, 95% CI 0.52 to 0.96). Very-low certainty evidence from one trial with 72 participants suggests a decrease in necrotizing enterocolitis by hospital discharge (RR 0.20, 95% CI 0.01 to 4.03). Very-low-certainty evidence from one trial with 72 participants suggests little to no effect on bronchopulmonary dysplasia at 26 weeks PMA (RR 0.93, 95% CI 0.53 to 1.64). Very-low-certainty evidence from one trial with 72 participants suggests little to no effect on retinopathy of prematurity by hospital discharge (RR 1.00, 95% CI 0.27 to 3.69).
- **Growth:** Low-certainty evidence from seven trials totalling 433 participants suggests an increase in the rate of weight gain (in grams per kilogram per day) by hospital discharge (MD 2.93, 95% CI 2.11 to 3.76). Moderate-certainty evidence from six trials totalling 358 participants suggests an increase in crown-heel length (millimetres

per week) by hospital discharge (MD 1.34, 95% CI 0.2 to 2.74). Low-certainty evidence from six trials totalling 358 participants suggests little to no effect on change in head circumference (in millimetres per week) by hospital discharge (MD 0.66, 95% CI 0.54 to 1.85).

- **Neurodevelopment:** Very-low-certainty evidence from one trial with 51 participants suggests little to no effect on cognitive developmental delay at 24 months of age (RR 0.25, 95% CI 0.06 to 1.11). Very-low-certainty evidence from one trial with 51 participants suggests little to no effect on language developmental delay at 24 months of age (RR 0.48, 95% CI 0.21 to 1.11). Very-low-certainty evidence from one trial with 51 participants suggests little to no effect on motor developmental delay at 24 months of age (RR 0.25, 95% CI 0.06 to 1.11). Very-low-certainty evidence from one trial with 51 participants suggests little to no effect on socio-emotional developmental delay at 24 months of age (RR 0.30, 95% CI 0.07 to 1.33). All neurodevelopmental outcomes were measured using BSID-III.

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Other studies

One additional trial also reported a decrease in infection-specific mortality by 28 days of age (adjusted odds ratio 0.72, 95% CI 0.39 to 1.34; 1 trial, 103 participants) and a decrease in nosocomial infections by 28 days of age (adjusted incidence ratio 0.46, 95% CI 0.26 to 0.81; 1 trial, 103 participants) (145).

Effectiveness: Comparison 2 – Topical ointment or cream versus no topical ointment or cream

Sources and characteristics of the evidence

For the second comparison, the effectiveness evidence was derived from a systematic review of eight RCTs including 2086 preterm or LBW infants from five countries (144) (Austria, Bangladesh, Saudi Arabia, Türkiye and the USA). Most trials enrolled very preterm babies born at gestational ages up to 30 weeks while others enrolled babies born before 31 weeks (1 study), before 33 weeks (3 studies), before 34 weeks (1 study) or up to 36 weeks' gestation (2 studies). The trials used commercially available ointments or creams. The intervention generally commenced within a few days after birth and continued until about 1–4 weeks postnatal age or until hospital discharge. The ointments or creams were applied 2–6 times each day onto the whole skin surface (except the face and head) by the family or health worker.

Critical outcomes

For topical ointment or cream compared with no topical ointment or cream, seven trials reported all-cause mortality and eight reported morbidity (8 reported invasive infection, 4 necrotizing enterocolitis, 2 bronchopulmonary dysplasia, 1 retinopathy of prematurity). Growth and neurodevelopment outcomes were not reported. (Full details are provided in GRADE Table A.12b, in the Web Supplement.)

- **Mortality:** Low-certainty evidence from seven trials totalling 2067 participants suggests little or no effect on all-cause mortality by hospital discharge (RR 0.87, 95% CI 0.75 to 1.03).
- **Morbidity:** Low-certainty evidence from eight trials totalling 2086 participants suggests little or no effect on invasive infection (at least one infection with any organism) by hospital discharge (RR 1.13, 95% CI 0.97 to 1.31). Low-certainty evidence from four trials totalling 1472 participants suggests little or no effect on necrotizing enterocolitis by hospital discharge (RR 1.25, 95% CI 0.89 to 1.76). Low-certainty evidence from two trials totalling 1009 participants suggests little or no effect on bronchopulmonary dysplasia by hospital discharge (RR 1.00, 95% CI 0.88 to 1.14).

Very-low-certainty evidence from one trial with 952 participants suggests little or no effect on retinopathy of prematurity by hospital discharge (RR 0.99, 95% CI 0.77 to 1.28).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, and want to take an active role in deciding what interventions are given to infants, including what and how they receive skin care (14). There was no specific evidence available about whether families value emollients for their preterm or LBW baby or find them more or less acceptable.

Resources required and implementation considerations

Organization of care

Emollients can be provided in the health-care facility or at home. They can be spread gently over the infant's abdomen, back and limbs. The family needs accurate information on how to apply the emollients gently. National or local guidance for health-care facilities should be used.

Infrastructure, equipment and supplies

Emollient preparations include sunflower and coconut oils. National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can support mothers and families. Standardized packages are needed for training, supervision and monitoring. Dispensing needs to be documented in clinical records.

Feasibility and equity

There was no specific evidence available about the feasibility and equity of topical emollient application for preterm or LBW babies.

Summary of judgements

	Comparison 1. Topical oil vs no topical oil (A.12a)	Comparison 2. Topical ointment or cream vs no topical ointment or cream (A.12b)
Justification	<ul style="list-style-type: none"> ▪ Evidence of moderate benefits: decreased severe infection (<i>low-certainty evidence</i>), increased weight (<i>low-certainty evidence</i>) and increased length (<i>moderate-certainty evidence</i>) ▪ No evidence of harms ▪ Evidence of little or no effect on: mortality (<i>low-certainty evidence</i>), necrotizing enterocolitis, bronchopulmonary dysplasia, retinopathy of prematurity (<i>low-certainty evidence</i>), head circumference (<i>low-certainty evidence</i>) and neurodevelopment (<i>very-low-certainty evidence</i>) ▪ No evidence on other critical outcomes 	<ul style="list-style-type: none"> ▪ Evidence of little or no effect on all-cause mortality, invasive infection, necrotizing enterocolitis, bronchopulmonary dysplasia and retinopathy of prematurity (<i>low-certainty evidence</i>) ▪ No evidence on other critical outcomes

Evidence-to-Decision summary		
Benefits	Moderate	Trivial or none
Harms	Trivial or none	Trivial or none
Certainty	Low	Low
Balance	Probably favours topical oils	Does not favour ointments or creams
Values	No uncertainty or variability about outcomes	No uncertainty or variability about outcomes
Acceptability	Probably yes	Probably yes
Resources	Low to moderate	Low to moderate
Feasibility	Probably yes	Varies
Equity	Probably equitable	Probably equitable

B. Care for complications

B.1 CONTINUOUS POSITIVE AIRWAY PRESSURE FOR RESPIRATORY DISTRESS SYNDROME

Recommendation and remarks

RECOMMENDATION B.1 (UPDATED)

Continuous positive airway pressure (CPAP) therapy is recommended in preterm infants with clinical signs of respiratory distress syndrome. (*Strong recommendation, moderate-certainty evidence*)

Remarks

- The GDG noted that the evidence on harms (increased pneumothorax) was of uncertain clinical significance and the overall certainty of the body of evidence was low due to imprecision and indirectness.
- The GDG noted that there were limited data on the timing of initiation and duration of CPAP. Based on most of the trials included in the evidence review, the GDG suggests that CPAP may be considered as soon as the diagnosis of respiratory distress syndrome (RDS) is clinically suspected, and that duration should be based on clinical judgement.
- The GDG also noted that CPAP implementation must be done with skilled staff, quality equipment and quality consumables (including humidified blended oxygen-air and monitors).
- The GDG decided not to make a separate recommendation on the timing of CPAP for infants with RDS.

Background and definitions

Respiratory distress syndrome (RDS) is a major cause of morbidity and mortality in preterm infants (146). RDS commonly develops in the first hours after birth and develops or “becomes established” over the first few days after birth (146-148). Until the 1970s, initial therapy for RDS was traditionally oxygen given through a head box or nasal prongs, and infants with severe disease received mechanical ventilation.

Continuous positive airway pressure (CPAP) involves connecting a nasal “interface” (prongs, face mask or head box) via tubing to a pressure source with an air-oxygen mix (149,150). CPAP provides distending pressure into the upper and lower airways preventing collapse, especially during expiration. CPAP devices were adapted for use in preterm babies in the 1970s and CPAP is now routinely used for preterm babies with RDS in many health-care facilities globally.

Summary of the evidence

OVERVIEW	B.1a Any CPAP	B.1b Early CPAP
PICO	Population – Preterm infants with RDS Intervention 1 – Any CPAP Comparator 1 – Usual supplemental oxygen therapy by head box, face mask or nasal cannula Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up	Population – Preterm infants with RDS Intervention 2 – Early CPAP Comparator 2 – Delayed CPAP Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – From birth Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none"> • Gestational age at birth (< 32 weeks, ≥ 32 weeks) • Birth weight (< 1.5 kg, ≥ 1.5 kg) 	

Effectiveness: Comparison 1 – Any CPAP versus supplemental oxygen

Sources and characteristics of the evidence

The effectiveness evidence for this comparison was derived from a Cochrane systematic review of five RCTs conducted in the 1970s and 1980s reporting on a total of 322 preterm infants (151). An updated search conducted on 1 October 2021 located no new trials.

Four studies were conducted in high-income settings (Australia, the United Kingdom and the USA) and one in a low-resource setting (the United Republic of Tanzania). Infants were included if they had RDS (defined as an infant needing FiO_2 [fraction of inspired oxygen] > 0.30). All trials used traditional CPAP as the intervention, none used “bubble” or newer types of CPAP. The comparator in all the trials was supplemental oxygen. No infants received mechanical ventilation in the control group. The mean age at study entry ranged from 10 to 150 hours post-birth. The mean birth weight of infants was 1.7–2.0 kg, with two trials excluding infants weighing less than 1.0 kg at birth.

Critical outcomes

For any CPAP compared with supplemental oxygen for RDS, five trials reported all-cause mortality outcomes, five trials reported morbidity (3 reported use of mechanical ventilation, 5 “failed treatment”, 4 pneumothorax, 2 bronchopulmonary dysplasia). No trials reported growth or neurodevelopment outcomes. (Full details are provided in GRADE Table B.1a, in the Web Supplement.)

- **Mortality:** Moderate-certainty evidence from five trials totalling 322 participants suggests a decrease in all-cause mortality by hospital discharge (RR 0.53, 95% CI 0.34 to 0.83).
- **Morbidity:** Very-low-certainty evidence from three trials totalling 233 participants suggests a decrease in the use of mechanical ventilation by hospital discharge (RR 0.72, 95% CI 0.54 to 0.96). Very-low-certainty evidence from five trials totalling 322 participants suggests a decrease in “failed treatment” (a composite outcome of death or the use of mechanical ventilation) by hospital discharge (RR 0.64, 95% CI 0.50 to 0.82). Low-

certainty evidence from four trials totalling 270 participants suggests an increase in pneumothorax by hospital discharge (RR 2.48, 95% CI 1.16 to 5.30). Very-low-certainty evidence from two trials totalling 209 participants suggests little or no effect on bronchopulmonary dysplasia (defined as oxygen dependency at 28 days) by 36 weeks PMA (RR 1.04, 95% CI 0.35 to 3.13).

Other outcomes

One trial reported a decrease in the composite outcome of death or abnormal blood gases by hospital discharge (RR 0.53, 95% CI 0.32 to 0.90; 1 trial, 24 infants). One trial reported a decrease in the outcome of “transfer to an NICU” by hospital discharge (RR 0.49, 95% CI 0.30 to 0.78; 1 trial, 24 infants). One trial reported a decrease in the duration of oxygen therapy by hospital discharge (MD 0.20 days, 95% CI -2.47 to 2.87; 1 trial, 24 infants).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Effectiveness: Comparison 2 – Early versus delayed CPAP

Sources and characteristics of the evidence

The effectiveness evidence for this comparison was derived from a systematic review of four trials (2 RCTs and 2 quasi-RCTs) that recruited 119 preterm infants with RDS (mean birth weight 1.5–2.1 kg, mean gestational age 31–34 weeks) conducted in the United Kingdom and the USA in the 1970s or the early 1980s (152). An updated search conducted on 1 October 2021 located no new trials. Infants were eligible for inclusion if they were given a diagnosis of RDS (based on clinical and radiological criteria) and were breathing spontaneously. Infants were randomized to receive CPAP immediately as soon as the diagnosis of RDS was made (“early group”) or for treatment to be delayed until deterioration as defined by the study (“delayed group”). The early CPAP group received CPAP at a mean age of 7–18 hours post-birth and required FiO_2 0.30 to 0.60. The delayed CPAP group required FiO_2 from 0.60 to 1.0 but the mean age of receipt of CPAP was not stated in any trial.

Critical outcomes

For early compared with delayed CPAP for RDS, four trials reported all-cause mortality, four trials reported morbidity (4 reported the use of mechanical ventilation, 3 pneumothorax, 1 bronchopulmonary dysplasia). No trials reported growth or neurodevelopment outcomes. (Full details are provided in GRADE Table B.1b, in the Web Supplement.)

- **Mortality:** Low-certainty evidence from four trials totalling 119 participants suggests little or no effect on all-cause mortality by hospital discharge (RR 0.93, 95% CI 0.43 to 2.03).
- **Morbidity:** Very-low-certainty evidence from four trials totalling 119 participants suggests a decrease in the use of mechanical ventilation by hospital discharge ((RR 0.77, 95% CI 0.43 to 1.38). Low-certainty evidence from two trials totalling 98 participants suggests little or no effect on pneumothorax (RR 1.09, 95% CI 0.39 to 3.04). Very-low-certainty evidence from one trial with 29 participants suggests an increase in bronchopulmonary dysplasia at 36 weeks PMA (RR 1.42, 95% CI 0.10 to 20.49).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Other studies

Two studies assessing the effect of continuous negative pressure (153,154) were included in the previous Cochrane review (155) but not in the updated Cochrane review (152), due to a change in the PICO intervention from negative pressure to continuous positive airway pressure (CPAP). The Cochrane review also excluded two RCTs (156,157) because they provided very early CPAP at 5 minutes of age, which was considered to be earlier than RDS could be established in the babies. These two RCTs were included in the 2021 Cochrane review of prophylactic and very early CPAP by Subramaniam et al. (see Recommendation B.1) (158).

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant

(see Table 1.1) reported that carers want assistance in interacting with their babies, especially when they are undergoing therapies that make it difficult to have physical contact (14). They want to learn about the health-care setting where they need to stay and care for their baby. They want to understand what medical equipment is being used and why. Studies report that families can find mechanical ventilation and CPAP intimidating and frightening and that these therapies can accentuate their feelings of inadequacy and lack of control over their baby's health care (147,159). Families also worry about the pain and discomfort their baby is experiencing in NICUs (14). No other specific evidence was located about whether families value CPAP rather than supplemental oxygen for their preterm or LBW baby or whether they find CPAP more or less acceptable than other supplemental oxygen.

Resources required and implementation considerations

Organization of care

CPAP for preterm or LBW infants should be done in special or intensive care units (level 2 or 3 facilities).

Infrastructure, equipment and supplies

CPAP devices include a pressure source with an air-oxygen mix. CPAP devices include "bubble" (underwater, water-seal) CPAP, ventilator CPAP and "Infant Flow Driver" CPAP. CPAP also requires an "interface", which is commonly a mask or nasal prongs. Disposable tubes and suction catheters are also needed. National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers who are qualified to work in level 2 (special newborn care units, special care nurseries) and level 3 (intensive care) facilities can support the provision of CPAP. Standardized packages are needed for training, supervision and monitoring.

Feasibility and equity

There was no specific evidence on the feasibility and equity of providing CPAP for preterm or LBW infants.

Summary of judgements

	Comparison 1. Any CPAP vs supplemental oxygen (B1.a)	Comparison 2. Early vs delayed CPAP (B1.b)
Justification	<ul style="list-style-type: none"> Evidence of moderate benefits: decreased mortality (<i>moderate-certainty evidence</i>), decreased mechanical ventilation (<i>very-low-certainty evidence</i>) and decreased “failed treatment”, i.e. death or use of mechanical ventilation (<i>very-low-certainty evidence</i>) Evidence of small harms: increased pneumothorax (<i>low-certainty evidence</i>) Evidence of little or no effect on bronchopulmonary dysplasia (<i>very-low-certainty evidence</i>) 	<ul style="list-style-type: none"> Evidence of small benefits: decrease in use of mechanical ventilation (<i>very-low-certainty</i>) Evidence of small harm: increase in bronchopulmonary dysplasia (<i>very-low-certainty evidence</i>) Evidence of little or no effect on mortality and air leak (pneumothorax) (<i>low-certainty evidence</i>) No evidence on other critical outcomes

Evidence-to-Decision summary		
Benefits	Moderate	Unknown
Harms	Small	Unknown
Certainty	Low	Very low
Balance	Favours CPAP	Unknown
Values	No uncertainty or variability about outcomes	Unknown
Acceptability	Probably yes	Unknown
Resources	Large	Negligible
Feasibility	Varies	Probably feasible
Equity	Varies	Probably equitable

B.2 CONTINUOUS POSITIVE AIRWAY PRESSURE IMMEDIATELY AFTER BIRTH

Recommendation and remarks

RECOMMENDATION B.2 (NEW)

Continuous positive airway pressure (CPAP) therapy may be considered immediately after birth for very preterm infants (< 32 weeks' gestation), with or without respiratory distress. (*Conditional recommendation, low-certainty evidence*)

Remarks

- The recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- The GDG noted that duration of CPAP (i.e. when to stop CPAP) should be based on clinical judgement.
- The GDG also noted that skilled staff and quality equipment (including humidified blended oxygen-air) are needed for the administration of CPAP to preterm and LBW infants.

Background and definitions

The benefits of CPAP for RDS in preterm infants are well established (151,155). However, it can be difficult to ascertain respiratory status in preterm babies soon after birth and to accurately predict the prognosis. Preterm babies with respiratory failure may not show signs of respiratory distress in the first hours after birth and babies with early respiratory distress may improve (147). Thus, health workers in NICUs

sometimes administer CPAP immediately after birth to all babies at risk, regardless of respiratory status (sometimes called immediate CPAP), rather than assessing for RDS. Benefits and harms of this practice have been unclear (147,148,150). However, recent trials have assessed the effectiveness of immediate CPAP compared with both supplemental oxygen and mechanical ventilation.

Summary of the evidence

OVERVIEW	B.2a Immediate CPAP vs supplemental oxygen	B.2b Immediate CPAP vs mechanical ventilation
PICO	<p>Population – Preterm infants immediately after birth</p> <p>Intervention 1 – CPAP commencing immediately after birth</p> <p>Comparator 1 – Supplemental oxygen by head box, face mask or nasal cannula</p> <p>Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up</p>	<p>Population – Preterm infants immediately after birth</p> <p>Intervention 2 – CPAP commencing immediately after birth</p> <p>Comparator 2 – Mechanical ventilation</p> <p>Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up</p>
Timing, setting, subgroups	<p>Timing of the intervention – Immediately after birth</p> <p>Setting – Health-care facility or home in any country or setting</p> <p>Subgroups</p> <ul style="list-style-type: none"> • Gestational age at birth (< 32 weeks, ≥ 32 weeks) • and birth weight (< 1.5 kg, ≥ 1.5 kg) 	

Effectiveness: Comparison 1 – Immediate CPAP versus supplemental oxygen

Sources and characteristics of the evidence

For this comparison, the effectiveness evidence was derived from a Cochrane systematic review of four trials enrolling a total of 765 infants under 32 weeks' gestation at birth from seven countries (Argentina,

Brazil, Canada, Italy, Paraguay, Peru and Uruguay) (158). An updated search conducted on 1 October 2021 located no new trials. The review found two types of trial: (i) trials that provided CPAP within 15 minutes of birth regardless of respiratory status, and (ii) trials that provided CPAP between 15 and 60 minutes after birth prior to the onset of RDS.

Critical outcomes

For comparison 1, four trials reported all-cause mortality, four reported morbidity (4 reported “failed treatment”, 4 bronchopulmonary dysplasia, 1 a composite outcome of death or bronchopulmonary dysplasia, 3 pneumothorax, 2 intraventricular haemorrhage). No trials reported growth or neurodevelopment. (Full details are provided in GRADE Table B.2a, in the Web Supplement.)

- **Mortality:** Moderate-certainty evidence from four trials totalling 765 participants suggests little or no effect on all-cause mortality by hospital discharge (RR 1.09, 95% CI 0.60 to 1.96).
- **Morbidity:** Very-low-certainty evidence from four trials totalling 765 participants suggests a decreased risk of “failed treatment” (defined as recurrent apnoea, hypoxia, hypercarbia, increasing oxygen requirement, or the need for mechanical ventilation) by hospital discharge (RR 0.60, 95% CI 0.49 to 0.74). Moderate-certainty evidence from three trials totalling 683 participants suggests decreased bronchopulmonary dysplasia by 36 weeks PMA (RR 0.76, 95% CI 0.51 to 1.14). Low-certainty evidence from one trial with 256 participants suggests decreased death or bronchopulmonary dysplasia by 36 weeks PMA (RR 0.69, 95% CI 0.40 to 1.19). Low-certainty evidence from three trials totalling 568 participants suggests decreased pneumothorax by hospital discharge (RR 0.75, 95% CI 0.35 to 1.61). Low-certainty evidence from two trials totalling 486 participants suggests little or no effect on intraventricular haemorrhage grade 3 or 4 by hospital discharge (RR 0.96, 95% CI 0.39 to 2.37).

Other outcomes

Three trials reported a decrease in surfactant use by hospital discharge (RR 0.75, 95% CI 0.58 to 0.96; 3 trials, 683 participants).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Effectiveness: Comparison 2 – Immediate CPAP versus mechanical ventilation

Sources and characteristics of the evidence

For this comparison, the effectiveness evidence was derived from the same Cochrane systematic review (158). Three trials were included, which enrolled a total of 2364 very preterm infants (< 32 weeks’ gestation) from 17 countries (Argentina, Australia,

Belgium, Brazil, Canada, Chile, France, Germany, Greece, Italy, the Islamic Republic of Iran, New Zealand, Norway, Paraguay, Peru, Uruguay and the USA). An updated search conducted on 1 October 2021 located no new trials. The review included the same two types of trials as described above.

Critical outcomes

For comparison 2, four trials reported all-cause mortality, four trials reported morbidity (4 reported “failed treatment”, 3 bronchopulmonary dysplasia, 1 a composite outcome of death or bronchopulmonary dysplasia, 3 pneumothorax, 2 intraventricular haemorrhage) and one trial reported on neurodevelopment (neurodevelopmental impairment). No trials reported growth. (Full details are provided in GRADE Table B.2b, in the Web Supplement.)

- **Mortality:** Moderate-certainty evidence from three trials totalling 2358 participants suggests little or no effect on all-cause mortality by hospital discharge (RR 0.82, 95% CI 0.66 to 1.03).
- **Morbidity:** Moderate-certainty evidence from two trials totalling 1042 participants suggests a decrease in risk of “failed treatment” (defined as recurrent apnoea, hypoxia, hypercarbia, increasing oxygen requirement, or the need for mechanical ventilation) by hospital discharge (RR 0.49, 95% CI 0.45 to 0.54). Moderate-certainty evidence from three trials totalling 2150 participants suggests a decrease in bronchopulmonary dysplasia at 36 weeks PMA (RR 0.89, 95% CI 0.80 to 0.99). Moderate-certainty evidence from three trials totalling 2358 participants suggests a decrease in the combined outcome of all-cause mortality and bronchopulmonary dysplasia at 36 weeks PMA (RR 0.89, 95% CI 0.81 to 0.97). Low-certainty evidence from three trials totalling 2357 participants suggests little or no effect on pneumothorax by hospital discharge (RR 1.24, 95% CI 0.91 to 1.69). Moderate-certainty evidence from three trials totalling 2301 participants suggests little or no effect on intraventricular haemorrhage grade 3 or 4 by hospital discharge (RR 1.09, 95% CI 0.86 to 1.39).
- **Neurodevelopment:** Moderate-certainty evidence from one trial with 976 participants suggests little or no effect on neurodevelopmental impairment (defined as cerebral palsy, developmental delay, intellectual impairment, blindness or sensorineural deafness) by 18–22 months of age (RR 0.91, 95% CI 0.62 to 1.32).

Other outcomes

There was decrease in surfactant use (RR 0.60, 95% CI 0.57 to 0.63; 3 trials, 2354 infants).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that carers want assistance in interacting with their babies, especially when they are undergoing therapies that make it difficult to have physical contact (14). They want to learn about the health-care setting where they need to stay and care for their baby. They want to understand what medical equipment is being used and why. Studies report that families can find mechanical ventilation and CPAP

intimidating and frightening and that these therapies can accentuate their feelings of inadequacy and lack of control over their baby's health care (147,159). Families also worry about the pain and discomfort their baby is experiencing in NICUs (14). No other specific evidence was located about whether families value immediate CPAP for their preterm or LBW baby or whether they find it more or less acceptable than supplemental oxygen.

Resources required and implementation considerations

Please refer to the information on this topic in section B.1.

Feasibility and equity

As described in section B.1, there was no specific evidence on the feasibility and equity of providing CPAP for preterm or LBW infants.

Summary of judgements

	Comparison 1. CPAP immediately after birth for very preterm infants vs supplemental oxygen (B2.a)	Comparison 2. CPAP immediately after birth for very preterm infants vs mechanical ventilation (B2.a)
Justification	<p>In trials where most participants are very preterm (< 32 weeks' gestation):</p> <ul style="list-style-type: none"> Evidence of small benefits: decreased "failed treatment" (defined as recurrent apnoea, hypoxia, hypercarbia, increasing oxygen requirement or the need for mechanical ventilation), decreased bronchopulmonary dysplasia (<i>moderate-certainty evidence</i>) and decreased pneumothorax (<i>low-certainty evidence</i>) No evidence of harms Evidence of little or no effect on mortality and intraventricular haemorrhage (<i>moderate-certainty evidence</i>) No evidence on other critical outcomes 	<p>In trials where most participants are very preterm (< 32 weeks' gestation):</p> <ul style="list-style-type: none"> Evidence of moderate benefits: decreased "failed treatment" (defined as recurrent apnoea, hypoxia, hypercarbia, increasing oxygen requirement or the need for mechanical ventilation), decreased bronchopulmonary dysplasia (<i>moderate-certainty evidence</i>) No evidence of harms Evidence of little or no effect on mortality (<i>moderate-certainty evidence</i>) pneumothorax (<i>low-certainty evidence</i>), intraventricular haemorrhage (<i>moderate-certainty evidence</i>) and neurodevelopment (<i>moderate-certainty evidence</i>) No evidence on other critical outcomes
Evidence-to-Decision summary		
Benefits	Small	Moderate
Harms	Trivial or none	Trivial or none
Certainty	Low	Moderate
Balance	Probably favours CPAP immediately after birth for very preterm infants (< 32 weeks' gestation)	Probably favours CPAP immediately after birth for very preterm infants (< 32 weeks' gestation)
Values	No uncertainty or variability about outcomes	Probable uncertainty or variability about outcomes
Acceptability	Varies	Varies
Resources	Vary	Vary
Feasibility	Varies	Varies
Equity	Varies	Varies

B.3 CONTINUOUS POSITIVE AIRWAY PRESSURE SOURCE

Recommendation and remarks

RECOMMENDATION B.3 (NEW)

For preterm infants who need continuous positive airway pressure (CPAP) therapy, bubble CPAP may be considered rather than other pressure sources (e.g. ventilator CPAP). (*Conditional recommendation, low-certainty evidence*)

Remarks

- The recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- Evidence was derived from trials that compared underwater (water-seal) “bubble” CPAP with mechanical ventilator CPAP or Infant Flow Driver (IFD) CPAP. All trials used commercially available devices and all used humidified blended oxygen-air.
- The GDG noted that the evidence on harms (increased nasal injury) was of uncertain clinical significance and the certainty of the body of evidence was low due to bias and imprecision.
- The GDG suggested that the nasal interfaces (i.e. prongs and cannulas) used with bubble CPAP machines should be carefully selected and that skilled nursing care is needed for the prongs and cannulas.
- The GDG also considered that careful selection, maintenance and monitoring of bubble CPAP devices is needed. Only commercially available bubble CPAP devices should be used; locally-manufactured or locally-adapted bubble CPAP devices should not be used.

Background and definitions

There are many different types of CPAP machines and pressure generation for ventilatory support of preterm infants. There is also considerable variation in practice and differing reports of benefits and harms (150,160,161). The older-style CPAP pressure sources

were mechanical ventilators; newer types include Infant Flow Driver (IFD) and bubble CPAP. Bubble CPAP uses an underwater water-seal method and is commonly used for providing CPAP to babies in LMICs (150,160,161).

Summary of the evidence

OVERVIEW	B.3 Continuous positive airway pressure source
PICO	Population – Preterm infants with respiratory distress syndrome or post-extubation Intervention – Bubble CPAP pressure source Comparator – Other CPAP pressure sources (ventilator CPAP or Infant Flow Driver CPAP) Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Immediately after birth Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none">• Gestational age at birth (< 32 weeks, ≥ 32 weeks)• Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – Bubble CPAP versus other CPAP pressure sources

Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review of 15 RCTs including a total of 1437 preterm infants (162). Most trials were small (median number of participants 88 [IQR 39–140]). They were conducted over the past 25 years in neonatal centres in seven countries (Albania, Armenia, Brazil, India, the Islamic Republic of Iran, Italy and the United Kingdom). The inclusion criteria were infants who required primary treatment for RDS after birth or following a period of mechanical ventilation (post-extubation). Most infants were born before 32 weeks' gestation (very preterm). Thirteen trials included both RDS and post-extubation infants, two trials included infants with RDS only and no trials included post-extubation infants only. All trials compared bubble CPAP with ventilator or IFD CPAP devices. The devices were all commercially manufactured; no locally manufactured or locally adapted devices were used. The interfaces in all trials were short nasal prongs. All infants received standard supportive care (i.e. supplemental oxygen).

Critical outcomes

For bubble CPAP compared with ventilator or IFD nasal CPAP, 10 trials reported all-cause mortality, 14 reported morbidity (13 reported "treatment failure", 14 pneumothorax, 7 bronchopulmonary dysplasia and 8 nasal injury). No trials reported growth or neurodevelopment. (Full details are provided in GRADE Table B.3, in the Web Supplement.)

- **Mortality:** Low-certainty evidence from 10 trials totalling 1189 participants suggests little or no effect on all-cause mortality by hospital discharge (RR 0.93, 95% CI 0.64 to 1.36).
- **Morbidity/adverse events:** Low-certainty evidence from 13 trials totalling 1230 participants suggests a decrease in "treatment failure" (defined as recurrent apnoea, hypoxia, hypercarbia, increasing oxygen requirement, or the receipt of mechanical ventilation within 72 hours after initiation of nasal CPAP) by hospital discharge (RR 0.76, 95% CI 0.60 to 0.95). Low-certainty evidence from 14 trials totalling 1340 participants suggests a decrease in pneumothorax (RR 0.73, 95% CI 0.40 to 1.34). Low-certainty evidence from seven trials totalling 603 participants suggests a decrease in bronchopulmonary dysplasia (oxygen dependency at 28 days) (RR 0.76, 95% CI 0.53

to 1.10). Low-certainty evidence from eight trials of 753 participants suggests an increase in nasal injury (defined as ulceration, bleeding, septal injury and scarring but excluding hyperaemia and erythema) by hospital discharge (RR 2.29, 95% CI 1.37 to 3.82).

Other outcomes

There was a decrease in length of hospital stay (in days) (MD -3.27, 95% CI -4.99 to -1.56 days; 5 trials, 591 participants).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (Table 1.1) reported that carers want assistance in interacting with their babies, especially when they are undergoing therapies that make it difficult to have physical contact (14). They want to learn about the health-care setting where they need to stay and care for their baby. They want to understand what medical equipment is being used and why. Studies report that families can find mechanical ventilation and CPAP intimidating and frightening and that these therapies can accentuate their feelings of inadequacy and lack of control over their baby's health care (147,159). Families also worry about the pain and discomfort their baby is experiencing in NICUs (14). Studies from LMICs indicate that bubble CPAP is both valued and acceptable to families and health workers (163,164). No other specific evidence was located about whether families value bubble CPAP rather than other types of CPAP for their preterm or LBW baby or whether they find bubble CPAP more or less acceptable.

Resources required and implementation considerations

Please refer to the information on this topic in section B.1.

Feasibility and equity

Studies from LMICs (165–168) report the low cost and feasibility of establishing bubble CPAP services. There was no other specific evidence on the feasibility and equity of providing CPAP for preterm or LBW infants.

Summary of judgements

Comparison: Bubble CPAP vs other CPAP pressure sources (B.3)

Justification	<ul style="list-style-type: none">• Evidence of small-to-moderate benefits: decreased pneumothorax, decreased bronchopulmonary dysplasia and decreased failed treatment (defined as recurrent apnoea, hypoxia, hypercarbia, increasing oxygen requirement or the need for mechanical ventilation) (<i>low-certainty evidence</i>)• Evidence of small harms: increased nasal injury (defined as ulceration, bleeding, septal injury and/or scarring but excluding hyperaemia and erythema) (<i>low-certainty evidence</i>)• Evidence of little or no effect on mortality (<i>low-certainty evidence</i>)• No evidence on other critical outcomes
----------------------	--

Evidence-to-Decision summary

Benefits	Small to moderate
Harms	Small
Certainty	Low
Balance	Probably favours bubble CPAP
Values	Uncertainty or variability about outcomes
Acceptability	Varies
Resources	Moderate
Feasibility	Varies
Equity	Varies

B.4 METHYLXANTHINES FOR TREATMENT OF APNOEA

Recommendation and remarks

RECOMMENDATION B.4 (NEW)

Caffeine is recommended for the treatment of apnoea in preterm infants. (*Strong recommendation, moderate-certainty evidence*)

Remarks

- The GDG noted that evidence was available for all preterm infants, so caffeine (or other methylxanthines) is recommended for treatment of apnoea in preterm infants.
- The GDG noted that there were limited data on the dose, timing of initiation and duration of administration. Based on the largest trial (169) included in the evidence review, the GDG suggested a 20 mg/kg loading dose and a 5 mg/kg per day maintenance dose for six weeks. The duration of caffeine administration should be based on clinical judgement.
- If caffeine is not available, other methylxanthines (aminophylline or theophylline) may be considered.

Background and definitions

Apnoea (temporary cessation of breathing) is common in preterm infants (170,171). The frequency of apnoea is inversely related to gestational age, and it occurs in almost all infants born before 28 weeks' gestation (extremely preterm) (170,171). Episodes of apnoea can result in hypoxaemia and bradycardia requiring mechanical ventilation. Intermittent hypoxic episodes in the first two months after birth are associated with increased risk of chronic

conditions, such as retinopathy of prematurity, and adverse neurodevelopmental outcomes (172,173). Since the 1970s, methylxanthine medicines such as theophylline, aminophylline and caffeine have been used to manage apnoea. More recently, large pragmatic studies have included methylxanthine treatment for a variety of indications, including the treatment and prevention of apnoea (174). Studies have also assessed the use of methylxanthines to prevent apnoea before and after extubation (169,175).

Summary of the evidence

OVERVIEW	B.4 Methylxanthines for treatment of apnoea
PICO	<p>Population – Preterm infants</p> <p>Intervention – Any methylxanthine (aminophylline, theophylline, caffeine) at any dose</p> <p>Comparator – Placebo or no methylxanthine treatment</p> <p>Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up</p>
Timing, setting, subgroups	<p>Timing of the intervention – Birth to 6 months of age</p> <p>Setting – Health-care facility or home in any country or setting</p> <p>Subgroups</p> <ul style="list-style-type: none"> • Gestational age at birth (< 32 weeks, ≥ 32 weeks) • Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – Methylxanthine for treatment of apnoea versus placebo or no methylxanthine treatment

Sources and characteristics of the evidence

The effectiveness evidence for this comparison was derived from a Cochrane review of 18 RCTs enrolling a total of 2705 preterm infants who received methylxanthines for any indication (174). For the indication relevant to this comparison (for

treatment of apnoea), the inclusion criteria for infants were gestational age at birth below 37 weeks and evidence of apnoea. Six RCTs were included, enrolling a total of 959 preterm infants from six countries (Australia, Canada, France, India, the United Kingdom and the USA). The largest study, the Caffeine for Apnoea of Prematurity (CAP) trial (169), enrolled 767 participants (birth weight 0.5–1.2 kg) from nine countries who received methylxanthines for

treatment of apnoea and conducted follow-up after five years. The other five trials were small, with fewer than 100 infants in each trial.

Critical outcomes

For methylxanthines for treatment of apnoea compared with placebo or no methylxanthine treatment, three trials reported all-cause mortality, five reported morbidity (1 reported apnoea, 5 use of mechanical ventilation, 1 bronchopulmonary dysplasia) and one trial reported a composite outcome of death or major neurodevelopmental disability. No trials reported growth outcomes. (Full details are provided in GRADE Table B.4, in the Web Supplement.)

- **Mortality:** Low-certainty evidence from three trials totalling 154 participants suggests a decrease in all-cause mortality by hospital discharge (RR 0.49, 95% CI 0.14 to 1.78).
- **Morbidity:** Very-low-certainty evidence from one trial with 43 participants suggests a decrease in any apnoeic episodes by hospital discharge (RR 0.70, 95% CI 0.30 to 1.62). Low-certainty evidence from five trials totalling 192 participants suggests a decrease in the use of mechanical ventilation by hospital discharge (RR 0.34, 95% CI 0.12 to 0.97). Moderate-certainty evidence from one trial with 805 participants suggests a decrease in bronchopulmonary dysplasia at 36 weeks PMA (RR 0.72, 95% CI 0.58 to 0.89).
- **Mortality or neurodevelopment:** Moderate-certainty evidence from one trial with 767 participants suggests little or no effect on the composite outcome of death or major neurodevelopmental disability by the latest follow-up (5 years) (RR 0.85, 95% CI 0.71 to 1.01). This composite outcome was defined as death or survival to 5 years with one or more of the following: motor impairment (defined as a gross motor function classification system level of 3–5), cognitive impairment (defined as a full-scale IQ < 70), behaviour problems, poor general health, deafness and/or blindness, all measured using validated tests.

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported (within a theme on the health-care environment) that carers want mechanisms and initiatives to help them to interact with their babies, especially when they are undergoing therapies that make it difficult to have physical contact with the infant (14). They also want to learn about the health-care setting (including the equipment in use) where they need to stay and care for the infant. No other specific evidence was located about whether families value methylxanthines for their preterm or LBW baby or whether they find them more or less acceptable than other medicines or no treatment.

Resources required and implementation considerations

Organization of care

Methylxanthines (caffeine, theophylline and aminophylline) must be dispensed by a health worker. They can be provided in the health-care facility or at home. Caffeine is given once a day and theophylline and aminophylline are given three times a day.

Infrastructure, equipment and supplies

Methylxanthines are available as intravenous and oral formulations. Caffeine citrate is available as 20 mg/ml and 10 mg/ml for intravenous and oral use, respectively. Oral caffeine comes as a ready-to-use formulation that needs no mixing. Theophylline is available as 50–60 mg/5 ml liquid. Aminophylline is available as 25 mg/ml ampoules.

Workforce, training, supervision and monitoring

Health workers at all levels can support mothers and families. Standardized packages are needed for training, supervision and monitoring. Dispensing needs to be documented in clinical records.

Feasibility and equity

Studies report that availability and cost are barriers for the use of caffeine citrate formulations in LMICs (176). Theophylline and aminophylline are more widely available than caffeine in LMICs (31,176). There was no specific evidence on the feasibility and equity of providing methylxanthines for preterm or LBW infants.

Summary of judgements

Comparison: Methylxanthine for the treatment of apnoea in preterm infants vs placebo or no methylxanthine treatment (B.4)

- Justification**
- Evidence of moderate benefits: decreased death, bronchopulmonary dysplasia (*moderate-certainty evidence*), decreased mechanical ventilation (*low-certainty evidence*) and decreased neurodevelopmental disability (*moderate-certainty evidence*)
 - No evidence of harms

Evidence-to-Decision summary

Benefits	Moderate
Harms	Trivial or none
Certainty	Moderate
Balance	Favours methylxanthines for treatment of apnoea in infants < 37 weeks
Values	No uncertainty or variability about outcomes
Acceptability	Acceptable
Resources	Low to moderate
Feasibility	Probably feasible
Equity	Probably equitable

B.5 METHYLYXANTHINES FOR EXTUBATION

Recommendation and remarks

RECOMMENDATION B.5 (NEW)

Caffeine is recommended for the extubation of preterm infants born before 34 weeks' gestation. (*Strong recommendation, moderate-certainty evidence*)

Remarks

- The GDG noted that evidence was available only for preterm infants born before 34 weeks' gestation, but suggests that caffeine (or other methylxanthines) may also be considered for extubation of preterm infants born at or after 34 weeks and before 37 weeks, depending on clinical judgement.
- The GDG noted that there were limited data on the timing of initiation and duration of administration. Based on the largest trials (169,175) included in the evidence review, the GDG suggested starting caffeine 24 hours before a planned extubation. If the extubation is unplanned, the infant should receive the caffeine as soon as possible after the extubation and within 6 hours, and should continue to receive it for six days.
- The GDG noted that there were limited data on the dosage. Based on the largest trials (169,175) included in the evidence review, the GDG suggested a 20 mg/kg loading dose and 5 mg/kg per day maintenance dose for six days.
- If caffeine is not available, other methylxanthines (aminophylline or theophylline) may be considered.

Background and definitions

Please refer to the information in section B.4.

Summary of the evidence

OVERVIEW	B.5 Methylxanthines for extubation
PICO	Population – Preterm infants (< 34 weeks) Intervention – Any methylxanthine (aminophylline, theophylline, caffeine) at any dose Comparator – Placebo or no methylxanthine treatment Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none">▪ Gestational age at birth (< 32 weeks, ≥ 32 weeks)▪ Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – Methylxanthine for extubation versus placebo or no methylxanthine treatment

Sources and characteristics of the evidence

The effectiveness evidence for this comparison was derived from the same Cochrane review of preterm infants who received methylxanthines for any indication (174). For the indication relevant to this comparison (for extubation), the inclusion criteria for infants were gestational age at birth below 34 weeks and planned extubation. Seven RCTs enrolling a total of 870 preterm infants were included from five countries (Australia, Canada, Spain, the United Kingdom and the USA). The largest study

was also the CAP trial (169), which followed up 676 participants who received methylxanthines for extubation. The other six trials were small, with fewer than 100 infants in each trial.

Critical outcomes

For methylxanthines for extubation compared with no methylxanthine treatment, six trials reported morbidity (6 reported “failed extubation”, 2 bronchopulmonary dysplasia) and one trial reported a composite outcome of death or major neurodevelopmental disability. No trials reported growth outcomes. (Full details are provided in GRADE Table B.5, in the Web Supplement.)

- **Morbidity:** Moderate-certainty evidence from six trials totalling 197 participants suggests decreased failed extubation (defined as the infant having to be re-intubated) by hospital discharge (RR 0.48, 95% CI 0.32 to 0.71). Moderate-certainty evidence from two trials totalling 704 participants suggests a decrease in bronchopulmonary dysplasia (defined as a need for supplemental oxygen) by 36 weeks PMA (RR 0.81, 95% CI 0.70 to 0.92).
- **Mortality or neurodevelopment:** Moderate-certainty evidence from one trial with 676 participants suggests decreased death or major neurodevelopmental disability (see section B.4 for

the definition of the composite outcome) by the latest follow-up (5 years) (RR 0.85, 95% CI 0.73 to 0.99).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Values and acceptability, resources, feasibility and equity

Please refer to the information on these topics in section B.4.

Summary of judgements

Comparison: Methylxanthine for extubation in preterm infants vs placebo or no methylxanthine treatment (B.5)

Justification

In trials where most participants are infants born < 34 weeks' gestation:

- Evidence of moderate benefits: decreased death, bronchopulmonary dysplasia, failed extubation and neurodevelopmental disability (*moderate-certainty evidence*)
- No evidence of harms

Evidence-to-Decision summary

Benefits	Moderate
Harms	Trivial or none
Certainty	Moderate
Balance	Favours methylxanthines for extubation in infants < 34 weeks
Values	No uncertainty or variability about outcomes
Acceptability	Acceptable
Resources	Low to moderate
Feasibility	Probably feasible
Equity	Probably equitable

B.6 METHYLYXANTHINES FOR PREVENTION OF APNOEA

Recommendation and remarks

RECOMMENDATION B.6 (NEW)

Caffeine may be considered for the prevention of apnoea in preterm infants born before 34 weeks' gestation. (Conditional recommendation, low-certainty evidence)

Remarks

- The GDG noted that the evidence on increased mortality came from three small trials totalling 129 infants (177-179) and was uncertain due to very low quality, and imprecision. Also, data on "death alone" were not available from a large trial of 423 infants (169), which reported no effect on a combined outcome of death and neurodevelopmental disability. The GDG also noted that the evidence on harms from increased use of mechanical ventilation was uncertain due to very low quality, and imprecision.
- The recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- The GDG noted that evidence was available only for preterm infants born before 34 weeks' gestation, but suggests that caffeine (or other methylxanthines) may also be considered for prevention of apnoea in preterm infants born at or after 34 weeks and before 37 weeks if there is clinical indication.
- The GDG noted that there were limited data on the dose, timing of initiation and duration of administration. Based on the largest trial (169) included in the evidence review, the GDG suggested a 20 mg/kg loading dose and a 5 mg/kg per day maintenance dose for six weeks. The duration of caffeine administration should be based on clinical judgement.
- If caffeine is not available, other methylxanthines (aminophylline or theophylline) may be considered.

Background and definitions

Please refer to the information in section B.4.

Summary of the evidence

OVERVIEW	B.6 Methylxanthines for prevention of apnoea
PICO	Population – Preterm infants (< 34 weeks) Intervention – Any methylxanthine (aminophylline, theophylline, caffeine) at any dose Comparator – Placebo or no methylxanthine treatment Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none">• Gestational age at birth (< 32 weeks, ≥ 32 weeks)• Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – Methylxanthines for prevention of apnoea versus placebo or no methylxanthine treatment

Sources and characteristics of the evidence

The effectiveness evidence for this comparison was derived the same Cochrane review of preterm infants who received methylxanthines for any indication (174). For the indication relevant to this comparison (for prevention of apnoea), the inclusion criteria for infants were gestational age at birth below 34 weeks

and no evidence of apnoea. Seven RCTs enrolling a total of 706 preterm infants were included from six countries (Australia, Canada, the Islamic Republic of Iran, Switzerland, the United Kingdom and the USA). The largest study was also the CAP trial, which followed up 423 participants who received methylxanthines for prevention of apnoea (169). The other six trials were small, with fewer than 100 infants in each.

Critical outcomes

For methylxanthines for prevention of apnoea compared with no methylxanthines, three trials reported all-cause mortality, four reported morbidity (2 reported apnoea, 4 use of mechanical ventilation, 3 bronchopulmonary dysplasia) and one reported a composite outcome of death or neurodevelopmental disability. No trials reported growth outcomes. (Full details are provided in GRADE Table B.6, in the Web Supplement.)

- **Mortality:** Low-certainty evidence from three trials (177-179) totalling 129 participants suggests little or no effect on mortality by hospital discharge (RR 2.19, 95% CI 0.85 to 5.68).
- **Morbidity:** Low-certainty evidence from two trials totalling 104 participants suggests a decrease in any apnoeic episodes by hospital discharge (RR 0.19, 95% CI 0.09 to 0.41). Low-certainty evidence from four trials totalling 208 participants suggests little or no effect on the use of mechanical ventilation by hospital discharge (RR 1.33, 95% CI 0.48 to 3.72). Moderate-certainty evidence from three trials totalling 541 participants suggests a

decrease in bronchopulmonary dysplasia (defined as the use of supplemental oxygen at 36 weeks PMA) (RR 0.78, 95% CI 0.63 to 0.97).

- **Mortality or neurodevelopment:** Moderate-certainty evidence from one trial with 423 participants suggests no effect on the composite outcome of death or neurodevelopmental disability (see section B.4 for the definition of the composite outcome) by latest follow-up (5 years) (RR 1.00, 95% CI 0.80 to 1.24). Data on death alone and neurodevelopmental disability alone were not available for this trial.

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Values and acceptability, resources, feasibility and equity

Please refer to the information on these topics in section B.4.

Summary of judgements

Comparison: Methylxanthine for the prevention of apnoea in preterm infants vs placebo or no methylxanthine treatment (B.6)

Justification

In trials where most participants are infants born < 34 weeks' gestation:

- Evidence of small-to-moderate benefit: decreased bronchopulmonary dysplasia (*moderate-certainty evidence*) and decreased apnoeic episodes (*low-certainty evidence*)
- Evidence of harms uncertain: little or no effect on mortality (*low-certainty evidence*), little or no effect on combined outcome of neurodevelopment or death (*moderate-certainty evidence*) and increase in use of mechanical ventilation (*low-certainty evidence*)
- No evidence on other critical outcomes

Evidence-to-Decision summary

Benefits	Small to moderate
Harms	Unknown
Certainty	Low
Balance	Probably favours methylxanthines for prevention of apnoea in infants < 34 weeks
Values	Uncertainty or variability about outcomes
Acceptability	Varies
Resources	Low to moderate
Feasibility	Probably feasible
Equity	Probably equitable

C. Family involvement and support

C.1 FAMILY INVOLVEMENT IN ROUTINE CARE

Recommendation and remarks

RECOMMENDATION C.1 (NEW)

Family involvement in the routine care of preterm or low-birth-weight infants in health-care facilities is recommended. (Strong recommendation, low- to moderate-certainty evidence)

Remarks

- The trials in the systematic review varied widely in intervention content, intensity and effect but all showed consistent and similar effects.
- The GDG noted that the resources needed for and the feasibility of implementing family involvement strategies vary according to setting but that simple family involvement interventions such as the delivery of direct bedside care and involvement in medical decision-making could be implemented in all settings. Other components that can be provided include chairs near the infant's cot, even in busy and crowded hospital wards.
- The GDG also noted that family involvement strategies reduced the length of hospital stay, improved breastfeeding and reduced parental anxiety and stress.

Background and definitions

Preterm and LBW infants commonly require specialized care, close monitoring and medical interventions (2,180). In some health-care facilities, families are not allowed any physical access to their infants and receive only intermittent verbal updates from health workers (181-184). Family involvement is often defined as the participation of mothers, fathers/partners and other family members in routine care of the newborn while the baby is in the health-care facility (180,185,186). It may include promotion of direct bedside care from the family (e.g. feeding and administration of medicines), inclusion of the family in medical decision-making, infrastructure

changes (e.g. beds and chairs near the baby's cot, family rooms), health-care facility culture change and health worker behaviour change. Strategies to increase family involvement have typically focused on packages of one or more of these interventions with the overall aims of increasing the amount of direct hands-on care that parents provide for their infant and empowering families to collaborate in health-care decision-making. Well known packages that are implemented in high-, middle- and low-income countries include family-centred care, family-participatory care and family-integrated care (180,185,186).

Summary of the evidence

OVERVIEW	C.1 Family involvement
PICO	Population – Hospitalized preterm or LBW infants Intervention – Interventions to involve families in their infant's routine health care Comparator – Usual hospital care Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Hospital in any country or setting Subgroups <ul style="list-style-type: none">• Gestational age at birth (< 32 weeks, ≥ 32 weeks)• Birth weight (< 1.5 kg, ≥ 1.5 kg)• Intensity of interventions (high intensity ≥ 12 hours per day, low intensity < 12 hours per day)

Effectiveness: Comparison – Family involvement in routine care versus usual hospital care

Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review of 15 RCTs enrolling a total of 5240 preterm or LBW infants from nine countries (Australia, Canada, China, India, the Islamic Republic of Iran, New Zealand, the Republic of Korea, Sweden and the USA) (187). Most infants were born before 32 weeks' gestation or had birth weight below 1.5 kg, and most trials excluded infants with major congenital anomalies. All trials were conducted in NICUs.

All trials evaluated the effect of family-centred models or packages for the hospital care of preterm or LBW infants on infant and parental outcomes. No studies of infrastructure or behaviour change interventions were located. The family-centred packages were heterogeneous, but their common core content was the involvement of family members in the provision of direct bedside care. Skin-to-skin care or kangaroo mother care (KMC) was included in nine trials, though frequency and duration were not described. Other common components included neurodevelopmental care (8 trials), preparation for transition to home (6 trials) and the involvement of parents in medical decision-making (4 trials).

Critical outcomes

For family involvement strategies compared with usual hospital care, four trials reported all-cause mortality outcomes, eight reported morbidity (6 reported serious infection, 6 necrotizing enterocolitis, 7 bronchopulmonary dysplasia, 8 retinopathy of prematurity and 5 intraventricular haemorrhage), three reported growth (weight gain) and two reported neurodevelopment. (Full details are provided in GRADE Table C.1, in the Web Supplement.)

- **Mortality:** Very-low-certainty evidence from four trials totalling 2378 participants suggests little or no effect on all-cause mortality by hospital discharge (OR 1.05, 95% CI 0.53 to 2.09).
- **Morbidity:** Low-certainty evidence from six trials totalling 2843 participants suggests a decrease in serious infection by hospital discharge (OR 0.79, 95% CI 0.53 to 1.16). Low-certainty evidence from six trials totalling 2809 participants suggests little or no effect on necrotizing enterocolitis by hospital discharge (OR 0.81, 95% CI 0.46 to 1.44). Low-certainty evidence from seven trials totalling 3085 participants suggests decreased bronchopulmonary dysplasia by hospital discharge (OR 0.74, 95% CI 0.53 to 1.03). Moderate-

certainty evidence from eight trials totalling 2551 participants suggests decreased retinopathy of prematurity by hospital discharge (OR 0.52, 95% CI 0.34 to 0.80). Very-low-certainty evidence from five trials totalling 2555 participants suggests decreased intraventricular haemorrhage by hospital discharge (OR 0.74, 95% CI 0.36 to 1.54).

- **Growth:** Moderate-certainty evidence from three trials totalling 2215 participants suggests increased in-hospital growth velocity (grams per day) (MD 2.09, 95% CI 1.27 to 2.91).
- **Neurodevelopment:** Low-certainty evidence from two trials totalling 422 participants suggests increased neurodevelopment (measured using the Neonatal Neurobehavioral Examination – Chinese version [NNE-C] test) by hospital discharge or term corrected age, i.e. 37 weeks PMA (MD 1.11, 95% CI 0.21 to 2.01) (187).

Other outcomes

There was a decrease in length of hospital stay (in days) (MD -2.91, 95% CI -5.15 to -0.68; 11 trials, 4452 participants). There was an increase in the proportion of infants predominantly or exclusively breastfeeding by hospital discharge (OR 1.34, 95% CI 1.10 to 1.65; 3 trials, 1739 participants). There was an increase in “any” breastfeeding by hospital discharge (OR 2.60, 95% CI 0.77 to 8.79; 5 trials, 2546 participants).

Subgroup analyses

For gestational age and birth weight, differences for weight gain and neurodevelopment could not be assessed as there were insufficient studies. For the other outcomes there was no evidence of a subgroup difference.

For the intensity of intervention, differences for intraventricular haemorrhage, weight gain and neurodevelopment could not be assessed as there were insufficient studies. For the other outcomes there was no evidence of a subgroup difference except for bronchopulmonary dysplasia, which decreased after high-intensity interventions (RR 0.18, 95% CI 0.05 to 0.66; 1 study, 366 participants) but not after low-intensity interventions (RR 1.04, 95% CI 0.68 to 1.58; 6 studies, 2719 participants) (test for subgroup differences, $\text{Chi}^2 = 7.22$, $P = 0.007$).

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, and want to take an

active role in deciding what interventions are given to infants, in the routine care of the newborn, in direct bedside care, including feeding their baby and in medical decision-making, and that they value hospital infrastructure changes (e.g. beds and chairs near the baby's cot, family rooms) (14). No other specific evidence was located.

Resources required and implementation considerations

Organization of care

Family involvement strategies can be implemented at all levels of newborn care (primary, secondary and tertiary). Health-care facilities should ensure that families have access to beds, food, bathing and toilet facilities throughout the infant's hospital stay.

Infrastructure, equipment and supplies

No special infrastructure, equipment or supplies are needed to support family involvement in the care of their preterm or LBW infants. However, many arrangements can make the infant and mother more comfortable, e.g. reclining beds and chairs. More structured packages may include infrastructure changes such as beds and chairs near the infant's cot, and family rooms.

If couplet care or maternal-newborn intensive care units (M-NICUs) are used, they should have all the infrastructure, equipment and supplies that NICUs have for small or sick babies and that maternity wards have for mothers. For infants, this includes CPAP machines, pulse oximeters, and radiant warmers or incubators if the infant is not in KMC. For mothers, this includes adult beds and an examination area where she can receive the health care she needs.

Workforce, training, supervision and monitoring

Health workers at all levels can support family involvement in the routine care of their preterm or LBW infant. Standardized packages can be used for training, supervision and monitoring. This can include the promotion of direct bedside care from the family (e.g. feeding and administration of medicines), inclusion of the family in medical decision-making, health-care facility culture change, health worker behaviour change and infrastructure change.

Feasibility and equity

There was no specific evidence on the feasibility and equity of promoting family involvement for preterm or LBW infants.

Summary of judgements

Comparison: Family involvement in routine care vs usual hospital care (C.1)

Justification

- Evidence of moderate benefits: decreased morbidity (infection, intraventricular haemorrhage, retinopathy of prematurity, bronchopulmonary dysplasia), increased weight and length, and increased neurodevelopment (*low- to moderate-certainty evidence*)
- No evidence of harms
- Evidence of little or no effect on: mortality, necrotizing enterocolitis, and weight and head circumference (*low- to very-low-certainty evidence*)
- No evidence on other critical outcomes

Evidence to-Decision summary

Benefits	Moderate
Harms	Trivial or none
Certainty	Low to moderate
Balance	Favours family involvement strategies
Values	No uncertainty or variability about outcomes
Acceptability	Acceptable
Resources	Vary
Feasibility	Varies
Equity	Probably equitable

C.2 FAMILY SUPPORT

Recommendation and remarks

RECOMMENDATION C.2 (NEW)

Families of preterm or low-birth-weight infants should be given extra support to care for their infants, starting in health-care facilities from birth, and continued during follow-up post-discharge. The support may include education, counselling and discharge preparation by health workers, and peer support.

(Conditional recommendation, very-low-certainty evidence)

Remarks

- The recommendation is conditional on shared decision-making with parents; this includes informing parents about the benefits and risks and the need for further research.
- The GDG noted that **education and counselling** also had important effects in improving parent-to-infant interaction, improving breastfeeding and decreasing parental anxiety, stress and depression, though these were not critical outcomes.
- The GDG noted that there were limited data on frequency, duration and intensity of **education and counselling**.
- The GDG noted that **discharge preparation** also had important effects in improving parent-to-infant interaction, improving breastfeeding and decreasing parental anxiety, stress and depression, though these were not critical outcomes.
- The GDG noted that there were limited data on the frequency, duration and intensity of **discharge preparation**.
- Preterm and LBW infants often require care from multiple health workers so the GDG also noted that careful coordination of care is needed post-discharge.
- The GDG made a conditional recommendation on **peer support**, although there were no data on critical outcomes; this was because of the effects on maternal anxiety and the importance of the intervention.
- The GDG noted that there were limited data on frequency, duration and intensity of **peer support**.
- The GDG decided not to make a recommendation on **digital information systems** as there was no evidence of benefits or harms on any critical outcome.

Background and definitions

Supporting families to care for their sick, vulnerable, preterm or LBW infant is a basic and integral component of any health system. However, many families still feel ill-equipped to care for their preterm or LBW newborn infant at home (188,189). Families need support at all stages, starting from before conception, and including at the identification of a high-risk pregnancy, at the birth of the baby, in the health-care facility, at discharge and when the baby reaches home. Much of the support that families need to care for their preterm and LBW infants is provided through social services in high-, middle- and low-income countries. However, “what the health system can do” and the “health system building blocks” they can use (i.e. service delivery, workforce, digital information systems, medical products and technologies, financing, leadership and governance) (190) are often overlooked. Two systematic

reviews have recently assessed the effectiveness of communication and peer-support interventions for families of preterm infants (191,192). However, there have been no recent systematic reviews of the effects of other health system “building blocks” on infant mortality, morbidity, growth and neurodevelopmental outcomes.

Overall, the effectiveness evidence was derived from a systematic review of 37 trials (35 RCTs and 2 non-randomized) enrolling a total of 11 758 preterm or LBW infants from 18 countries (193) (Australia, Bangladesh, Canada, China, Denmark, Egypt, Finland, Greece, India, the Islamic Republic of Iran, Jamaica, the Netherlands, Norway, the Philippines, the Republic of Korea, Sweden, the United Kingdom and the USA). No studies were based in low-income settings. Interventions commenced either in the facility (24 trials) or in the home (13 trials). All

began after birth; no intervention started during pregnancy. No studies assessed the effect of the “usual support” that is provided to all babies, while all studies assessed only “extra support” (i.e. additional or strengthened support) needed for preterm and LBW infants. The interventions included in the studies were education and counselling (18 trials), peer support (2), discharge preparation (1), digital information systems (4) and home visits by a trained

health worker or volunteer (9). No studies on the other health system building blocks – including financing, leadership or governance – were identified. The education and counselling, peer support and discharge preparation interventions are described below. Home visiting interventions are described in section C.3. Parental leave, financing and entitlements are described in section C.4.

Summary of the evidence

OVERVIEW	C.2a Education and counselling	C.2b Peer support	C.2c Discharge preparation	C.2d Digital information
PICO	<p>Population – Families of preterm or LBW infants</p> <p>Intervention 1 – Education and counselling interventions</p> <p>Intervention 2 – Peer support interventions</p> <p>Intervention 3 – Discharge preparation interventions</p> <p>Intervention 4 – Digital information interventions</p> <p>Comparator – Usual care</p> <p>Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up</p>			
Timing, setting, subgroups	<p>Timing of the intervention – Birth to 6 months of age</p> <p>Setting – Health-care facility or home in any country or setting</p> <p>Subgroups</p> <ul style="list-style-type: none"> Gestational age at birth (< 32 weeks, ≥ 32 weeks) Birth weight (< 1.5 kg, ≥ 1.5 kg) 			

Effectiveness: Comparison 1 – Education and counselling versus usual care

Sources and characteristics of the evidence

For comparison 1, the effectiveness evidence was derived from the systematic review, which included four trials enrolling a total of 312 preterm or LBW infants (193). The interventions included individual or group education or training (provided by health workers) of families to care for their preterm or LBW infant. Content included well-being strategies (e.g. strategies for managing stress, anxiety, depression, self-efficacy) and basic newborn-care practices (e.g. positioning, bathing, breastfeeding, thermal care, responsiveness and sensitivity). The interventions began in the facility, with some continuing at home following discharge.

Critical outcomes

For education and counselling compared with usual care, two trials reported growth (weight gain, length gain) and three reported neurodevelopment (cognitive and motor development). No trials reported mortality or morbidity. (Full details are provided in GRADE Table C.2a, in the Web Supplement.)

- **Growth:** Very-low-certainty evidence from one trial with 184 participants suggests an increase in infant weight (in grams) at 60 days (MD 305, 95% CI 228 to 382). Very-low-certainty evidence from one trial with 57 participants suggests an increase in infant weight (in grams) at 120 days (MD 410, 95% CI 406.03 to 413.97). Very-low-certainty evidence from one trial with 184 participants suggests an increase in infant length (in centimetres) at 60 days (MD 1.5, 95% CI 1.08 to 1.92). Very-low-certainty evidence from one trial with 57 participants suggests an increase in infant length (in centimetres) at 120 days (MD 1.2, 95% CI 0.2 to 2.6).
- **Neurodevelopment:** Very-low-certainty evidence from one trial with seven participants suggests little or no effect on motor development (BSID-III) at 6 months of age (MD 0.38, 95% CI -1.15 to 1.19). Low-certainty evidence from three trials totalling 64 participants suggests an increase in cognitive development (BSID-III) at 4–6 months of age (SMD 0.67, 95% CI 0.16 to 1.17).

Other outcomes

There was little or no effect on infant temperament at 6 months of age (SMD 0.26, 95% CI -0.29 to 0.81; 2 trials, 155 participants). There was an increase in mother-infant interaction at 6 weeks (MD 1.8, 95% CI 0.21 to 3.81; 1 trial, 142 participants), 3 months (MD 0.8, 95% CI 0.6 to 2.2; 1 trial, 196 participants) and 6 months of age (MD 0.21, 95% CI 0.11 to 0.67; 1 trial, 63 participants), but there was little to no effect at follow-up at 12 months of age (MD 0.1, 95% CI -0.01 to 0.21; 1 trial, 93 participants). There was little to no effect on duration of exclusive breastfeeding (EBF) (MD 2.0, 95% CI -5.48 to 9.48; 1 trial, 128 participants), but there was an increase in EBF at 2–3 months (RR 1.71, 95% CI 1.26 to 2.31; 2 trials, 244 participants).

Effectiveness: Comparison 2 – Peer support versus usual care

Sources and characteristics of the evidence

For comparison 2, the effectiveness evidence was derived from the systematic review, which included two trials enrolling a total of 118 preterm or LBW infants (193). The peer supporters were all women who had cared for a preterm or LBW infant in a similar environment and were willing to use their experiences to support others. The interventions all commenced in the facility and took place either following agreement from the parent or were initiated by the parent. Content included well-being strategies and newborn-care practices.

Critical outcomes

For peer support compared with usual care, no trials reported mortality, morbidity, growth or neurodevelopment. (Full details are provided in GRADE Table C.2b, in the Web Supplement.)

Other outcomes

There was a decrease in maternal anxiety when the baby reached 4 months of age (SMD 0.74 lower, 95% CI 1.32 lower to 0.16 lower; 1 trial, 49 participants). There was little or no effect on EBF (intervention group: median 3 months [range 0–14]; control group: median 4.3 [range 0–13]; 1 trial, 69 participants).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Effectiveness: Comparison 3 – Discharge preparation versus usual care

Sources and characteristics of the evidence

For comparison 3, the effectiveness evidence was derived from the systematic review, which included one trial enrolling 173 preterm or LBW infants (193). The interventions were delivered by health workers in the days just prior to hospital discharge and focused specifically on preparing parents for the discharge of their infant. The content included well-being strategies and newborn-care practices, but also “anticipatory guidance” (i.e. what to expect), financial and social support information, and referral pathways.

Critical outcomes

For discharge preparation compared with usual care, one trial reported morbidity (emergency department presentations). No trials reported mortality, growth or neurodevelopment outcomes. (Full details are provided in GRADE Table C.2c, in the Web Supplement.)

- **Morbidity:** Very-low-certainty evidence from one observational study with 173 participants suggests a decrease in emergency hospital visits by 2 months of age (RR 0.62, 95% CI 0.39 to 1.00).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Effectiveness: Comparison 4 – Digital information systems versus usual care

Sources and characteristics of the evidence

For comparison 4, the effectiveness evidence was derived from the systematic review, which included four trials enrolling a total of 902 preterm or LBW infants (193). The interventions used electronic web-based applications, including Skype, audiovisual workshops and telephone media. Content included well-being strategies and newborn-care practices. The interventions commenced either in the facility or at home.

Critical outcomes

One trial reported morbidity (emergency department presentations). No trials reported mortality, growth or neurodevelopment outcomes. (Full details are provided in GRADE Table C.2d, in the Web Supplement.)

■ **Morbidity:** Very-low-certainty evidence from one trial with 89 participants suggests little to no effect on emergency hospital visits by two months post-discharge (usual care group: median 1 visit [range 0–6 visits] versus digital information systems group: median 1 visit [range 0–7 visits]).

Other outcomes

There was little or no effect on maternal–infant interaction by 1 month of age (MD -0.8, 95% CI -1.84 to 0.24; 1 trial, 129 participants) or by 4 months of age (MD -0.9, 95% CI -2.09 to 0.29; 1 trial, 85 participants). There was little or no effect on EBF by 2 months of age (RR 1.02, 95% CI 0.89 to 1.16; 2 trials, 688 participants).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting all newborn-care practices, and want to take an active role in deciding what interventions are given to infants, including what newborn-care practices

they receive and how they are implemented (14). No specific evidence was located about the kinds of support families of preterm or LBW babies value or find acceptable.

Resources required and implementation considerations

Organization of care

Families may need education, counselling, discharge preparation and peer support at all levels of health facility care. Education, counselling and peer support may be needed at home. Support and planning should be started in the antenatal period where possible. Services should follow national and local guidance for health-care facilities.

Infrastructure, equipment and supplies

National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can provide family support. Standardized packages can be used for training, supervision and monitoring.

Feasibility and equity

There was no specific evidence on the feasibility and equity of providing family support for preterm or LBW infants.

Summary of judgements

	Comparison 1. Education and counselling vs usual care (C.2a)	Comparison 2. Peer support vs usual care (C.2b)	Comparison 3. Discharge preparation vs usual care (C.2c)	Comparison 4. Digital information systems vs usual care (C.2d)
Justification	<ul style="list-style-type: none"> Evidence of moderate benefits: increase in weight, length and neurodevelopment (<i>very-low-certainty evidence</i>) No evidence of harms No evidence on other critical outcomes 	<ul style="list-style-type: none"> Evidence of small benefits: decrease in maternal anxiety (<i>very-low-certainty evidence</i>) No evidence of harms No evidence on critical outcomes 	<ul style="list-style-type: none"> Evidence of small benefits: decrease in emergency department presentations (<i>very-low-certainty evidence</i>) No evidence of harms No evidence on other critical outcomes 	<ul style="list-style-type: none"> Evidence of little to no effect on emergency hospital visits (<i>very-low-certainty evidence</i>) No evidence on other critical outcomes
Evidence-to-Decision summary				
Benefits	Moderate	Small	Small	Unknown
Harms	None	None	None	Unknown
Certainty	Very low	Very low	Very low	Very low
Balance	Probably favours education and counselling	Probably favours peer support	Probably favours discharge preparation	Unknown
Values	No uncertainty or variability about outcomes	No uncertainty or variability about outcomes	No uncertainty or variability about outcomes	No uncertainty or variability about outcomes
Acceptability	Probably acceptable	Probably acceptable	Probably acceptable	Probably acceptable
Resources	Moderate	Moderate	Moderate	Moderate
Feasibility	Probably feasible	Probably feasible	Probably feasible	Varies
Equity	Probably equitable	Probably equitable	Probably equitable	Varies

C.3 HOME VISITS

Recommendation and remarks

RECOMMENDATION C.3 (NEW)

Home visits by trained health workers are recommended to support families to care for their preterm or low-birth-weight infant. (*Strong recommendation, moderate-certainty evidence*)

Remarks

- Trained health workers can include nurses, midwives, doctors and community health workers.
- The GDG noted that there were limited data on the content, frequency, duration and intensity of home visits for preterm and LBW infants. Based on the trials included in the evidence review, the GDG recommended that extra home visits (i.e. additional to the routine scheduled postnatal contacts for all infants [22]) should be made, and that their content, frequency, duration and intensity should be based on clinical judgement.
- The GDG noted that home visits also increased exclusive breastfeeding, immunization visits and parental-infant attachment and decreased parental stress, though these were not critical outcomes.

Background and definitions

Families need support at all stages, from before conception, and including at the identification of a high-risk pregnancy, at the birth of the baby, in the health-care facility, at discharge, and especially when the baby reaches home (189,194). Studies over the last 10 years in high-, middle- and low-income

countries have shown that home visiting during the antenatal and postnatal periods can improve both the demand for and the use of antenatal, delivery and postnatal services and reduce maternal and newborn mortality (22,195). However, there is limited information on the effects of home visiting for preterm and LBW infants.

Summary of the evidence

OVERVIEW	C.3 Home visits
PICO	Population – Families of preterm or LBW infants Intervention – Home visits to support families to care for their preterm or LBW infant in the home Comparator – Usual care Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none">• Gestational age at birth (< 32 weeks, ≥ 32 weeks)• Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – Home visits to support families to provide care versus usual care

Sources and characteristics of the evidence

The effectiveness evidence was derived from a systematic review of nine trials enrolling a total

of 8742 preterm or LBW infants from India, the Netherlands, Taiwan (China) and the USA (193). The interventions were delivered by health workers, community health workers, trained intervention workers or trained volunteers. They started and continued in the home, immediately following

discharge from the facility. The content included well-being strategies and newborn-care practices but also “anticipatory guidance” (i.e. what to expect), financial and social support information, and referral pathways.

Critical outcomes

For home visits to support families to provide care compared with usual care, two trials reported all-cause mortality, one trial reported morbidity (hospitalizations) and two trials reported neurodevelopment (cognitive and motor neurodevelopment). No trials reported growth outcomes. (Full details are provided in GRADE Table C.3, in the Web Supplement.)

- **Mortality:** Moderate-certainty evidence from one trial with 6984 participants suggests decreased all-cause mortality by 180 days of age (RR 0.71, 95% CI 0.57 to 0.89). Low-certainty evidence from one observational study with 970 participants suggests decreased all-cause mortality by 12 months (RR 0.14, 95% CI 0.02 to 1.16).
- **Morbidity:** Low-certainty evidence from one observational study with 970 participants suggests a decrease in hospitalizations by 12 months (MD 0.34, 95% CI 0.16 to 0.52).
- **Neurodevelopment:** Moderate-certainty evidence from two trials totalling 652 participants suggests little or no effect on cognitive neurodevelopment (BSID-III) by 12 months (SMD 0.03, 95% CI -0.12 to 0.19). Low-certainty evidence from one trial with 136 participants suggests little or no effect on motor neurodevelopment (BSID-III) by 12 months (MD 0.02, 95% CI -0.35 to 0.32).

Other outcomes

There was little or no effect on infant temperament at 6 months of age (MD 0.70, 95% CI -0.60 to 1.46; 1 trial, 161 participants) or parent-infant attachment at 6 months of age (MD -1.20, 95% CI -2.79 to 0.39; 1 trial, 136 participants).

There was an increase in EBF at 6 months (RR 4.48, 95% CI 0.28 to 72.9; 3 trials, 7221 participants) and an increase in immunization visits in the first

year (MD 1.21, 95% CI 0.93 to 1.49; 1 trial, 970 participants).

Subgroup analyses

The effect of gestational age and birth weight could not be assessed as there were insufficient trials for any critical outcome.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want to be involved in delivering care to infants, including supporting all newborn-care practices, and want to take an active role in deciding what interventions are given to infants, including what newborn-care practices they receive and how they are implemented (14). No specific evidence was located about whether families value home visiting for their preterm or LBW baby or whether they find it more or less acceptable than other care.

Resources required and implementation considerations

Organization of care

A minimum of four postnatal care contacts is recommended for all infants (22). Extra home visits (i.e. additional to the routine scheduled postnatal contacts for all infants) are needed for preterm and LBW babies. Their content, frequency, duration and intensity should follow national and local guidance for health-care facilities and should be based on clinical judgement.

Infrastructure, equipment and supplies

National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can provide home visits. However, standardized packages are needed for training, supervision and monitoring. Further guidance on follow-up care is being developed and will be published separately.

Feasibility and equity

There was no specific evidence about the feasibility and equity of home visiting interventions for preterm or LBW infants. Home visiting is a core part of the

health programmes for both term and preterm infants in many high-, middle- and low-income countries (22,195).

Summary of judgements

Comparison: Home visits to support families to provide care vs usual care (C.3)	
Justification	<ul style="list-style-type: none">▪ Evidence of moderate benefits: moderate decrease in mortality (<i>moderate-certainty evidence</i>) and small decrease in number of hospitalizations (<i>very-low-certainty evidence</i>)▪ Evidence of little or no effect on cognitive or motor neurodevelopment (<i>low- to moderate-certainty evidence</i>)▪ No evidence of harms▪ No evidence on other critical outcomes
Evidence-to-Decision summary	
Benefits	Moderate
Harms	Trivial or none
Certainty	Low to moderate
Balance	Favours home visits
Values	No uncertainty or variability about outcomes
Acceptability	Probably acceptable
Resources	Moderate
Feasibility	Probably feasible
Equity	Probably equitable

C.4 PARENTAL LEAVE AND ENTITLEMENTS

Good practice statement and remarks

GOOD PRACTICE STATEMENT C.4 (NEW)

Parental leave and entitlements should address the special needs of mothers, fathers and other primary caregivers of preterm or low-birth-weight infants.

Remarks

- The GDG made this good practice statement in recognition of the costs and burdens to parents and families of implementing preterm and LBW infant care.
- Based on the studies in the review, the GDG considered that parental leave and entitlements should include additional days of leave from work and additional financial payments. However, there was insufficient information available to enable the GDG to make recommendations about the number of days of leave parents should be given or what type of financial entitlements they should receive.
- The GDG also noted that the special needs of mothers and fathers/partners of preterm and LBW infants vary according to individual preferences and setting. They include: support for long hospital stays, multiple medical appointments, transport and equipment; support to help manage stress and anxiety about the infant; and support for caring for other children and family members.
- The GDG noted that parental leave and entitlements are in place in some countries but recommended that they should be expanded globally across high-, middle- and low-income countries.

Background and definitions

Families of preterm and LBW infants are well known to have increased risks of financial impoverishment, stress, anxiety and depression (188,195,196). Leave from work is needed to help families care for the infant. Families may also need financial support for transport and equipment as well as for the costs of the hospitalization and caring for other children

or family members (189,191,197). Government and regulatory policies and entitlements are important ways to ensure families receive the financial and workplace support they need. However, there have been few reviews of policies for parental leave and entitlements for families of preterm or LBW infants across high-, middle- and low-income countries.

Summary of the evidence

OVERVIEW	C.4 Parental leave and entitlements
PICO	Population – Preterm or LBW infants Intervention – Parental leave and entitlements Comparator – Usual care Outcomes – All-cause mortality, morbidity, growth, neurodevelopment at latest follow-up
Timing, setting, subgroups	Timing of the intervention – Birth to 6 months of age Setting – Health-care facility or home in any country or setting Subgroups <ul style="list-style-type: none"> • Gestational age at birth (< 32 weeks, ≥ 32 weeks) • Birth weight (< 1.5 kg, ≥ 1.5 kg)

Effectiveness: Comparison – Parental leave and entitlements versus usual care

Sources and characteristics of the evidence

A systematic review of 37 trials (35 RCTs and 2 non-randomized studies) located no studies of the effectiveness of parental leave and entitlements in terms of critical infant outcomes (mortality, morbidity, growth, neurodevelopment) or family outcomes (stress, anxiety, depression) (193).

An additional policy review was done of the most recent relevant policy reports:

- (i) WHO sexual, reproductive, maternal, newborn, child and adolescent health policy survey, 2018–2019 (2018) (198);
- (ii) International Labour Organization database on conditions of work and employment programmes (2022) (199);
- (iii) International Network on Leave Policies and Research (2021) (200).

One hundred and forty countries had policies for parental leave for childhood illness or complications. Twenty-eight countries had a parental leave policy specifically formulated for families of preterm infants: 20 high-income countries (Austria, Canada, Chile, Croatia, Cyprus, Finland, France, Germany, Hungary, Israel, Romania, Italy, Latvia, Lithuania, Luxembourg, New Zealand, Portugal, Slovenia, Spain and the United Kingdom), 6 upper-middle-income countries (Argentina, Belarus, Bulgaria, India, South Africa and Uruguay), 1 lower-middle-income (India) and 1 low-income country (Yemen). Seventeen countries only had policies for maternity leave (Argentina, Austria, Bulgaria, Canada, Chile, Croatia, Finland, France, Hungary, India, Italy, Latvia, Lithuania, Luxembourg, New Zealand, South Africa and Spain) and six had policies for both maternity and paternity leave (Cyprus, Germany, Portugal, Slovenia, the United Kingdom and Uruguay). Five countries did not specify whether the leave was maternal, paternal or both (Belarus, Israel, Romania, Türkiye and Yemen). The amount of leave time was equivalent to the number of weeks early that the baby was born in most cases.

Two countries – Canada and Germany – reported that they provided families with additional financial support for their preterm infants, called “parental allowance”, but details were not available.

Values and acceptability

The systematic review about what matters to families about the care of the preterm or LBW infant (see Table 1.1) reported that families want workplace support, parental leave and financial incentives – especially support for the costs of accommodation, treatment, hospitalization and transport (14). No other specific evidence was located about what types of policies and entitlements for parental leave and financial support families value or find acceptable.

Resources required and implementation considerations

Organization of care

Families need leave and entitlements when the infant is in the health-care facility and also at home, after discharge. Support and planning should be started in the antenatal period where possible or from the time of birth. Services should follow national and local guidance for health-care facilities.

Infrastructure, equipment and supplies

National or local guidance for health-care facilities should be used.

Workforce, training, supervision and monitoring

Health workers at all levels can provide support and referral for parental leave and entitlements, though detailed discussions are often managed by social care staff. Services should follow national and local guidance for health-care facilities. Standardized packages can be used for training, supervision and monitoring.

Feasibility and equity

There was no specific evidence about the feasibility and equity of parental leave and entitlements for preterm or LBW infants.

Summary of judgements

Comparison: Parental leave and entitlements vs usual care (C.4)

Justification

- There were no studies comparing the benefits and harms of parental leave and entitlements.
- This good practice statement was based on a review of 27 global policies for parental leave and entitlements for families of preterm and LBW Infants.

Evidence-to-Decision summary

Benefits	Large
Harms	None
Certainty	Unknown
Balance	Favours parental leave and entitlements
Values	No uncertainty or variability about outcomes
Acceptability	Acceptable
Resources	Moderate
Feasibility	Varies
Equity	Equitable

4. Implementation

The recommendations should be adapted to the needs of different countries, local contexts, and individual families and infants. The Guideline Development Group proposed implementation considerations for each recommendation and also reflected on adoption, adaptation and implementation to ensure availability, accessibility, acceptability and quality of care, in accordance with a human rights-based approach. Providers of services for the preterm or low-birth-weight (LBW) infant must consider the needs of, and provide equal care to, all individuals and their newborns.

Health policy considerations for the adoption and scale-up of recommended interventions for the care of the preterm or LBW infant:

- A firm government commitment to scale-up and increased coverage of these interventions is needed, irrespective of social, economic, ethnic, racial or other factors. National support must be secured for all recommendations, not just for specific components.
- To set the policy agenda, to secure broad anchoring and to ensure progress in policy formulation and decision-making, representatives of training facilities and the relevant medical specialties and professional societies should be included in participatory processes at all stages, including prior to an actual policy decision, to secure broad support for scaling up.
- To facilitate negotiations and planning, situation-specific information on the expected impact of the implementation of the recommendations on service users, health workers and costs should be compiled and disseminated.

Health system or organization-level considerations for implementation:

- Derivative tools should be updated, such as *Integrated management of childhood illness: management of the sick and young infant aged up to 2 months* (201), *Pocket book of hospital care for children* and *Guidelines for the management of common childhood illnesses* (202), and global and national essential medicines lists.
- National and subnational subgroups may be established to adapt and implement these recommendations, including the development

or revision of existing national or subnational guidelines or protocols.

- Long-term planning is needed for resource generation and budget allocation to address the shortage of health workers and trained community health workers, to improve facility infrastructure and referral pathways, and to strengthen and sustain high-quality small and sick newborn care services.
- Implementation of the recommendations should involve pre-service training institutions and professional bodies, so that training curricula for small and sick newborn care services can be updated as quickly and smoothly as possible.
- In-service training and supervisory courses will need to be developed according to health workers' professional requirements, considering the content and duration of the courses and the procedures for the selection of health workers for training. These courses can also be explicitly designed to address staff turnover, particularly in low-resource settings.
- Standardized tools will need to be developed for supervision, ensuring that supervisors are able to support and enable health workers to deliver integrated, comprehensive small and sick newborn care services.
- A strategy to optimize the use of human resources.
- Tools or "job aids" for implementation at the different levels of health-care facility and in communities will need to be developed or updated with all the key information.
- Strategies will need to be devised to improve supply chain management according to local requirements, such as developing protocols for the procedures of obtaining and maintaining the stock of supplies, encouraging health workers to collect and monitor data on the stock levels and strengthening the provider-level coordination and follow-up of medicines and health-care supplies required for implementation.
- Development or revision of national guidelines and/or health-care facility-based protocols is needed.
- Good-quality supervision, communication and transport links between community, primary- and higher-level facilities need to be established to ensure that referral pathways are efficient.

- Successful implementation strategies should be documented and shared as examples of best practice for other implementers.

User-level considerations for implementation:

- Community-sensitizing activities should be undertaken to disseminate information about the importance of each component of care, and infants' rights to receive care for their health and well-being. This information should provide details about the timing and content of the recommended contacts, and about the expected user fees.

Considerations for humanitarian emergencies:

- The adaptation of the recommendations should consider their integration and alignment with other response strategies. Additional considerations should be made for the unique needs of families and infants in emergency settings, including their values and preferences. Context-specific tools may be needed in addition to standard tools to support the implementation by stakeholders of the recommendations in humanitarian emergencies.

5. Applicability issues

A number of factors may hinder the effective implementation and scale-up of the recommendations in this guideline. These barriers may be related to the behaviours of families or health workers and to the organization of care or health service delivery. As part of efforts to implement these recommendations, health system stakeholders may wish to consider the following potential barriers:

- difficult access to health services and health workers for families and newborns, including lack of transport, geographical conditions and financial barriers;
- lack of human resources with the necessary expertise and skills to implement, supervise and support recommended practices, including client counselling;
- lack of infrastructure to support interventions (e.g. lack of electricity for refrigeration, lack of access to clean water and sanitation, lack of access to digital interventions and devices, lack of physical space to conduct individual care and counselling);
- lack of time or understanding of the value of newly recommended interventions among health workers and health system administrators;

- lack of physical resources (such as equipment, supplies, medicines and nutritional supplements);
- lack of opportunities for continuing education and professional development for health workers;
- resistance of health workers to change from non-evidence-based to evidence-based practices (such as providing home visits or ensuring family involvement);
- lack of effective referral mechanisms and care pathways for families and newborns identified as needing additional care (e.g. continuous positive airway pressure or methylxanthines);
- lack of health management information systems designed to document and monitor recommended practices (e.g. patient records and registers).

Given these potential barriers, a phased approach to the adoption, adaptation and implementation of the recommendations in this guideline may be helpful. Various strategies for addressing these barriers and facilitating implementation are provided in each chapter.

6. Research implications

The Guideline Development Group (GDG) identified important knowledge gaps that need to be addressed through primary research. The research questions were prioritized by the GDG based on consideration of whether they would: (i) contribute to improvements in care and outcomes for preterm or low-birth-weight (LBW) infants; (ii) be likely to result in significant public health impacts; (iii) be answerable; (iv) inform a new recommendation or change an existing recommendation; (v) result in findings that would be feasible to implement; and (vi) be likely to promote equity. The full list of research gaps can be found in Web Annex B, but the prioritized research questions are listed below.

A.1 Kangaroo mother care (KMC)

A.1a Any KMC

- What is the effectiveness of KMC on longer-term (i.e. up to 2 years of age, school-age, adolescence) growth, neurodevelopment, behaviour, mental health and disability outcomes?
- What are the key components of an implementation model that achieves high population-level coverage of KMC for more than 8 hours per day in high-income countries?

A.1b Immediate KMC

- What is the effectiveness of immediate KMC in critically ill preterm or LBW infants, such as infants who are mechanically ventilated or on blood pressure support (e.g. vasopressors)?
- How can immediate KMC be scaled up in routine health systems?

A.2 Mother's own milk

- How can exclusive breastfeeding be promoted, supported and scaled up for preterm or LBW infants, especially those who are very preterm or very LBW?
- What are the most effective early feeding strategies for very preterm or very LBW infants, infants with illnesses (e.g. post-surgery), and infants with other conditions (e.g. doppler abnormalities, severe growth restriction)?

A.3 Donor human milk

- What is the effectiveness, safety and feasibility of human milk banks in low- and middle-income countries?

A.11 Probiotics

- What is the effectiveness and safety of probiotics in human-milk-fed infants?
- What is the effect of probiotics on immune function and gut microbiome in preterm or LBW infants?
- What are the most optimal probiotic compositions for preterm or LBW infants – that is, the optimal combination of genera, species and strains?
- What is the optimal probiotics regimen (dosage and duration) for preterm or LBW infants?
- What is the effectiveness of probiotics alone compared with a combination of probiotics and prebiotics for preterm or LBW infants?
- What is the role of probiotics in the prevention and management of postnatal growth restriction in preterm infants?

A.12 Emollients

- What is the effect of emollients on mortality, invasive infection, sepsis, growth and longer-term neurodevelopment in preterm or LBW infants in high-, middle- and low-income countries, especially in Africa?
- What is the effect of emollients on thermoprotection and the microbiome in preterm or LBW infants?
- Which emollients (which oils, which composition) are most effective and safe for preterm or LBW infants?
- What is the optimal regime (dose, frequency, duration) and mode of application (e.g. non-touch applications) for very or extremely preterm infants?

B.1 Continuous positive airway pressure (CPAP) for respiratory distress syndrome

- What is the effectiveness of CPAP compared with humidified high-flow nasal cannulae and other forms of non-invasive ventilation in preterm or LBW infants with respiratory distress syndrome?

C.1 Family involvement

- What strategies can be used to increase family participation in the care of their preterm or LBW infants in intensive and special care units, and in settings without dedicated newborn units?

C.2 Family support

- What is the most effective type of family support (including education, counselling, discharge preparation, peer support) for families of preterm or LBW infants?

C.3 Home visits

- What is the effectiveness of standard in-person home visits compared with digital home visits (e.g. online video, mobile application [app], mHealth) for post-discharge follow-up of preterm or LBW infants?
- What is the feasibility of digital home visits in low-, middle- and high-income countries?

7. Dissemination

The recommendations will be disseminated through WHO regional and country offices, ministries of health, professional associations, WHO collaborating centres, other United Nations agencies and nongovernmental organizations. The recommendations will be available on the WHO website and also as a printed publication. Online versions will be available via the websites of the relevant WHO departments. Technical meetings will be held between WHO and stakeholders to share the recommendations and derivative products.

Evidence briefs for policy-makers, programme managers and health workers will be developed. They will focus on selected recommendations and context-specific issues, and will be developed and disseminated in collaboration with United Nations agencies and partners.

The executive summary and recommendations from this publication will be translated into the six United Nations languages for dissemination through the WHO regional and country offices, and web versions

will be available via the websites of the WHO departments, as above.

In addition, a number of articles presenting the recommendations and key implementation considerations will be published, in compliance with WHO's open access and copyright policies. Relevant WHO clusters, departments and partnerships, such as the Partnership for Maternal, Newborn and Child Health (PMNCH), will also be part of the dissemination process.

WHO, in collaboration with other partners, will support national and subnational working groups to adopt, adapt and implement the guideline. This will include the development or revision of existing national policies, guidelines or protocols in line with the WHO recommendations, and tools to support the adaptation and implementation processes. This also includes technical support for local guideline implementers in the development of training materials and quality indicators.

8. Monitoring and evaluating the impact of the recommendations

The implementation and impact of these recommendations will be monitored at the health service, subnational and national levels, based on clearly defined criteria and indicators that are associated with locally agreed targets. In collaboration with the monitoring and evaluation teams of the WHO Departments of Maternal, Newborn, Child and Adolescent Health and Ageing, and Sexual and Reproductive Health and Research, the data on country- and regional-level adoption of the recommendations will be collected and evaluated in the short to medium term across individual WHO Member States, through the WHO sexual, reproductive, maternal, newborn, child and adolescent health (SRMNCAH) policy survey (198). A full monitoring framework will be developed. In the meantime, the Guideline Development Group for this guideline suggests the consideration of the following indicators, which have been adapted from current global recommended indicators (53,203), including

the *Every Newborn Action Plan* (ENAP) indicators for mortality and coverage of postnatal care.

- Neonatal mortality – the proportion of preterm or low-birth-weight (LBW) infants dying in the first 28 days after birth.
- Early breastfeeding – the proportion of preterm or LBW infants put to the breast within the first 24 hours after birth.
- Early postnatal care for infants within two days of birth – the proportion of preterm or LBW infants who had postnatal contact with a health worker within two days of delivery.
- Kangaroo mother care (KMC) – the proportion of preterm infants who receive KMC.

These indicators should be considered preliminary and will undergo further review. New indicators will be added, including those for the measurement of coverage and quality of care.

9. Updating of the guideline

In accordance with the process for updating WHO guidelines, the “living guidelines” approach will be used (204). This is a systematic and continual process of identifying and bridging evidence gaps, at least every six months following guideline publication and dissemination. A Guideline Steering Group for maternal and newborn health recommendations will convene regularly to review WHO’s current portfolio of relevant recommendations, and to prioritize new and existing questions for recommendation development and updating. The focus will be on recommendations supported by very-low- or low-certainty evidence and where new recommendations or a change in the published recommendations may be needed. When new evidence that could potentially impact the current evidence base for any of the recommendations is identified, the recommendation

will be updated. If no new reports or information are identified for a particular recommendation, the recommendation will be revalidated.

Any concern about the validity of any recommendation should be promptly communicated by email to the WHO Department of Maternal, Newborn, Child and Adolescent Health and Ageing (mncah@who.int). All communications will be reviewed and plans will be made to update the recommendation as needed.

WHO welcomes suggestions regarding additional questions for inclusion in future updates of this guideline; suggestions can be addressed by email to the same department (mncah@who.int).

References

1. Blencowe H, Krusevec J, de Onis M, Black RE, An X, Stevens GA, et al. National, regional, and worldwide estimates of low birthweight in 2015, with trends from 2000: a systematic analysis. *Lancet Glob Health*. 2019;(7):e849-e60. doi:10.1016/S2214-109X(18)30565-5.
2. Katz J, Lee AC, Kozuki N, Lawn JE, Cousens S, Blencowe H, et al. Mortality risk in preterm and small-for-gestational-age infants in low-income and middle-income countries: a pooled country analysis. *Lancet*. 2013;382(9890):417-25. doi:10.1016/S0140-6736(13)60993-9.
3. Lee AC, Kozuki N, Cousens S, Stevens GA, Blencowe H, Silveira MF, et al. Estimates of burden and consequences of infants born small for gestational age in low and middle income countries with INTERGROWTH-21st standard: analysis of CHERG datasets. *BMJ*. 2017;358:j3677. doi:10.1136/bmj.j3677.
4. Global strategy for women's, children's and adolescents' health (2016–2030). New York (NY): Every Woman Every Child; 2015 (<http://globalstrategy.everywomaneverychild.org>).
5. Escobar GJ, McCormick MC, Zupancic JA, Coleman-Phox K, Armstrong MA, Greene JD, et al. Unstudied infants: outcomes of moderately premature infants in the neonatal intensive care unit. *Arch Dis Child Fetal Neonatal Ed*. 2006;91(4):F238-44. doi:10.1136/adc.2005.087031.
6. Wang ML, Dorer DJ, Fleming MP, Catlin EA. Clinical outcomes of near-term infants. *Pediatrics*. 2004;114(2):372-6. doi:10.1542/peds.114.2.372.
7. Kinney HC. The near-term (late preterm) human brain and risk for periventricular leukomalacia: a review. *Semin Perinatol*. 2006;30(2):81-8. doi:10.1053/j.semperi.2006.02.006.
8. Blencowe H, Lee AC, Cousens S, Bahalim A, Narwal R, Zhong N, et al. Preterm birth-associated neurodevelopmental impairment estimates at regional and global levels for 2010. *Pediatr Res*. 2013;74(Suppl 1):17-34. doi:10.1038/pr.2013.204.
9. Gluckman PD, Hanson MA, Beedle AS. Early life events and their consequences for later disease: a life history and evolutionary perspective. *Am J Hum Biol*. 2007;19(1):1-19. doi:10.1002/ajhb.20590.
10. World Health Organization, United Nations Children's Fund. Low birth weight: country, regional and global estimates. Geneva: World Health Organization; 2004 (<https://apps.who.int/iris/handle/10665/43184>).
11. Lawn JE, Cousens S, Zupan J. 4 million neonatal deaths: When? Where? Why? *Lancet*. 2005;365(9462):891-900. doi:10.1016/s0140-6736(05)71048-5.
12. Perin J, Mulick A, Yeung D, Villavicencio F, Lopez G, Strong KL, et al. Global, regional, and national causes of under-5 mortality in 2000-19: an updated systematic analysis with implications for the Sustainable Development Goals. *Lancet Child Adolesc Health*. 2022;6(2):106-15. doi:10.1016/S2352-4642(21)00311-4.
13. McCormick MC. The contribution of low birth weight to infant mortality and childhood morbidity. *N Engl J Med*. 1985;312(2):82-90. doi:10.1056/NEJM198501103120204.
14. Hurt L, Odd D, Mann M, Beetham H, Dorgeat E, Isaac TC, et al. What matters to families about the healthcare of preterm or low birth weight babies: a qualitative evidence synthesis. *medRxiv*. 2022:2022.06.22.22276770. doi:10.1101/2022.06.22.22276770.
15. Every child alive: the urgent need to end newborn deaths. New York (NY): United Nations Children's Fund; 2018 (<https://www.unicef.org/reports/every-child-alive>).

16. Sixty-fifth World Health Assembly, Geneva, 21-26 May 2012: resolutions and decisions: annexes. Geneva: World Health Organization; 2012 (<https://apps.who.int/iris/handle/10665/80058>).
17. Every newborn: an action plan to end preventable deaths. Geneva: World Health Organization; 2014 (<https://apps.who.int/iris/handle/10665/127938>).
18. Sixty-seventh World Health Assembly, Geneva, 19-24 May 2014: resolutions and decisions: annexes. Geneva: World Health Organization; 2014 (<https://apps.who.int/iris/handle/10665/260211>).
19. Guidelines on optimal feeding of low birth-weight infants in low- and middle-income countries. Geneva: World Health Organization; 2011 (<https://apps.who.int/iris/handle/10665/85670>).
20. WHO recommendations on interventions to improve preterm birth outcomes. Geneva: World Health Organization; 2015 (<https://apps.who.int/iris/handle/10665/183037>).
21. Recommendations for management of common childhood conditions: evidence for technical update of pocket book recommendations: newborn conditions, dysentery, pneumonia, oxygen use and delivery, common causes of fever, severe acute malnutrition and supportive care. Geneva: World Health Organization; 2012 (<https://apps.who.int/iris/handle/10665/44774>).
22. WHO recommendations on maternal and newborn care for a positive postnatal experience. Geneva: World Health Organization; 2022 (<https://apps.who.int/iris/handle/10665/352658>).
23. WHO recommendations on antenatal care for a positive pregnancy experience. Geneva: World Health Organization; 2016 (<https://apps.who.int/iris/handle/10665/250796>).
24. WHO recommendations: intrapartum care for a positive childbirth experience. Geneva: World Health Organization; 2018 (<https://apps.who.int/iris/handle/10665/260178>).
25. WHO handbook for guideline development, second edition. Geneva: World Health Organization; 2014 (<https://apps.who.int/iris/handle/10665/145714>).
26. Edmond K, Strobel N. Evidence for global health care interventions for preterm or low birth weight infants: an overview of systematic reviews. *Pediatrics*. 2022;150(Suppl 1). doi:10.1542/peds.2022-057092C.
27. Declarations of interest. In: About WHO - ethics. 2020 [website]. Geneva: World Health Organization; 2020 (<https://www.who.int/about/ethics/declarations-of-interest>, accessed 10 September 2022).
28. DECIDE 2011-2015 [website]. The Cochrane Collaboration; 2022 (<https://www.decide-collaboration.eu/>, accessed 1 July 2022).
29. Grading of Recommendations Assessment, Development and Evaluation [website]. GRADE Working Group; 2022 (<http://gradeworkinggroup.org/>, accessed 11 April 2022).
30. GRADE-CERQual project group [website]. GRADE-CERQual; 2018 (<https://www.cerqual.org/the-grade-cerqual-project-group/>, accessed 20 January 2022).
31. UNICEF supply catalogue [website]. New York (NY): United Nations Children's Fund; 2018 (<https://supply.unicef.org/>, accessed 31 September 2021).
32. International medical products price guide, 2015 edition. Medford (MA): Management Sciences for Health; 2016 (<https://mshpriceguide.org/en/home>).
33. WHO compendium of innovative health technologies for low-resource settings: 2022. Geneva: World Health Organization; 2022 (<https://apps.who.int/iris/handle/10665/355162>).
34. Reporting the effects of an intervention in EPOC reviews. Cochrane Effective Practice and Organisation of Care Group; 2018 (https://epoc.cochrane.org/sites/epoc.cochrane.org/files/public/uploads/Resources-for-authors2017/how_to_report_the_effects_of_an_intervention.pdf).

35. Prady SL, Uphoff EP, Power M, Golder S. Development and validation of a search filter to identify equity-focused studies: reducing the number needed to screen. *BMC Med Res Methodol*. 2018;18(1):106. doi:10.1186/s12874-018-0567-x.
36. Langlois EV, Miskurka M, Zunzunegui MV, Ghaffar A, Ziegler D, Karp I. Inequities in postnatal care in low- and middle-income countries: a systematic review and meta-analysis. *Bull World Health Organ*. 2015;93(4):259-70G. doi:10.2471/BLT.14.140996.
37. Anderzen-Carlsson A, Lamy Z, Tingvall M, Eriksson M. Parental experiences of providing skin-to-skin care to their newborn infant – Part 2: a qualitative meta-synthesis. *Int J Qual Stud Health Well-being*. 2014;9:24907. doi:10.3402/qhw.v9.24907.
38. Smith ER, Bergelson I, Constantian S, Valsangkar B, Chan GJ. Barriers and enablers of health system adoption of kangaroo mother care: a systematic review of caregiver perspectives. *BMC Pediatr*. 2017;17(1):35. doi:10.1186/s12887-016-0769-5.
39. WHO Immediate KMC Study Group, Arya S, Naburi H, Kawaza K, Newton S, Anyabolu CH, et al. Immediate “kangaroo mother care” and survival of infants with low birth weight. *N Engl J Med*. 2021;384(21):2028-38. doi:10.1056/NEJMoa2026486.
40. Mazumder S, Taneja S, Dube B, Bhatia K, Ghosh R, Shekhar M, et al. Effect of community-initiated kangaroo mother care on survival of infants with low birthweight: a randomised controlled trial. *Lancet*. 2019;394(10210):1724-36. doi:10.1016/S0140-6736(19)32223-8.
41. Sivanandan S, Sankar MJ. Kangaroo mother care for preterm or low birth weight infants: a systematic review and meta-analysis. *medRxiv*. 2022:2022.09.14.22279053. doi:10.1101/2022.09.14.22279053.
42. Blunt JW, DeLuca HF, Schnoes HK. 25-hydroxycholecalciferol: biologically active metabolite of vitamin D3. *Biochemistry*. 1968;7(10):3317-22. doi:10.1021/bi00850a001.
43. Mony PK, Tadele H, Gobezaeyehu AG, Chan GJ, Kumar A, Mazumder S, et al. Scaling up kangaroo mother care in Ethiopia and India: a multi-site implementation research study. *BMJ Glob Health*. 2021;6(9):e005905. doi:10.1136/bmjgh-2021-005905.
44. Klemming S, Lilliesköld S, Westrup B. Mother-Newborn Couplet Care from theory to practice to ensure zero separation for all newborns. *Acta Paediatrica*. 2021;110(11):2951-7. doi:10.1111/apa.15997.
45. Arora P, Kommalur A, Devadas S, Kariyappa M, Rao SPN. Quality improvement initiative to improve the duration of kangaroo mother care for twin preterm neonates born at a tertiary care hospital in resource-limited settings. *J Paediatr Child Health*. 2021;57(7):1082-8. doi:10.1111/jpc.15406.
46. Minot KL, Kramer KP, Butler C, Foster M, Gregory C, Haynes K, et al. Increasing early skin-to-skin in extremely low birth weight infants. *Neonatal Netw*. 2021;40(4):242-50. doi:10.1891/11-T-749.
47. Calibo AP, De Leon Mendoza S, Silvestre MA, Murray JCS, Li Z, Mannava P, et al. Scaling up kangaroo mother care in the Philippines using policy, regulatory and systems reform to drive changes in birth practices. *BMJ Global Health*. 2021;6(8):e006492. doi:10.1136/bmjgh-2021-006492.
48. Hendricks-Munoz KD, Mayers RM. A neonatal nurse training program in kangaroo mother care (KMC) decreases barriers to KMC utilization in the NICU. *Am J Perinatol*. 2014;31(11):987-92.
49. Kapoor R, Verma A, Dalal P, Gathwala G, Dalal J. Enhancing kangaroo mother care uptake through implementation of an education protocol. *Indian J Pediatr*. 2021;88(6):544-9. doi:10.1007/s12098-020-03537-z.
50. Mondkar J, Chawla D, Sachdeva RC, Manerkar S, Shanbhag S, Khan A, et al. Impact of mother-baby friendly initiative plus approach on improving human milk feeding for neonates in hospital: a quality improvement before-and-after uncontrolled study. *Eur J Pediatr*. 2022;181(1):107-16. doi:10.1007/s00431-021-04141-9.

51. Vesel L, ten Asbroek AH, Manu A, Soremekun S, Tawiah Agyemang C, Okyere E, et al. Promoting skin-to-skin care for low birthweight babies: findings from the Ghana Newhints cluster-randomised trial. *Trop Med Int Health*. 2013;18(8):952-61. doi:10.1111/tmi.12134.
52. Requejo J, Diaz T, Park L, Chou D, Choudhury A, Guthold R, et al. Assessing coverage of interventions for reproductive, maternal, newborn, child, and adolescent health and nutrition. *BMJ*. 2020;368:l6915. doi:10.1136/bmj.l6915.
53. Maternal and newborn – Coverage. In: Maternal, newborn, child and adolescent health and ageing: data portal [website]. Geneva: World Health Organization; 2022 (www.who.int/data/maternal-newborn-child-adolescent-ageing/maternal-and-newborn-data/maternal-and-newborn---coverage, accessed 11 April 2022).
54. Grummer-Strawn LM, Rollins N. Summarising the health effects of breastfeeding. *Acta Paediatr*. 2015;104(467):1-2. doi:10.1111/apa.13136.
55. Walsh V, McGuire W. Immunonutrition for Preterm Infants. *Neonatology*. 2019;115(4):398-405. doi:10.1159/000497332.
56. Optimal feeding of low-birth-weight infants: technical review. Geneva: World Health Organization; 2006 (<https://apps.who.int/iris/handle/10665/43602>).
57. Embleton NE, Pang N, Cooke RJ. Postnatal malnutrition and growth retardation: an inevitable consequence of current recommendations in preterm infants? *Pediatrics*. 2001;107(2):270-3. doi:10.1542/peds.107.2.270.
58. Embleton ND, Berrington JE, Dorling J, Ewer AK, Juszczak E, Kirby JA, et al. Mechanisms affecting the gut of preterm infants in enteral feeding trials. *Front Nutr*. 2017;4:14. doi:10.3389/fnut.2017.00014.
59. Gidrewicz DA, Fenton TR. A systematic review and meta-analysis of the nutrient content of preterm and term breast milk. *BMC Pediatr*. 2014;14(1):216. doi:10.1186/1471-2431-14-216.
60. Strobel NA, Adams C, McAullay DR, Edmond KM. Mother's own milk compared with formula milk for feeding preterm or low birth weight infants: systematic review and meta-analysis. *Pediatrics*. 2022;150(Suppl 1). doi:10.1542/peds.2022-057092D.
61. Mrelus E, Brogren S, Andersson S, Alehagen S. Fathers' experiences of feeding their extremely preterm infants in family-centred neonatal intensive care: a qualitative study. *Int Breastfeed J*. 2021;16(1):46. doi:10.1186/s13006-021-00394-0.
62. Russell G, Sawyer A, Rabe H, Abbott J, Gyte G, Duley L, et al. Parents' views on care of their very premature babies in neonatal intensive care units: a qualitative study. *BMC Pediatr*. 2014;14(1):230. doi:10.1186/1471-2431-14-230.
63. Guideline: protecting, promoting and supporting breastfeeding in facilities providing maternity and newborn services. Geneva: World Health Organization; 2017 (<https://apps.who.int/iris/handle/10665/259386>).
64. Baker P, Santos T, Neves PA, Machado P, Smith J, Piwoz E, et al. First-food systems transformations and the ultra-processing of infant and young child diets: the determinants, dynamics and consequences of the global rise in commercial milk formula consumption. *Matern Child Nutr*. 2021;17(2):e13097. doi:10.1111/mcn.13097.
65. Neves PAR, Vaz JS, Maia FS, Baker P, Gatica-Domínguez G, Piwoz E, et al. Rates and time trends in the consumption of breastmilk, formula, and animal milk by children younger than 2 years from 2000 to 2019: analysis of 113 countries. *Lancet Child Adolesc Health*. 2021;5(9):619-30. doi:10.1016/s2352-4642(21)00163-2.
66. Tyebally Fang M, Grummer-Strawn L, Maryuningsih Y, Biller-Andorno N. Human milk banks: a need for further evidence and guidance. *Lancet Glob Health*. 2021;9(2):e104-e5. doi:10.1016/S2214-109X(20)30468-X.
67. Quigley M, Embleton ND, McGuire W. Formula versus donor breast milk for feeding preterm or low birth weight infants. *Cochrane Database Syst Rev*. 2019;(7):CD002971.

68. Chagwena DT, Mugariri F, Sithole B, Mataga SF, Danda R, Matsungu TM, et al. Acceptability of donor breastmilk banking among health workers: a cross-sectional survey in Zimbabwean urban settings. *Int Breastfeed J*. 2020;15(1):37. doi:10.1186/s13006-020-00283-y.
69. Esquerre-Zwiers A, Rossman B, Meier P, Engstrom J, Janes J, Patel A. "It's somebody else's milk": unraveling the tension in mothers of preterm infants who provide consent for pasteurized donor human milk. *J Hum Lact*. 2016;32(1):95-102. doi:10.1177/0890334415617939.
70. Lubbe W, Oosthuizen CS, Dolman RC, Covic N. Stakeholder attitudes towards donating and utilizing donated human breastmilk. *Int J Environ Res Public Health*. 2019;16(10):1838. doi:10.3390/ijerph16101838.
71. Magowan S, Burgoine K, Ogara C, Ditai J, Gladstone M. Exploring the barriers and facilitators to the acceptability of donor human milk in eastern Uganda – a qualitative study. *Int Breastfeed J*. 2020;15(1):28. doi:10.1186/s13006-020-00272-1.
72. Tyebally Fang M, Chatzixiros E, Grummer-Strawn L, Engmann C, Israel-Ballard K, Mansen K, et al. Developing global guidance on human milk banking. *Bull World Health Organ*. 2021;99(12):892-900. doi:10.2471/BLT.21.286943.
73. Rochow N, Fusch G, Choi A, Chessell L, Elliott L, McDonald K, et al. Target fortification of breast milk with fat, protein, and carbohydrates for preterm infants. *J Pediatr*. 2013;163(4):1001-7. doi:10.1016/j.jpeds.2013.04.052.
74. Agostoni C, Buonocore G, Carnielli V, De Curtis M, Darmaun D, Decsi T, et al. Enteral nutrient supply for preterm infants: commentary from the European Society of Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. *J Pediatr Gastroenterol Nutr*. 2010;50(1):85-91. doi:10.1097/MPG.0b013e3181adaee0.
75. Arslanoglu S, Moro GE, Ziegler EE, World Association of Perinatal Medicine Working Group On Nutrition. Optimization of human milk fortification for preterm infants: new concepts and recommendations. *J Perinat Med*. 2010;38(3):233-8. doi:10.1515/jpm.2010.073.
76. Imdad A, Bhutta ZA. Nutritional management of the low birth weight/preterm infant in community settings: a perspective from the developing world. *J Pediatr*. 2013;162(Suppl 3):S107-14. doi:10.1016/j.jpeds.2012.11.060.
77. Klingenberg C, Embleton ND, Jacobs SE, O'Connell LA, Kuschel CA. Enteral feeding practices in very preterm infants: an international survey. *Arch Dis Child Fetal Neonatal Ed*. 2012;97(1):F56-61. doi:10.1136/adc.2010.204123.
78. Brown JV, Lin L, Embleton ND, Harding JE, McGuire W. Multi-nutrient fortification of human milk for preterm infants. *Cochrane Database Syst Rev*. 2020;6(7):CD000343.
79. Hay WW, Jr., Hendrickson KC. Preterm formula use in the preterm very low birth weight infant. *Semin Fetal Neonatal Med*. 2017;22(1):15-22. doi:10.1016/j.siny.2016.08.005.
80. Horbar JD, Ehrenkranz RA, Badger GJ, Edwards EM, Morrow KA, Soll RF, et al. Weight growth velocity and postnatal growth failure in infants 501 to 1500 grams: 2000-2013. *Pediatrics*. 2015;136(1):e84-92. doi:10.1542/peds.2015-0129.
81. Walsh V, Brown JVE, Askie LM, Embleton ND, McGuire W. Nutrient-enriched formula versus standard formula for preterm infants. *Cochrane Database Syst Rev*. 2019;(7):CD004204.
82. Pollitt E, Gilmore M, Valcarcel M. The stability of sucking behavior and its relationship to intake during the first month of life. *Infant Behav Dev*. 1978;1:347-57. doi:10.1016/s0163-6383(78)80046-0.
83. Shulhan J, Dicken B, Hartling L, Larsen BM. Current knowledge of necrotizing enterocolitis in preterm infants and the impact of different types of enteral nutrition products. *Adv Nutr*. 2017;8(1):80-91. doi:10.3945/an.116.013193.

84. Fang L, Wu L, Han S, Chen X, Yu Z. Quality improvement to increase breastfeeding in preterm infants: systematic review and meta-analysis. *Front Pediatr*. 2021;9:681341. doi:10.3389/fped.2021.681341.
85. Kinshella MW, Prasad S, Hiwa T, Vidler M, Nyondo-Mipando AL, Dube Q, et al. Barriers and facilitators for early and exclusive breastfeeding in health facilities in Sub-Saharan Africa: a systematic review. *Glob Health Res Policy*. 2021;6(1):21. doi:10.1186/s41256-021-00206-2.
86. Chitale R, Ferguson K, Talej M, Yang WC, He S, Edmond KM, et al. Early enteral feeding for preterm or low birth weight infants: a systematic review and meta-analysis. *Pediatrics*. 2022;150(Suppl 1). doi:10.1542/peds.2022-057092E.
87. Black MM, Aboud FE. Responsive feeding is embedded in a theoretical framework of responsive parenting. *J Nutr*. 2011;141(3):490-4. doi:10.3945/jn.110.129973.
88. Glodowski KR, Thompson RH, Martel L. The rooting reflex as an infant feeding cue. *J Appl Behav Anal*. 2019;52(1):17-27. doi:10.1002/jaba.512.
89. McCormick FM, Tosh K, McGuire W. Ad libitum or demand/semi-demand feeding versus scheduled interval feeding for preterm infants. *Cochrane Database Syst Rev*. 2010;(2):CD005255.
90. Watson J, McGuire W. Responsive versus scheduled feeding for preterm infants. *Cochrane Database Syst Rev*. 2016;(8):CD005255.
91. Fry TJ, Marfurt S, Wengier S. Systematic review of quality improvement initiatives related to cue-based feeding in preterm infants. *Nurs Womens Health*. 2018;22(5):401-10. doi:10.1016/j.nwh.2018.07.006.
92. Talej M, Smith ER, Lauria ME, Chitale R, Ferguson K, He S. Responsive feeding for preterm or low birth weight infants: a systematic review and meta-analysis. *Pediatrics*. 2022;150(Suppl 1). doi:10.1542/peds.2022-057092F.
93. Brown EG, Sweet AY. Preventing necrotizing enterocolitis in neonates. *JAMA*. 1978;240(22):2452-4. doi:10.1001/jama.1978.03290220064019.
94. McKeown RE, Marsh TD, Amarnath U, Garrison CZ, Addy CL, Thompson SJ, et al. Role of delayed feeding and of feeding increments in necrotizing enterocolitis. *J Pediatr*. 1992;121(5 Pt 1):764-70. doi:10.1016/s0022-3476(05)81913-4.
95. Oddie SJ, Young L, McGuire W. Slow advancement of enteral feed volumes to prevent necrotising enterocolitis in very low birth weight infants. *Cochrane Database Syst Rev*. 2021;(8):CD001241.
96. Yang WC, Fogel A, Lauria ME, Ferguson K, Smith ER. Fast feed advancement for preterm and low birth weight infants: a systematic review and meta-analysis. *Pediatrics*. 2022;150(Suppl 1). doi:10.1542/peds.2022-057092G.
97. Dorling J, Hewer O, Hurd M, Bari V, Bosiak B, Bowler U, et al. Two speeds of increasing milk feeds for very preterm or very low-birthweight infants: the SIFT RCT. *Health Technol Assess*. 2020;24(18):1-94. doi:10.3310/hta24180.
98. Victora CG, Christian P, Vidaletti LP, Gatica-Dominguez G, Menon P, Black RE. Revisiting maternal and child undernutrition in low-income and middle-income countries: variable progress towards an unfinished agenda. *Lancet*. 2021;397(10282):1388-99. doi:10.1016/S0140-6736(21)00394-9.
99. Yang WC, Lauria ME, Fogel A, Ferguson K, Smith ER. Duration of exclusive breastfeeding for preterm or low birth weight infants: a systematic review and meta-analysis. *Pediatrics*. 2022;150(Suppl 1). doi:10.1542/peds.2022-057092H.
100. McCarthy EK, Dempsey EM, Kiely ME. Iron supplementation in preterm and low-birth-weight infants: a systematic review of intervention studies. *Nutr Rev*. 2019;77(12):865-77. doi:10.1093/nutrit/nuz051.
101. Mills RJ, Davies MW. Enteral iron supplementation in preterm and low birth weight infants. *Cochrane Database Syst Rev*. 2012;(3):CD005095.

102. Manapurath RM, Gadapani Pathak B, Sinha B, Upadhyay RP, Choudhary TS, Chandola TR, et al. Enteral iron supplementation in preterm or low birth weight infants: a systematic review and meta-analysis. *Pediatrics*. 2022;150(Suppl 1). doi:10.1542/peds.2022-057092I.
103. Wessells KR, Brown KH. Estimating the global prevalence of zinc deficiency: results based on zinc availability in national food supplies and the prevalence of stunting. *PLoS One*. 2012;7(11):e50568. doi:10.1371/journal.pone.0050568.
104. Hambidge M. Human zinc deficiency. *J Nutr*. 2000;130(5S Suppl):1344S-9S. doi:10.1093/jn/130.5.1344S.
105. King JC, Shames DM, Woodhouse LR. Zinc homeostasis in humans. *J Nutr*. 2000;130(5S Suppl):1360S-6S. doi:10.1093/jn/130.5.1360S.
106. Gulani A, Bhatnagar S, Sachdev HP. Neonatal zinc supplementation for prevention of mortality and morbidity in breastfed low birth weight infants: systematic review of randomized controlled trials. *Indian Pediatr*. 2011;48(2):111-7. doi:10.1007/s13312-011-0043-8.
107. Staub E, Evers K, Askie LM. Enteral zinc supplementation for prevention of morbidity and mortality in preterm neonates. *Cochrane Database Syst Rev*. 2021;(3):CD012797.
108. Sinha B, Dudeja N, Chowdhury R, Choudhary TS, Upadhyay RP, Rongsen-Chandola T, et al. Enteral zinc supplementation in preterm or low birth weight infants: a systematic review and meta-analysis. *Pediatrics*. 2022;150(Suppl 1). doi:10.1542/peds.2022-057092J.
109. Huey SL, Acharya N, Silver A, Shen R, Yu EA, Pena-Rosas JP, et al. Effects of oral vitamin D supplementation on linear growth and other health outcomes among children under five years of age. *Cochrane Database Syst Rev*. 2020;(12):CD012875.
110. Misra M, Pacaud D, Petryk A, Collett-Solberg PF, Kappy M, Drug, et al. Vitamin D deficiency in children and its management: review of current knowledge and recommendations. *Pediatrics*. 2008;122(2):398-417. doi:10.1542/peds.2007-1894.
111. Tan ML, Abrams SA, Osborn DA. Vitamin D supplementation for term breastfed infants to prevent vitamin D deficiency and improve bone health. *Cochrane Database Syst Rev*. 2020;(12):CD013046.
112. Specker BL, Tsang RC, Hollis BW. Effect of race and diet on human-milk vitamin D and 25-hydroxyvitamin D. *Am J Dis Child*. 1985;139(11):1134-7. doi:10.1001/archpedi.1985.02140130072032.
113. Zittermann A, Pilz S, Berthold HK. Serum 25-hydroxyvitamin D response to vitamin D supplementation in infants: a systematic review and meta-analysis of clinical intervention trials. *Eur J Nutr*. 2020;59(1):359-69. doi:10.1007/s00394-019-01912-x.
114. Kumar M, Shaikh S, Sinha B, Upadhyay RP, Choudhary TS, Chandola TR, et al. Enteral vitamin D supplementation in preterm or low birth weight infants: a systematic review and meta-analysis. *Pediatrics*. 2022;150(Suppl 1). doi:10.1542/peds.2022-057092K.
115. Kumar GT, Sachdev HS, Chellani H, Rehman AM, Singh V, Arora H, et al. Effect of weekly vitamin D supplements on mortality, morbidity, and growth of low birthweight term infants in India up to age 6 months: randomised controlled trial. *BMJ*. 2011;342:d2975. doi:10.1136/bmj.d2975.
116. Brown CC, Noelle RJ. Seeing through the dark: new insights into the immune regulatory functions of vitamin A. *Eur J Immunol*. 2015;45(5):1287-95. doi:10.1002/eji.201344398.
117. Shenai JP. Vitamin A supplementation in very low birth weight neonates: rationale and evidence. *Pediatrics*. 1999;104(6):1369-74. doi:10.1542/peds.104.6.1369.
118. Khwaja O, Volpe JJ. Pathogenesis of cerebral white matter injury of prematurity. *Arch Dis Child Fetal Neonatal Ed*. 2008;93(2):F153-61. doi:10.1136/adc.2006.108837.

119. Lee J, Dammann O. Perinatal infection, inflammation, and retinopathy of prematurity. *Semin Fetal Neonatal Med.* 2012;17(1):26-9. doi:10.1016/j.siny.2011.08.007.
120. Darlow BA, Graham PJ, Rojas-Reyes MX. Vitamin A supplementation to prevent mortality and short- and long-term morbidity in very low birth weight infants. *Cochrane Database Syst Rev.* 2016;(8):CD000501.
121. Rakshasbhuwankar AA, Pillow JJ, Simmer KN, Patole SK. Vitamin A supplementation in very-preterm or very-low-birth-weight infants to prevent morbidity and mortality: a systematic review and meta-analysis of randomized trials. *Am J Clin Nutr.* 2021;114(6):2084-96. doi:10.1093/ajcn/nqab294.
122. Manapurath RM, Kumar M, Pathak BG, Chowdhury R, Sinha B, Choudhary T, et al. Enteral low-dose vitamin A supplementation in preterm or low birth weight infants to prevent morbidity and mortality: a systematic review and meta-analysis. *Pediatrics.* 2022;150(Suppl 1):e2022057092L. doi:10.1542/peds.2022-057092L.
123. Atkinson SA, Radde IC, Anderson GH. Macromineral balances in premature infants fed their own mothers' milk or formula. *J Pediatr.* 1983;102(1):99-106. doi:10.1016/s0022-3476(83)80302-3.
124. Harding JE, Wilson J, Brown J. Calcium and phosphorus supplementation of human milk for preterm infants. *Cochrane Database Syst Rev.* 2017;(2):CD003310.
125. Kumar M, Chowdhury R, Sinha B, Upadhyay RP, Chandola TR, Mazumder S, et al. Enteral calcium or phosphorus supplementation in preterm or low birth weight infants: a systematic review and meta-analysis. *Pediatrics.* 2022;150(Suppl 1). doi:10.1542/peds.2022-057092M.
126. Kumar M, Chowdhury R, Sinha B, Upadhyay RP, Chandola TR, Mazumder S, et al. Enteral multiple micronutrient supplementation in preterm and low birth weight infants: a systematic review and meta-analysis. *Pediatrics.* 2022;150(Suppl 1). doi:10.1542/peds.2022-057092N.
127. Sharif S, Meader N, Oddie SJ, Rojas-Reyes MX, McGuire W. Probiotics to prevent necrotising enterocolitis in very preterm or very low birth weight infants. *Cochrane Database Syst Rev.* 2020;(10):CD005496.
128. van den Akker CHP, van Goudoever JB, Szajewska H, Embleton ND, Hojsak I, Reid D, et al. Probiotics for preterm infants: a strain-specific systematic review and network meta-analysis. *J Pediatr Gastroenterol Nutr* 2018;67(1):103-22. doi:10.1097/MPG.0000000000001897.
129. Embleton ND, Zalewski S, Berrington JE. Probiotics for prevention of necrotizing enterocolitis and sepsis in preterm infants. *Curr Opin Infect Dis.* 2016;29(3):256-61. doi:10.1097/QCO.0000000000000269.
130. Thomas DW, Greer FR, American Academy of Pediatrics Committee on Nutrition, American Academy of Pediatrics Section on Gastroenterology Hepatology and Nutrition. Probiotics and prebiotics in pediatrics. *Pediatrics.* 2010;126(6):1217-31. doi:10.1542/peds.2010-2548.
131. Bron PA, Kleerebezem M, Brummer RJ, Cani PD, Mercenier A, MacDonald TT, et al. Can probiotics modulate human disease by impacting intestinal barrier function? *Br J Nutr.* 2017;117(1):93-107. doi:10.1017/S0007114516004037.
132. Koretz RL. Probiotics in gastroenterology: how pro is the evidence in adults? *Am J Gastroenterol.* 2018;113(8):1125-36. doi:10.1038/s41395-018-0138-0.
133. Lerner A, Shoenfeld Y, Matthias T. Probiotics: if it does not help it does not do any harm. Really? *Microorganisms.* 2019;7(4):104. doi:10.3390/microorganisms7040104.
134. Bertelli C, Pillonel T, Torregrossa A, Prod'homme G, Fischer CJ, Greub G, et al. *Bifidobacterium longum* bacteremia in preterm infants receiving probiotics. *Clin Infect Dis.* 2015;60(6):924-7. doi:10.1093/cid/ciu946.

135. Esaiassen E, Cavanagh P, Hjerde E, Simonsen GS, Stoen R, Klingenberg C. Bifidobacterium longum subspecies infantis bacteremia in 3 extremely preterm infants receiving probiotics. *Emerg Infect Dis*. 2016;22(9):1664-6. doi:10.3201/eid2209.160033.
136. Jenke A, Ruf EM, Hoppe T, Heldmann M, Wirth S. Bifidobacterium septicaemia in an extremely low-birthweight infant under probiotic therapy. *Arch Dis Child Fetal Neonatal Ed*. 2012;97(3):F217-8. doi:10.1136/archdischild-2011-300838.
137. Zbinden A, Zbinden R, Berger C, Arlettaz R. Case series of Bifidobacterium longum bacteremia in three preterm infants on probiotic therapy. *Neonatology*. 2015;107(1):56-9. doi:10.1159/000367985.
138. Alcon-Giner C, Dalby M, Caim S, Ketskemety J, Shaw A, Sim K, et al. Microbiota supplementation with bifidobacterium and lactobacillus modifies the preterm infant gut microbiota and metabolome: an observational study. *Cell Rep Med*. 2020;1(5):100077. doi:10.1016/j.xcrm.2020.100077.
139. Boyle RJ, Robins-Browne RM, Tang ML. Probiotic use in clinical practice: what are the risks? *Am J Clin Nutr*. 2006;83(6):1256-64. doi:10.1093/ajcn/83.6.1256.
140. Sesham R, Oddie S, Embleton ND, Clarke P. Probiotics for preterm neonates: parents' perspectives and present prevalence. *Arch Dis Child Fetal Neonatal Ed*. 2014;99(4):F345. doi:10.1136/archdischild-2014-306344.
141. Dyer JA. Newborn skin care. *Semin Perinatol*. 2013;37(1):3-7. doi:10.1053/j.semperi.2012.11.008.
142. Evans NJ, Rutter N. Development of the epidermis in the newborn. *Biol Neonate*. 1986;49(2):74-80. doi:10.1159/000242513.
143. Collins A, Weitkamp JH, Wynn JL. Why are preterm newborns at increased risk of infection? *Arch Dis Child Fetal Neonatal Ed*. 2018;103(4):F391-F4. doi:10.1136/archdischild-2017-313595.
144. Cleminson J, McGuire W. Topical emollient for preventing infection in preterm infants. *Cochrane Database Syst Rev*. 2021;(5):CD001150.
145. Darmstadt GL, Badrawi N, Law PA, Ahmed S, Bashir M, Iskander I, et al. Topically applied sunflower seed oil prevents invasive bacterial infections in preterm infants in Egypt: a randomized, controlled clinical trial. *Pediatr Infect Dis J*. 2004;23(8):719-25. doi:10.1097/01.inf.0000133047.50836.6f.
146. Fraser J, Walls M, McGuire W. ABC of preterm birth: respiratory complications of preterm birth. *BMJ*. 2004;329(7472):962-5. doi:10.1136/sbmj.0507278
147. Stoll BJ, Hansen NI, Bell EF, Walsh MC, Carlo WA, Shankaran S, et al. Trends in care practices, morbidity, and mortality of extremely preterm neonates, 1993-2012. *JAMA*. 2015;314(10):1039-51. doi:10.1001/jama.2015.10244.
148. Sweet DG, Carnielli V, Greisen G, Hallman M, Ozek E, Te Pas A, et al. European consensus guidelines on the management of respiratory distress syndrome - 2019 update. *Neonatology*. 2019;115(4):432-50. doi:10.1159/000499361.
149. Gregory GA, Kitterman JA, Phibbs RH, Tooley WH, Hamilton WK. Treatment of the idiopathic respiratory-distress syndrome with continuous positive airway pressure. *N Engl J Med*. 1971;284(24):1333-40. doi:10.1056/NEJM197106172842401.
150. Soll RF, Barkhuff W. Noninvasive ventilation in the age of surfactant administration. *Clin Perinatol*. 2019;46(3):493-516. doi:10.1016/j.clp.2019.05.002.
151. Ho JJ, Subramaniam P, Davis PG. Continuous positive airway pressure (CPAP) for respiratory distress in preterm infants. *Cochrane Database Syst Rev*. 2020;(10):CD002271.
152. Ho JJ, Subramaniam P, Sivakaanathan A, Davis PG. Early versus delayed continuous positive airway pressure (CPAP) for respiratory distress in preterm infants. *Cochrane Database Syst Rev*. 2020;(10):CD002975.

153. Gerad P, Fox WW, Outerbridge EW, Beaudry PH, Stern L. Early versus late introduction of continuous negative pressure in the management of the idiopathic respiratory distress syndrome. *J Pediatr*. 1975;87(4):591-5. doi:10.1016/s0022-3476(75)80832-8.
154. Mockrin LD, Bancalari EH. Early versus delayed initiation of continuous negative pressure in infants with hyaline membrane disease. *J Pediatr*. 1975;87(4):596-600. doi:10.1016/s0022-3476(75)80833-x.
155. Ho JJ, Henderson-Smart DJ, Davis PG. Early versus delayed initiation of continuous distending pressure for respiratory distress syndrome in preterm infants. *Cochrane Database Syst Rev*. 2002;(2):CD002975.
156. Badiie Z, Naseri F, Sadeghnia A. Early versus delayed initiation of nasal continuous positive airway pressure for treatment of respiratory distress syndrome in premature newborns: a randomized clinical trial. *Adv Biomed Res*. 2013;2:4. doi:10.4103/2277-9175.107965.
157. Morley CJ, Davis PG, Doyle LW, Brion LP, Hascoet JM, Carlin JB, et al. Nasal CPAP or intubation at birth for very preterm infants. *N Engl J Med*. 2008;358(7):700-8. doi:10.1056/NEJMoa072788.
158. Subramaniam P, Ho JJ, Davis PG. Prophylactic or very early initiation of continuous positive airway pressure (CPAP) for preterm infants. *Cochrane Database Syst Rev*. 2021;(10):CD001243.
159. Dada S, Ashworth H, Sobitschka A, Raguveer V, Sharma R, Hamilton RL, et al. Experiences with implementation of continuous positive airway pressure for neonates and infants in low-resource settings: a scoping review. *PLoS One*. 2021;16(6):e0252718. doi:10.1371/journal.pone.0252718.
160. Benveniste D, Berg O, Pedersen JE. A technique for delivery of continuous positive airway pressure to the neonate. *J Pediatr*. 1976;88(6):1015-9. doi:10.1016/s0022-3476(76)81066-9.
161. Gupta S, Donn SM. Continuous positive airway pressure: physiology and comparison of devices. *Semin Fetal Neonatal Med*. 2016;21(3):204-11. doi:10.1016/j.siny.2016.02.009.
162. Prakash R, De Paoli AG, Davis PG, Oddie SJ, McGuire W. Bubble devices versus other pressure sources for nasal continuous positive airway pressure in preterm infants. *Cochrane Database of Syst Rev*. 2022 (in press).
163. Kinshella MW, Walker CR, Hiwa T, Vidler M, Nyondo-Mipando AL, Dube Q, et al. Barriers and facilitators to implementing bubble CPAP to improve neonatal health in sub-Saharan Africa: a systematic review. *Public Health Rev*. 2020;41(1):6. doi:10.1186/s40985-020-00124-7.
164. Salimu S, Kinshella MW, Vidler M, Banda M, Newberry L, Dube Q, et al. Health workers' views on factors affecting caregiver engagement with bubble CPAP. *BMC Pediatr*. 2020;20(1):180. doi:10.1186/s12887-020-02088-8.
165. Aneji C, Hartman T, Olutunde O, Okonkwo I, Ewumwen E, Adetiloye O, et al. Implementing bubble continuous positive airway pressure in a lower middle-income country: a Nigerian experience. *Pan Afr Med J*. 2020;37:10. doi:10.11604/pamj.2020.37.10.24911.
166. Ekhuagere OA, Mairami AB, Kirpalani H. Risk and benefits of bubble continuous positive airway pressure for neonatal and childhood respiratory diseases in low- and middle-income countries. *Paediatr Respir Rev*. 2019;29:31-6. doi:10.1016/j.prrv.2018.04.004.
167. McAdams RM, Hedstrom AB, DiBlasi RM, Mant JE, Nyonyintono J, Otai CD, et al. Implementation of bubble CPAP in a rural Ugandan neonatal ICU. *Respir Care*. 2015;60(3):437-45. doi:10.4187/respcare.03438.
168. Nyondo-Mipando AL, Woo Kinshella ML, Bohne C, Suwedi-Kapesa LC, Salimu S, Banda M, et al. Barriers and enablers of implementing bubble continuous positive airway pressure (CPAP): perspectives of health professionals in Malawi. *PLoS One*. 2020;15(2):e0228915. doi:10.1371/journal.pone.0228915.

169. Schmidt B, Roberts RS, Davis P, Doyle LW, Barrington KJ, Ohlsson A, et al. Caffeine therapy for apnea of prematurity. *N Engl J Med*. 2006;354(20):2112-21. doi:10.1056/NEJMoa054065.
170. Eichenwald EC, Aina A, Stark AR. Apnea frequently persists beyond term gestation in infants delivered at 24 to 28 weeks. *Pediatrics*. 1997;100(3 Pt 1):354-9. doi:10.1542/peds.100.3.354.
171. Henderson-Smart DJ. Recurrent apnea in the newborn (Chapter 47). In: Evidence-based paediatrics, second edition. Moyer VA, Elliott EJ, Gilbert R, Klassen T, Logan S, Mellis C, et al., editors. BMJ Books; 2004:483-9.
172. Di Fiore JM, Bloom JN, Orge F, Schutt A, Schluchter M, Cheruvu VK, et al. A higher incidence of intermittent hypoxemic episodes is associated with severe retinopathy of prematurity. *J Pediatr*. 2010;157(1):69-73. doi:10.1016/j.jpeds.2010.01.046
173. Poets CF, Roberts RS, Schmidt B, Whyte RK, Asztalos EV, Bader D, et al. Association between intermittent hypoxemia or bradycardia and late death or disability in extremely preterm infants. *JAMA*. 2015;314(6):595-603. doi:10.1001/jama.2015.8841.
174. Marques K, Roehr CC, Bruschetti M, Davis PG, Soll R. Methylxanthine for the prevention and treatment of apnea in preterm infants. *Cochrane Database Syst Rev*. 2022 (in press).
175. Steer PA, Flenady VJ, Shearman A, Lee TC, Tudehope DI, Charles BG. Periextubation caffeine in preterm neonates: a randomized dose response trial. *J Paediatr Child Health*. 2003;39(7):511-5. doi:10.1046/j.1440-1754.2003.00207.x.
176. Dukhovny D, Lorch SA, Schmidt B, Doyle LW, Kok JH, Roberts RS, et al. Economic evaluation of caffeine for apnea of prematurity. *Pediatrics*. 2011;127(1):e146-55. doi:10.1542/peds.2010-1014.
177. Armanian AM, Badiie Z, Afghari R, Salehimehr N, Hassanzade A, Sheikhzadeh S, et al. Prophylactic aminophylline for prevention of apnea at higher-risk preterm neonates. *Iran Red Crescent Med J*. 2014;16(8):e12559. doi:10.5812/ircmj.12559.
178. Armanian AM, Iranpour R, Faghihian E, Salehimehr N. Caffeine administration to prevent apnea in very premature infants. *Pediatr Neonatol*. 2016;57(5):408-12. doi:10.1016/j.pedneo.2015.10.007.
179. Hegyi T, Hiatt IM, Stile IL, Zolfaghari S. Effects of postnatal aminophylline on the course of respiratory distress syndrome in premature infants. *Clin Ther*. 1986;8(4):439-49.
180. Kumar H, Bhat A, Alwadhi V, Maria A, Khanna R, Neogi SB, et al. An assessment of implementation of family participatory care in special newborn care units in three states of India. *Indian Pediatr*. 2021;58(4):349-53. doi:10.1007/s13312-021-2194-6.
181. Bastani F, Abadi TA, Haghani H. Effect of family-centered care on improving parental satisfaction and reducing readmission among premature infants: a randomized controlled trial. *J Clin Diagn Res*. 2015;9(1):SC04-8. doi:10.7860/JCDR/2015/10356.5444.
182. Committee on Hospital Care, American Academy of Pediatrics. Family-centered care and the pediatrician's role. *Pediatrics*. 2003;112(3 Part 1):691-7. doi:10.1542/peds.112.3.691.
183. Maria A, Litch JA, Stepanchak M, Sarin E, Wadhwa R, Kumar H. Assessment of feasibility and acceptability of family-centered care implemented at a neonatal intensive care unit in India. *BMC Pediatr*. 2021;21(1):171. doi:10.1186/s12887-021-02644-w.
184. Obeidat HM, Bond EA, Callister LC. The parental experience of having an infant in the newborn intensive care unit. *J Perinat Educ*. 2009;18(3):23-9 doi:10.1624/105812409X461199.
185. Ding X, Zhu L, Zhang R, Wang L, Wang TT, Latour JM. Effects of family-centred care interventions on preterm infants and parents in neonatal intensive care units: a systematic review and meta-analysis of randomised controlled trials. *Aust Crit Care*. 2019;32(1):63-75. doi:10.1016/j.aucc.2018.10.007.

186. Zhang R, Huang RW, Gao XR, Peng XM, Zhu LH, Rangasamy R, et al. Involvement of parents in the care of preterm infants: a pilot study evaluating a family-centered care intervention in a Chinese neonatal ICU. *Pediatr Crit Care Med*. 2018;19(8):741-7. doi:10.1097/PCC.0000000000001586.
187. North K, Whelan R, Folger LV, Lawford H, Olson I, Driker S, et al. Family involvement in the routine care of hospitalized preterm or low birth weight infants: a systematic review and meta-analysis. *Pediatrics*. 2022;150(Suppl 1). doi:10.1542/peds.2022-0570920.
188. Garti I, Donkor E, Musah N, Appiah EO, Gyekye S, Menlah A, et al. Mothers' experiences of caring for preterm babies at home: qualitative insights from an urban setting in a middle-income country. *BMC Pregnancy Childbirth*. 2021;21(1):395. doi:10.1186/s12884-021-03872-9.
189. Lakshmanan A, Kubicek K, Williams R, Robles M, Vanderbilt DL, Mirzaian CB, et al. Viewpoints from families for improving transition from NICU-to-home for infants with medical complexity at a safety net hospital: a qualitative study. *BMC Pediatr*. 2019;19(1):223. doi:10.1186/s12887-019-1604-6.
190. de Savigny D, Adam T, Alliance for Health Policy and Systems Research, World Health Organization. Systems thinking for health systems strengthening. Geneva: World Health Organization; 2009 (<https://apps.who.int/iris/handle/10665/44204>).
191. Hunt H, Abbott R, Boddy K, Whear R, Wakely L, Bethel A, et al. "They've walked the walk": a systematic review of quantitative and qualitative evidence for parent-to-parent support for parents of babies in neonatal care. *J Neonatal Nurs*. 2019;25(4):166-76. doi:10.1016/j.jnn.2019.03.011.
192. Labrie NHM, van Veenendaal NR, Ludolph RA, Ket JCF, van der Schoor SRD, van Kempen A. Effects of parent-provider communication during infant hospitalization in the NICU on parents: a systematic review with meta-synthesis and narrative synthesis. *Patient Educ Couns*. 2021;104(7):1526-52. doi:10.1016/j.pec.2021.04.023.
193. Bedwell C, Lavender T, Tate N, Danna VA. Interventions to support parents, families and carers in caring for premature or low birth weight (LBW) infants in the home: a systematic review and meta-analysis. *medRxiv*. 2022:2022.10.25.22281452v1. doi:10.1101/2022.10.25.22281452.
194. Franck LS, McNulty A, Alderdice F. The perinatal-neonatal care journey for parents of preterm infants: what is working and what can be improved. *J Perinat Neonatal Nurs*. 2017;31(3):244-55. doi:10.1097/JPN.0000000000000273.
195. Black RE, Taylor CE, Arole S, Bang A, Bhutta ZA, Chowdhury AMR, et al. Comprehensive review of the evidence regarding the effectiveness of community-based primary health care in improving maternal, neonatal and child health: 8. summary and recommendations of the Expert Panel. *J Glob Health*. 2017;7(1):010908. doi:10.7189/jogh.07.010908.
196. Vigod SN, Villegas L, Dennis CL, Ross LE. Prevalence and risk factors for postpartum depression among women with preterm and low-birth-weight infants: a systematic review. *BJOG*. 2010;117(5):540-50. doi:10.1111/j.1471-0528.2009.02493.x.
197. Koenraads M, Phuka J, Maleta K, Theobald S, Gladstone M. Understanding the challenges to caring for low birthweight babies in rural southern Malawi: a qualitative study exploring caregiver and health worker perceptions and experiences. *BMJ Glob Health*. 2017;2(3):e000301. doi:10.1136/bmjgh-2017-000301.
198. Sexual, reproductive, maternal, newborn, child and adolescent health: policy survey, 2018–2019: summary report. Geneva: World Health Organization; 2020 (<https://apps.who.int/iris/handle/10665/331847>).
199. TRAVAIL: Conditions of Work and Employment Programme [website]. International Labour Organization; 2017 (https://www.ilo.org/dyn/travail/travmain.sectionChoice?p_structure=, accessed 11 April 2022).

200. Koslowski A, Blum S, Dobroćić I, Kaufman G, Moss P, editors. 17th international review of leave policies and related research 2021: research report. International Network on Leave Policies and Research; 2021 (https://ub-deposit.fernuni-hagen.de/receive/mir_mods_00001739).
201. Integrated management of childhood illness: management of the sick young infant aged up to 2 months: IMCI chart booklet. Geneva: World Health Organization; 2019 (<https://apps.who.int/iris/handle/10665/326448>).
202. Pocket book of hospital care for children: guidelines for the management of common childhood illnesses, second edition. Geneva: World Health Organization; 2013 (<https://apps.who.int/iris/handle/10665/81170>).
203. Every Newborn Action Plan [website]. Geneva: World Health Organization; 2022 (<https://www.who.int/initiatives/every-newborn-action-plan>, accessed 1 January 2022).
204. Vogel JP, Dowswell T, Lewin S, Bonet M, Hampson L, Kellie F, et al. Developing and applying a “living guidelines” approach to WHO recommendations on maternal and perinatal health. *BMJ Global Health*. 2019;4(4):e001683. doi:10.1136/bmjgh-2019-001683.
205. Guidelines on basic newborn resuscitation. Geneva: World Health Organization; 2012 (<https://apps.who.int/iris/handle/10665/75157>).

Annexes

Annex 1: Current WHO recommendations for care of the preterm or low-birth-weight infant

	Recommendation
A. Preventive and promotive care	
Cord care	Included in WHO resuscitation guideline (see next page)
Kangaroo mother care	Included in this guideline
Thermal care	Included in WHO preterm guideline (see next page)
Feeding	Included in this guideline
Micronutrients	Included in this guideline
Probiotics	Included in this guideline
Emollients	Included in this guideline
Developmental care	Forthcoming in new WHO guidelines
Massage	Forthcoming in new WHO guidelines
Positioning	Forthcoming in new WHO guidelines
Immunization	Included in WHO immunization guideline (see next page)
Surveillance of growth, neurodevelopment, hearing, vision, disability	Forthcoming in new WHO guidelines
B. Care for complications	
Resuscitation	Forthcoming in update of WHO guidelines
Surfactant	Included in WHO preterm guideline (see next page)
Continuous positive airway pressure (CPAP)	Included in this guideline
Oxygen	Included in WHO preterm guideline (see next page)
Apnoea	Included in this guideline
Hypoglycaemia	Forthcoming in update of WHO guidelines
Hyperbilirubinaemia	Forthcoming in update of WHO guidelines
Infections	Forthcoming in update of WHO guidelines
Necrotizing enterocolitis	Forthcoming in update of WHO guidelines
Anaemia	Forthcoming in update of WHO guidelines
Growth, neurodevelopment, hearing, vision, disability	Forthcoming in new WHO guidelines
C. Family involvement and support	
Family involvement	Included in this guideline
Education and counselling	Included in this guideline
Discharge preparation	Included in this guideline
Peer support	Included in this guideline
Home visits	Included in this guideline
Parental leave and entitlements	Included in this guideline

Delayed cord clamping recommendations:¹

- In term or preterm newborns who do not require positive-pressure ventilation, the cord should not be clamped earlier than one minute after birth.
- When term or preterm newborns require positive-pressure ventilation, the cord should be clamped and cut to allow effective ventilation to be performed.

Thermal care recommendations:²

- During stabilization and transfer of preterm newborns to specialized neonatal care wards, wrapping in plastic bags or wraps may be considered as an alternative to prevent hypothermia.

Immunization recommendations:³

- Newborn immunization should be promoted in accordance with the latest existing WHO recommendations for routine immunization.

Surfactant recommendations:²

- Surfactant replacement therapy is recommended for intubated and ventilated newborns with respiratory distress syndrome.

- Either animal-derived or protein-containing synthetic surfactants can be used for surfactant replacement therapy in ventilated preterm newborns with respiratory distress syndrome.
- Administration of surfactant before the onset of respiratory distress syndrome (prophylactic administration) in preterm newborns is not recommended.
- In intubated preterm newborns with respiratory distress syndrome, surfactant should be administered early (within the first 2 hours after birth) rather than waiting for the symptoms to worsen before giving rescue therapy.

Oxygen recommendations:²

- During ventilation of preterm babies born at or before 32 weeks of gestation, it is recommended to start oxygen therapy with 30% oxygen or air (if blended oxygen is not available), rather than with 100% oxygen.
- The use of progressively higher concentrations of oxygen should only be considered for newborns undergoing oxygen therapy if their heart rate is less than 60 beats per minute after 30 seconds of adequate ventilation with 30% oxygen.

1 Guidelines on basic newborn resuscitation. Geneva: World Health Organization; 2012 (<https://apps.who.int/iris/handle/10665/75157>).

2 WHO recommendations on interventions to improve preterm birth outcomes. Geneva: World Health Organization; 2015 (<https://apps.who.int/iris/handle/10665/183037>).

3 WHO recommendations for routine immunization - summary tables [website]. Geneva: World Health Organization; 2021 (<https://www.who.int/teams/immunization-vaccines-and-biologicals/policies/who-recommendations-for-routine-immunization---summary-tables>, accessed 1 November 2022).

Annex 2: Contributors to the guideline

Members of the Guideline Development Group

Name	Affiliation
Nafisa Hamoud Al Jaifi	Sana'a University, Sanaa, Yemen
Shabina Ariff	Aga Khan University, Karachi, Pakistan
Mats Blennow	Karolinska Institutet and University Hospital, Stockholm, Sweden
Liz Comrie-Thomson	Burnet Institute, Melbourne, Australia
Gary Darmstadt (Co-chair)	Stanford University School of Medicine, Stanford, United States of America (USA)
Qi Feng	Peking University First Hospital, Beijing, China
Patricia Fernandez Riera	Ministry of Health, Buenos Aires, Argentina
Rizwan Ullah Hassan	Ministry of Public Health, Kabul, Afghanistan
Zelee Hill	University College London, London, United Kingdom of Great Britain and Northern Ireland
Ayede Adejumo Idowu	University of Ibadan, Ibadan, Nigeria
Carole Kenner	Council of International Neonatal Nurses, USA
Victoria Nakibuuka Kirabira	Nsambya Hospital, Kampala, Uganda
Reinhard Klinkott	Garmisch-Partenkirchen Hospital, Garmisch-Partenkirchen, Germany
Socorro De Leon-Mendoza	Kangaroo Mother Care Foundation, Philippines, Inc, Manila, Philippines
Silke Mader	European Foundation for the Care of Newborn Infants, Munich, Germany
Karim Manji	Muhimbili University of Health and Allied Sciences, Dar es Salaam, United Republic of Tanzania
Rhonda Marriott	Murdoch University, Perth, Australia
Monica Morgues	Universidad de Chile, Santiago, Chile
Sushma Nangia	Lady Hardinge Medical College & Kalawati Saran Children's Hospital, New Delhi, India
Vinod Paul (Co-chair)	National Institution for Transforming India (NITI Aayog), New Delhi, India
Mohammad Shaidullah	Bangabandhu Sheikh Mujib Medical University, Dhaka, Bangladesh
Hoang Thi Tran	Da Nang Hospital for Women and Children, Da Nang, Viet Nam
Andrew Weeks	University of Liverpool, Liverpool, United Kingdom
Bogale Worku	Ethiopian Pediatric Society, Addis Ababa, Ethiopia
Khalid Yunis	American University of Beirut, Beirut, Lebanon

Members of the Evidence Synthesis Team (EST)

Name	Affiliation
Rajiv Bahl	WHO Department of Maternal, Newborn, Child and Adolescent Health and Ageing (MCA)
Roger Chou (guideline methodologist)	Oregon Health and Science University, Portland, USA
Karen Edmond	WHO Department of MCA
Shuchita Gupta	WHO Department of MCA
Suman Rao	WHO Department of MCA
Natalie Strobel	Edith Cowan University, Perth, Australia

Members of the External Review Group (ERG)

Name	Affiliation
Ebunoluwa Adejuyigbe	Obafemi Awolowo University, Ife, Nigeria
Alexander Manu	Liverpool School of Tropical Medicine, Liverpool, United Kingdom
Carolyn McLennan	Northern Territory Department of Health, Alice Springs, Australia
Abiy Seifu	Addis Ababa University, Addis Ababa, Ethiopia

WHO Regional Office Representatives

Jamela Al-Raiby	WHO Regional Office for the Eastern Mediterranean
Rajesh Mehta	WHO Regional Office for South-East Asia
Assumpta Muriithi	WHO Regional Office for Africa
Khalid Siddeeg	WHO Regional Office for the Eastern Mediterranean
Howard Sobel	WHO Regional Office for the Western Pacific
Martin Weber	WHO Regional Office for Europe
Teshome Desta Woldehanna	WHO Regional Office for Africa

Members of the WHO Steering Group

Name	WHO Cluster/Department/Unit
Rajiv Bahl	Universal Health Coverage (UHC)/Life Course Cluster, Department of Maternal, Newborn, Child and Adolescent Health and Ageing, Newborn Health Unit (UHL/MCA/NBH)
Vanessa Cavallera	UHC/Communicable and Non-communicable Diseases Cluster, Department of Mental Health and Substance Use, Brain Health Unit (UCN/MSD/BRH)
Doris Chou	UHL, Department of Sexual and Reproductive Health and Research, Maternal and Perinatal Health Unit (UHL/SRH/MPH)
Karen Edmond*	UHL/MCA/NBH
Shuchita Gupta	UHL/MCA/NBH
Anayda Portela	UHL, MCA, Maternal and Adolescent Health Unit (UHL/MCA/MAH)
Suman Rao	UHL/MCA/NBH
Larry Grummer Strawn	Healthier Populations Cluster, Department of Nutrition and Food Safety, Food and Nutrition Actions in Health Systems Unit (HEP/NFS/AHS)

* Responsible technical officer

Annex 3: Summary of declarations of interest from the Guideline Development Group (GDG) members and how they were managed

Name	Declared interest(s)	Management of conflict(s) of interest
Nafisa Hamoud Al Jaifi	None declared	Not applicable
Shabina Ariff	None declared	Not applicable
Mats Blennow	He worked for Medecins Sans Frontiers (MSF) missions as a neonatal expert for a telemedicine project, with a monthly salary of US\$ 1000.	This declared conflict of interest was not considered significant enough to pose any risk to the guideline development process or to reduce its credibility.
Liz Comrie-Thomson	As a sole trader/consultant, she provided technical review of project design documents for an early child development project that included a nutrition component.	This declared conflict of interest was not considered significant enough to pose any risk to the guideline development process or to reduce its credibility.
Gary Darmstadt	None declared	Not applicable
Socorro De Leon-Mendoza	Attended the technical meeting on “Standards of care for Small and Sick Newborns”, 10–12 April 2019, at the Department of Maternal, Newborn, Child and Adolescent Health and Ageing, World Health Organization, Geneva, Switzerland.	This declared conflict of interest was not considered significant enough to pose any risk to the guideline development process or to reduce its credibility.
Qi Feng	None declared	Not applicable
Patricia Fernandez Riera	None declared	Not applicable
Zelee Hill	None declared	Not applicable
Ayede Adejumo Idowu	None declared	Not applicable
Carole Kenner	None declared	Not applicable
Victoria Nakibuuka Kirabira	None declared	Not applicable
Reinhard Klinkott	None declared	Not applicable
Silke Mader	None declared	Not applicable
Karim Manji	None declared	Not applicable
Rhonda Marriot	None declared	Not applicable
Monica Morgues	None declared	Not applicable
Sushma Nangia	None declared	Not applicable
Vinod Paul	None declared	Not applicable
Mohammod Shaidullah	None declared	Not applicable

Name	Declared interest(s)	Management of conflict(s) of interest
Hoang Tran	None declared	Not applicable
Rizwan Ullah Hassan	None declared	Not applicable
Andrew Weeks	He was a co-investigator on a grant to develop a low-cost neonatal resuscitation platform for low-resource settings. The total grant was 50 000 Canadian dollars from Grand Challenges Canada. It was paid to the Sanyu Research Institute in Uganda. He was an unpaid co-investigator and received no personal remuneration.	This declared conflict of interest was not considered significant enough to pose any risk to the guideline development process or to reduce its credibility.
Bogale Worku	None declared	Not applicable
Khalid Yunis	None declared	Not applicable

For more information, please contact:

World Health Organization
Avenue Appia 20, CH-1211 Geneva 27
Switzerland

Department of Maternal, Newborn, Child and Adolescent Health and Ageing
E-mail: mncah@who.int
Website: www.who.int/teams/maternal-newborn-child-adolescent-health-and-ageing/

ISBN 978-92-4-005826-2

